



**HAL**  
open science

## From a Sequential Chemo-Enzymatic Approach to a Continuous Process for HMF Production from Glucose

Alexandra Gimbernat, Marie Guehl, Nicolas Lopes Ferreira, Egon Heuson, Pascal Dhulster, Mickaël Capron, Franck Dumeignil, Damien Delcroix, Jean Girardon, Renato Froidevaux

► **To cite this version:**

Alexandra Gimbernat, Marie Guehl, Nicolas Lopes Ferreira, Egon Heuson, Pascal Dhulster, et al.. From a Sequential Chemo-Enzymatic Approach to a Continuous Process for HMF Production from Glucose. *Catalysts*, 2018, 8 (8), 10.3390/catal8080335 . hal-01915536v1

**HAL Id: hal-01915536**

**<https://ifp.hal.science/hal-01915536v1>**

Submitted on 8 Nov 2018 (v1), last revised 29 Jan 2019 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

1 Article

# 2 From a sequential chemo-enzymatic approach to a 3 continuous process for HMF production from glucose

4 Alexandra Gimbernat<sup>1</sup>, Marie Guehl<sup>2,3</sup>, Nicolas Lopes Ferreira<sup>4</sup>, Egon Heuson<sup>1</sup>, Pascal Dhulster<sup>1</sup>,  
5 Mickael Capron<sup>2</sup>, Franck Dumeignil<sup>2</sup>, Damien Delcroix<sup>3</sup>, Jean Sébastien Girardon<sup>2</sup> and Renato  
6 Froidevaux<sup>1\*</sup>

7 <sup>1</sup>Univ. Lille, INRA, ISA, Univ. Artois, Univ. Littoral Côte d'Opale, EA7394 –ICV- Institut Charles Violette,  
8 F-59000 Lille, France.

9 <sup>2</sup>Univ. Lille, CNRS, Centrale Lille, ENSCL, Univ. Artois, UMR 8181- UCCS- Unité de Catalyse et Chimie du  
10 Solide, F-59000, Lille, France.

11 <sup>3</sup>IFP Energies nouvelles, Rond-point de l'échangeur de Solaize, BP 3, 69360 Solaize, France

12 <sup>4</sup>IFP Energies nouvelles, 1 et 4 avenue de Bois-Préau, 92852 Rueil Malmaison, France

13  
14 \*Correspondence: Renato.froidevaux@univ-lille1.fr; Tel.: +33 320 41 75 66

15 **Abstract:** Notably available from the cellulose contained in lignocellulosic biomass, glucose is a  
16 highly attractive substrate for eco-efficient processes towards high-value chemicals. A recent  
17 strategy for biomass valorization consists on combining biocatalysis and chemocatalysis to realise  
18 the so-called chemo-enzymatic or hybrid catalysis. Optimisation of the glucose conversion to  
19 5-hydroxymethylfurfural (HMF) is the object of many research efforts. HMF can be produced by  
20 chemo-catalyzed fructose dehydration, while fructose can be selectively obtained from enzymatic  
21 glucose isomerization. Despite recent advances in HMF production, a fully integrated efficient  
22 process remains to be demonstrated. Our innovative approach consists on a continuous process  
23 involving enzymatic glucose isomerization, selective arylboronic-acid mediated fructose  
24 complexation/transportation and chemical fructose dehydration to HMF. We designed a novel  
25 reactor based on two aqueous phases dynamically connected via an organic liquid membrane,  
26 which enabled substantial enhancement of glucose conversion (70%) while avoiding intermediate  
27 separation steps. Further, in the as-combined steps, the use of an immobilized glucose isomerase  
28 and an acidic resin facilitates catalyst recycling.

29 **Keywords:** (bio)catalysis; Biomass; Glucose; HMF; chemo-enzymatic catalysis  
30

## 31 1. Introduction

32 In a context of fast depletion of fossil resources, lignocellulosic biomass possesses a high potential as  
33 a sustainable raw material for fuels [1] and fine chemicals [2, 3] production in industrially significant  
34 volumes. Upon specific physical, chemical and/or biological treatments of lignocellulose, access can  
35 be given to monomeric sugars [4], like glucose as its main saccharidic component. Glucose is, as a  
36 result, currently considered as one of the most promising starting materials for the conversion of  
37 biomass into drop-in or new high-value platform chemicals [5].

38 Efficient and eco-responsible valorization routes for this new renewable carbon source are highly  
39 desirable, and every field of catalysis should be mobilized as the key to success for a selective  
40 valorization of such a polyfunctional substrate. Biocatalytic processes are already or about to be  
41 industrialized [6] and heterogeneous or homogeneous catalysis, alone or in combination, have  
42 shown high efficiency for biomass valorization, but lack of selectivity still causes problems in the  
43 products separation steps [7]. In the very last years, a new paradigm of catalysis has emerged to


44 answer the challenges of selectivity and productivity by combining biocatalysis and chemocatalysis  
45 [8-10]. This combination is known as chemo-enzymatic catalysis or hybrid catalysis [11, 12].

46 In the field of sustainable and biosourced chemistry, glucose conversion to 5-hydroxymethylfurfural  
47 (HMF) drives a specific attention as a platform molecule [13]. Its hydrogenation, for instance, can  
48 lead to 2,5-dimethylfuran (2,5-DMF), a new fuel additive [14] and a precursor of terephthalic acid  
49 after dehydrative Diels-Alder reaction with ethylene to p-xylene and subsequent oxidation [15, 16].  
50 HMF oxidation is also envisioned to obtain 2,5-furandicarboxylic acid (FDCA), a biosourced  
51 alternative to terephthalic acid in the polyesters plastic production with a high market potential [17,  
52 18]. Unfortunately, abundant glucose is not the substrate of choice for efficient dehydration to HMF  
53 compared to its more expensive fructose isomer. The latter presents a fructofuranose mutamer more  
54 prone to dehydration to a furanic ring with lower energetic barriers [19]. Isomerization of glucose to  
55 fructose thus appears as a key step that can hardly be circumvented. A well-known transformation  
56 of glucose to fructose is the enzymatic glucose isomerization, employed in the industrial production  
57 of high-fructose corn syrup "HFCS" [20]. Despite the need for expensive high purity glucose, for  
58 buffers and multiple ion-exchange resins to get rid of all the metallic residues in alimentary HFCS,  
59 enzymatic isomerization remains the preferred process compared to chemical isomerization, even if  
60 the latter receives a recent renewed interest [21-25]. The aforementioned enzymatic reaction reaches  
61 a thermodynamic equilibrium ( $K_{eq} \sim 1$ ), which limits glucose conversion [26-28]. Whereas this  
62 equilibrium is particularly well-adapted for HFCS production involving glucose/fructose mixtures,  
63 HMF synthesis requires a pure fructose feed for the subsequent dehydration step, where  
64 compatibility issues between catalysts arise: Huang & al indeed shown incompatibility between a  
65 thermophilic glucose isomerase immobilized on aminopropyl-functionalized mesoporous silica  
66 (FMS) and a heterogeneous Brønsted acid propylsulfonic-FMS-SO<sub>3</sub>H in a THF:H<sub>2</sub>O (4:1 v/v) mixture  
67 [29]. Isomerization is carried out at 363 K to reach 61% fructose yield, and then the temperature has  
68 to be increased to 403 K to reach a 30% HMF yield. In the meantime, glucose isomerase is fully  
69 denatured.

70 Strategies have thus been set up to combine enzymatic isomerization of glucose to fructose and  
71 chemical by separating bio- and chemo-catalysis involving fructose transportation between the  
72 isomerization aqueous phase and an organic phase for subsequent reactivity. Huang *et al* [30]  
73 describe this concept for the first time by adding sodium tetraborate in the aqueous isomerization  
74 medium to form a fructoborate compound by complexation with fructose. Transportation of this  
75 complex to a separated organic phase is assisted by a cationic quaternary ammonium and enables  
76 enhanced glucose conversion by shifting the isomerization equilibrium towards fructose. The  
77 separate dehydration of fructose in the organic phase gives an increase in HMF yield up to 63%  
78 compared to a yield of 28% HMF without borate addition, associated to a glucose conversion of 88%  
79 instead of 53%. However, in this work, the complexation selectivity between glucose and fructose  
80 was not satisfactory and was thus further improved by Palkovits & al, who optimized the chemical  
81 nature of the complexing boronate species [31]. A global process exploiting this concept for glucose  
82 to HMF production has concomitantly been proposed by Alipour *et al* [32] Therein, fructose is  
83 complexed with phenylboronic acid and transferred to an organic phase, which is separated and  
84 contacted with an acidic ionic liquid phase to promote the release of free fructose. This fructose-rich  
85 ionic liquid is further used for dehydration to HMF in a biphasic medium. The produced HMF is  
86 then back-extracted to a last low-boiling point organic phase. This method involves the use of four  
87 different media with intermediate phase separations. Inspired by these sequential processes, we  
88 propose here to circumvent this drawback by setting up an innovative integrated implementation  
89 minimizing the number of steps and preventing among others the use of costly and hard to recover  
90 ionic liquids.

91 Herein, we highlight an efficient and fully-integrated cascade reactions process to convert glucose to  
92 HMF using an organic liquid membrane separating two aqueous phases for fructose transportation.  
93 This approach that we first proposed as a concept exemplifying the potential of hybrid catalysis [33]


94 is based on continuous aqueous glucose isomerization, fructose complexation, extraction and  
 95 transportation towards an intermediate organic phase. Fructose is then released at the interface with  
 96 the second aqueous phase (aqueous receiving phase), where it is subsequently dehydrated to HMF.  
 97 This integrated process without intermediate phase recovery, separation and recycling is presented  
 98 in Figure 1. Before fully designing the entire process, where all the reactions are carried out  
 99 simultaneously, we separately investigated each reaction to notably determine their respective  
 100 optimal ranges of conditions before combination: (i) Glucose isomerization to fructose occurs in a  
 101 primary aqueous feed phase; (2) The as-formed fructose transportation is then made possible  
 102 through selective complexation with phenylboronic acid as a carrier associated to a quaternary  
 103 ammonium into an organic solvent (4-methyl-2-pentanone “MIBK”); (3) Eventually, the  
 104 fructoboronate complex is hydrolyzed by acidic conditions in a receiving aqueous phase containing  
 105 an acidic resin, which also promotes dehydration of fructose to HMF. Our approach is not only  
 106 focused on the compatibility issues of bio- and chemocatalytic reactions with the aim of overcoming  
 107 the isomerization equilibrium limitation, but also pays a particular attention in integrating all the  
 108 steps to minimize separation and recycling burdens, which can be detrimental for the overall  
 109 economics and efficiency of the process. The methodology envisioned to move from a sequential  
 110 approach towards an integrated continuous process is schematized in Figure 2.



111

112 **Figure 1.** Hybrid catalysis simultaneous process applied to the transformation of glucose to HMF  
 113 (D-Glc = D-glucose, D-Fru = D-Fructose, HMF = 5-hydroxymethylfurfural)


114 **Figure 2.** Methodology from a sequential approach towards an integrated continuous process.



115

116 **2. Results and Discussion**117 *2.1. Study of D-glucose enzymatic isomerization in the aqueous feed phase*

118 The optimal temperature of use ranges from 333 to 353 K and the optimum pH ranges from pH 7.0 to  
 119 9.0 [34]. We first checked the influence of pH and temperature using the lot of IGI received from our  
 120 supplier. A glucose solution at pH 7.5 (Tris-HCl buffer, 100 mM) was prepared and isomerized from  
 121 323 K to 363 K. The relative enzymatic activity is presented in Figure 3, which highlights an optimal  
 122 enzymatic activity at 343 K. A glucose solution of a pH from 4.5 to 9 (Tris-HCl buffer, 100 mM from  
 123 pH 7 to 9; sodium phosphate buffer, 100 mM pH 6.5; sodium citrate buffer, 100 mM pH 4.5) was thus  
 124 prepared and isomerized at 343 K. The relative enzymatic activity presented in Figure 4 shows an  
 125 optimal IGI activity at a pH around 7.5. It is noteworthy that the IGI maintained more than 80% of its  
 126 optimal activity between pH 7 and 9. Below a pH of 6.5, the IGI lost more than 50% of its activity.  
 127 The selected pH in the aqueous feed phase for the enzymatic isomerization reaction was 8.5. A  
 128 D-glucose enzymatic isomerization was then performed in the selected conditions (Tris-HCl 100  
 129 mM, pH 8.5, 343 K). The results are presented in Figure 5 as a function of time. A blank reaction  
 130 (grey dots) was ran in the absence of IGI and showed the absence of D-glucose isomerization or  
 131 degradation in the chosen experimental conditions. When IGI was present in the system, the  
 132 D-fructose amount increased from 0% to 42% during the first 30 min. From 30 to 60 min, the  
 133 D-glucose conversion was slower, while the D-Fructose amount increased from 42% to 55%. A  
 134 maximum D-glucose conversion of 55% was finally reached in agreement with the glucose-fructose  
 135 thermodynamic equilibrium expected at this temperature ( $K_{eq} = 1,23$ ), as already observed by


136 McKay *et al* [35].

137 **Figure 3.** Relative enzymatic activity of IGI in D-glucose isomerization at pH 7.5 as a function of  
 138 temperature.


139 **Figure 4.** Relative enzymatic activity of IGI in D-glucose isomerization at 343 K as a function of pH

140

141 **Figure 5.** D-glucose consumption during the isomerization step as a function of time; ( $\Delta$ ), D-fructose




142 formation ( $\square$ ), D-glucose amount in the blank experiment at pH 8.5 ( $\circ$ ). Initial conditions: D-Glucose



143 (100 mM), IGI (0,5 g), Tris-HCl 100 mM, pH 8.5, V = 100 mL, 343 K, 750 rpm.

#### 144 2.2. Organic liquid membrane for D-fructose complexation/transportation

145 In order to enhance the isomerization yield, simultaneous isomerization and  
 146 complexation/transportation of D-fructose in a liquid/liquid aqueous/organic biphasic system was  
 147 further studied. The temperature was set at 343 K and the pH was chosen as a trade-off to  
 148 simultaneously maximize the enzymatic activity ( $7 < \text{pH} < 9$ ), to avoid monosaccharide degradation,  
 149 and to enhance the extraction process as reported previously [36]: the pH was thus kept at a value of  
 150 8.5 by using a Tris-HCl 100 mM buffer. The extraction was conducted in an organic solvent with low  
 151 water miscibility (MIBK) containing a lipophilic arylboronic acid (carrier) and a quaternary  
 152 ammonium salt (Aliquat336<sup>®</sup>) as a phase transfer agent [37]. At pH 8.5, aryl boronic acid  $\text{ArB}(\text{OH})_2$   
 153 was actually present under its hydroxylated anionic form, as a tetrahedral aryltrihydroxyborate  
 154  $\text{ArB}(\text{OH})_3^-$  [38]. At the interface between the aqueous and organic phases, D-fructose further reacts  
 155 with the arylborate to form a tetrahedral fructoboronate ester (Figure 6) [38]. The fructoboronate  
 156 complex then forms an intimate ion pair with Aliquat336<sup>®</sup>, which enables its transportation to the  
 157 organic phase [39].



158 **Figure 6.** Fructoboronate complex formation at the interphase between the first aqueous phase and  
 159 the organic liquid membrane.

160

#### 161 2.2.1. Influence of the boronic acid structure

162 The influence of the boronic acid structure was investigated in order to optimize kinetics and  
 163 maximize selectivity of D-fructose complexation/transportation. Seven arylboronic acids differing  
 164 by the electronic properties of their substituents and thus by their pKa were screened in the  
 165 complexation reaction with D-fructose: 2,3-DCPBA, 2,4-DCPBA, 3,4-DCPBA, 3,5-DCPBA, PBA,  
 166 4-TBPBA, 4-TFMPBA. The complexation/transportation, which will be referred to as “extraction” in  
 167 the following sections, was carried out in a biphasic system, as described before. The pKa of the  
 168 different boronic acids and the relative extraction yields are summarized in Table 1 and reported in  
 169 Figure 7.

170

171 **Table 1.** Influence of the boronic acid structure on the extraction yield and rate.

Boronic acid	$pK_a$	Extraction		Extraction	
		yield		rate	
		%		$\mu\text{mol}/\text{min}$	
4-TBPBA	9.3	8.3	$\pm 5.6$	0.52	$\pm 0.075$
PBA	9.1	32.4	$\pm 0.3$	1.27	$\pm 0.396$
2,4-DCPBA	8.9	43.3	$\pm 1.6$	1.56	$\pm 0.34$
3,4-DCPBA	7.4	46.5	$\pm 4.9$	1.48	$\pm 0.23$
2,3-DCPBA	7.4	49.2	$\pm 1.6$	1.99	$\pm 0.178$
4-TFMPBA	9.1	50.3	$\pm 2.2$	1.26	$\pm 0.129$
3,5-DCPBA	7.4	55.3	$\pm 0.9$	1.94	$\pm 0.280$

181


182

183

184

185

186



187

188 **Figure 7.** Influence of the arylboronic acid structure on the complexation/transportation process  
 189 after 45 min. [D-Fructose]<sub>I</sub> = 100 mM, [Boronic acid] = 100 mM, [Aliquat336®] = 200 mM, Tris HCl 100  
 190 mM pH 8.5, MIBK, 343 K, 750 rpm.


191

192 A group composed by 2,4-DCPBA, 3,4-DCPBA, 2,3-DCPBA, 4TFMPBA and 3,5-DCPBA showed the  
 193 best results with extraction yields between 43% and 55%, while with 4-TBPBA and PBA only 8% and  
 194 32% were achieved, respectively. These results also show that the D-fructose extraction yield  
 195 increases when the pKa of the boronic acid used decreases. Better extraction yields were observed  
 196 when the pKa of the boronic acid was lower than (or close to) the pH of the aqueous feed phase (8.5).  
 197 This analysis supports the hypothesis from Morin *et al.* [36] and Karpa *et al* [37] indicating that  
 198 higher extraction yields are obtained when  $\text{pH} \geq \text{pKa}$ , favoring a predominant tetrahedral anionic  
 199 borate form of the complex, which enables its transportation by association with the ammonium  
 200 carrier. An exception was however observed with 4-TFMPBA with a pKa of 9.1. This might be due to  
 201 the fact that 4-TFMPBA has a higher dipolar moment (2.90 D, against 0.43 D to 1.86 D for the other  
 202 used boronic acids) that could increase the reactivity. The extraction rates and the pKas (Table 1)  
 203 seem to be correlated. A higher pKa of the boronic acid was linked with a lower initial extraction  
 204 rate. Among the most effective boronic acids, 3,4-DCPBA was selected for further studies.

205

### 206 2.2.2. Influence of the boronic acid:Aliquat336® molar ratio

207 With 3,4-DCPBA as the selected boronic acid, the influence of the molar ratio between the boronic  
 208 acid and the ammonium salt Aliquat336® in the organic solvent was investigated. Figure 8 shows the  
 209 evolution of D-Fructose concentration in the aqueous phase as a function of time for different  
 210 Aliquat336® concentrations (from 0 to 400 mM). D-Fructose and 3,4-DCPBA were initially  
 211 introduced at concentrations of 100 mM. In the first 15 minutes of reaction, the concentration of  
 212 D-Fructose decreased irrespective of the Aliquat336® concentration with various initial rates. An  
 213 equilibrium seemed to be progressively reached between 15 and 45 minutes, as no more evolution of  
 214 the D-Fructose concentration was then observed. A blank reaction (blue dots) without Aliquat336®  
 215 nor 3,4-DCPBA was also tested, which showed no evolution. From these observations, it can be  
 216 considered that the molar ratio 3,4-DCPBA:Aliquat336® is a relevant parameter to control the initial  
 217 D-fructose extraction rate. Figure 9 shows the evolution of the initial extraction rate and of the  
 218 D-fructose extraction yield for different 3,4-DCPBA:Aliquat336® molar ratios (from 1:0.5 to 1:4). The  
 219 initial extraction rate continuously increased when the 3,4-DCPBA:Aliquat336® molar ratio  
 220 increased from 1:0.5 to 1:2 with respectively 0.37  $\mu\text{mol}/\text{min}$  and 1.50  $\mu\text{mol}/\text{min}$ . Above a 1:2 molar  
 221 ratio, no further kinetic benefits could be clearly observed from such an excess of Aliquat336® in the  
 222 reaction medium. This result is consistent with an ion-pairing phenomenon between the  
 223 fructoboronate complex and the Aliquat336® occurring at the interphase enabling D-fructose  
 224 transportation. The D-fructose extraction yield increased when the 3,4-DCPBA:Aliquat336® molar  
 225 ratio increased from 1:0.5 to 1:2 with respectively 21.5% and 50.0%. Like in the case of the initial  
 226 extraction rate, above a 1:2 molar ratio no further benefits on the extraction yield were displayed by  
 227 using an excess of Aliquat336®. Therefore, in order to maximize the initial extraction yield and the  
 228 initial rate while optimizing the quantity of ammonium salt added, an arylboronic acid:Aliquat336®  
 229 ratio of 1:2 was selected for the following studies.






230 **Figure 8.** Evolution of [D-fructose] in the aqueous phase as a function of time for different  
 231 3,4-DCPBA/Aliquat336® ratios. [D-fructose]<sub>i</sub> = 100 mM, [3,4-DCPBA] = 100 mM, [Aliquat336®] = X  
 232 mM (X varies from 0 to 400 mM), Tris-HCl 100 mM pH 8.5, MIBK, 343 K, 750 rpm.

233

234

235



236 **Figure 9.** Initial extraction rate in aqueous phase (blue rods) and extraction yield (red dots) in  
 237 function of 3,4-DCPBA:Aliquat336® molar ratio. [D-Fructose]<sub>i</sub> = 100 mM, [3,4-DCPBA] = 100 mM,  
 238 [Aliquat336®] = X mM (X varies from 0 to 400 mM), Tris-HCl

### 239 2.2.3. Influence of boronic acid and Aliquat336® concentrations

240 Keeping a molar ratio of arylboronic acid:Aliquat336® equal to 1:2 and at fixed initial D-Fructose  
 241 amount, both 3,4-DCPBA and Aliquat336® concentrations were varied, considering the best  
 242 compromise between component concentrations and costs (Figure 10). From 0 to 15 min, the  
 243 D-Fructose extraction yield increases rapidly and then stabilizes for lowest 3,4-DCPBA  
 244 concentrations less than 100 mM. The best extraction yields are thus obtained for higher 3,4-DCPBA  
 245 concentrations, the final yield being the highest at 300 mM. The slight difference in extraction yields  
 246 between 100 mM and 300 mM 3,4-DCPBA concentration does not justify the use of such a  
 247 concentrated solution. From Figure 11, the initial extraction rate values for different  
 248 3,4-DCPBA:Aliquat336® concentrations have been calculated (Table 2). Two behaviors can be  
 249 observed, the first one for the 25:50 ratio and the second one for all the other ratios, namely from  
 250 50:100 to 300:600. We observed that the larger the 3,4-DCPBA amount, the higher the initial  
 251 extraction rate. However, after a certain amount of 3,4-DCPBA, the interface seems to be saturated  
 252 by 3,4-DCPBA:Aliquat336® complexes and then the initial rate does not increase anymore.  
 253 Therefore, further optimization steps considered a concentration of 100 mM while keeping a molar  
 254 ratio of 1:2 with Aliquat 336®.

255

256


257

258

259

260 **Table 2.** Initial extraction rates for various concentrations in the system. [D-Fructose]<sub>I</sub> = 100 mM,  
 261 [3,4-DCPBA] = Y mM (Y varies from 25 to 300 mM), [Aliquat336<sup>®</sup>] = Y×2 mM, Tris-HCl 100 mM, pH  
 262 8.5, MIBK, 343 K, 750 rpm.

[3,4-DCPBA] (mM)	Extraction rate ( $\mu\text{mol}/\text{min}$ )
25	0.89 $\pm$ 0.15
50	1.37 $\pm$ 0.15
150	1.45 $\pm$ 0.05
100	1.22 $\pm$ 0.15
200	1.40 $\pm$ 0.21
300	1.33 $\pm$ 0.14




268

269 **Figure 10.** Evolution of the D-Fructose extraction yield for different 3,4-DCPBA and Aliquat336<sup>®</sup>  
 270 concentrations for a 1:2 molar ratio. [D-Fructose]<sub>I</sub> = 100 mM, [3,4-DCPBA] = Y mM (Y varies from 25  
 271 to 300 mM), [Aliquat336<sup>®</sup>] = Y × 2 mM, Tris-HCl 100 mM, pH 8.5, MIBK, 343 K, 750 rpm.

272 2.2.4. Influence of the D-fructose:boronic acid molar ratio

273 The influence of the D-Fructose:3,4-DCPBA molar ratio was further investigated. Figure 11 shows  
 274 the D-fructose extraction yield as a function of this molar ratio. The yield ranged between 40% and  
 275 60%, for ratios from 0.25:1 to 1:1 and between 18% and 28% for ratios from 2:1 to 10:1. When the  
 276 3,4-DCPBA was introduced in excess or stoichiometrically compared to D-Fructose (red rods), the  
 277 extraction yield was larger than 40%, reaching a maximum of 60% for a two-fold excess of boronic

278 acid. However, when it was introduced in default (blue rods), the extraction yield did not exceed  
 279 35% without any significant further evolution between a molar ratio of 2:1 to 10:1.




280

281 **Figure 11.** Influence of the initial D-Fructose:3,4-DCPBA molar ratio on the the extraction yield.  
 282 [D-Fructose]<sub>i</sub> = X mM, [3,4-DCPBA] = 100 mM, [Aliquat336<sup>®</sup>] = 200 mM, Tris-HCl 100 mM, pH 8.5,  
 283 MIBK, 343 K, 750 rpm.

284 2.2.5. Influence of D-fructose and boronic acid concentrations

285 The evolution of the extracted D-fructose concentration as a function of the initial D-fructose  
 286 concentration, for the same 3,4-DCPBA concentration, is shown in Figure 12. Logically, when the  
 287 initial concentration of D-fructose increased from 25 mM to 1000 mM, the extracted concentration  
 288 linearly increased, according to a straight correlation between the initial amount of D-fructose and  
 289 the amount of extracted D-fructose. Keeping a molar D-fructose:boronic acid ratio equal to 1:1 and a  
 290 molar ratio of arylboronic acid:Aliquat336<sup>®</sup> equal to 1:2, a benchmark experiment was set with the  
 291 following parameters: T = 34 K, [D-Fructose]<sub>i</sub> = 100 mM, [3,4-DCPBA] = 100 mM, [Aliquat 336<sup>®</sup>] = 200  
 292 mM. The other experiments have been set up by doubling all concentrations. Table 3 presents the  
 293 extraction yields and initial extraction rates depending on the variation of the initial D-fructose  
 294 concentration while keeping all the molar ratios relative to 3,4-DCPBA and Aliquat336<sup>®</sup> as constant.  
 295 An increase in the concentrations (D-Fructose, 3,4-DCPBA and Aliquat336<sup>®</sup>) led to a decrease in the  
 296 D-Fructose extraction yield. Indeed, at 100 mM of D-Fructose, 43.3% of the fructoboronate complex  
 297 were extracted whereas only 32.7% and 27.3% were respectively extracted at 200 mM and 300 mM of  
 298 D-fructose. Therefore, the optimal conditions ([D-Fru]/[3,4-DCPBA]/[Aliquat336] ratio of  
 299 100/100/200 mM) were kept for the following studies.

300



301 **Figure 12.** Influence of the initial D-Fructose concentration on the amount of D-fructose extracted.  
 302 [D-Fructose]<sub>I</sub> = X mM (X varies from 25 to 1000), [3,4-DCPBA] = 100 mM, [Aliquat336®] = 200 mM,  
 303 Tris-HCl 100 mM, pH 8.5, MIBK, 343 K, 750 rpm.

304 **Table 3.** D-fructose extraction yield and initial extraction rate for different concentrations.


[D-Fru]	[3,4-DCPBA]	[Aliquat336®]	Extraction yield	
mM	mM	mM	%	
100	100	200	43.34	± 4.22
200	200	400	32.61	± 1.22
300	300	600	27.31	± 4.6

305

306 2.2.6. Hydrolysis of the fructoboronate complex for D-fructose release in the second aqueous phase

307 The further stage of the D-fructose transportation is the fructoboronate complex hydrolysis for  
 308 D-fructose release in the second aqueous phase (the aqueous receiving phase). Therefore, the  
 309 influence of the pH of the second aqueous phase on the release yield was investigated. pHs of 1, 3, 5  
 310 and 8 were tested by using with H<sub>2</sub>SO<sub>4</sub> for pH 1, citrate buffer for pH 3 and 5 and Tris HCl buffer for  
 311 pH 8. Prior to these experiments, a D-fructose extraction experiment was performed  
 312 complexation/transportation) in the optimized conditions (100 mM D-fructose and a  
 313 D-fructose:3,4-DCPBA:Aliquat336® molar ratio of 1:1:2). The organic phase was then recovered and  
 314 contacting with the second aqueous phase for the release step. As observed by Paugam *et al* [40] and  
 315 keeping in mind the final objective of this work that is the obtaining of a simultaneous process, the  
 316 saccharide flow from the basic aqueous donor phase to the acidic receiving phase through an  
 317 organic phase is favored when a pH gradient is applied between the two phases. Figure 13 shows  
 318 the evolution of the D-fructose release yield as a function of time for different pH of the aqueous  
 319 receiving phase. In the cases of pH 8 and pH 1, an increase was observed until 30 minutes (with an  
 320 initial rate of about 0.6 and 1.8 μmol/min, respectively, Table 4) and a steady-state was reached for a  
 321 yield of 22.7% and 54.5% for pH 8 and pH 1, respectively. Concerning pH 3 and pH 5, an increase  
 322 was observed up to 60 minutes (with an initial rate of 1.5 μmol/min, Figure 13) to obtain respectively  
 323 91% and 100% of release. Considering that the pK<sub>a</sub> of 3,4-DCPBA is around 7.4, when the pH of the  
 324 receiving aqueous phase is 8, the pH is then superior to pK<sub>a</sub>. Then, the release mechanism is  
 325 unfavorable [41]. With pH 3 and 5, more than 90% of the extracted D-Fructose were released.  
 326 Moreover, the experiment with H<sub>2</sub>SO<sub>4</sub> at pH 1 shows full hydrolysis of the fructoboronate complex  
 327 followed by conversion of D-Fructose to HMF, levulinic and formic acid, which explains the high  
 328 release rate and the decrease in the actual D-fructose concentration after 15 minutes.

329



330 **Figure 13.** Influence of the pH of the aqueous receiving phase on the D-fructose complex  
 331 transportation. Extraction process: Tris-HCl 100 mM pH 8.5, [D-Fru]I = 100 mM, [3,4-DCPBA] = 100  
 332 mM, [Aliquat336®] = 200 mM, MIBK, 343 K. Receiving phase: Citrate buffer

333 **Table 4.** Transport yield and extraction rate of D-fructoboronate hydrolysis to D-fructose and  
 334 D-fructose release in the aqueous phase as a function of its pH.

pH	Transport Yield (%)		Extraction rate ( $\mu\text{mol}/\text{min}$ )	
5	100	$\pm 4.2$	1.58	$\pm 0.14$
3	91.5	$\pm 6.3$	1.53	$\pm 0.13$
1	54.5	$\pm 3.7$	1.84	$\pm 0.09$
8	22.7	$\pm 5.2$	0.63	$\pm 0.14$

335

336

### 337 2.3. Study of D-Fructose dehydration in the receiving aqueous phase

338 D-Fructose dehydration to HMF was carried out in the receiving aqueous phase. We previously  
 339 observed when studying fructoboronate complex transportation and hydrolysis in a highly acidic  
 340 aqueous phase the formation of formic and levulinic acids. This confirmed that solutions to  
 341 minimize HMF degradation by rehydration in the aqueous phase are requested. An alternative to  
 342 strong homogeneous acid has been to use an acidic resin containing strong sulfonic groups, its  
 343 supported character facilitating a potential regenerability and recyclability of the catalyst.


344

### 345 2.4. Study of sequential process

346 In the formerly optimized conditions, a sequential process was tested. The first step consisted on a  
 347 simultaneous D-glucose isomerization and D-fructose transportation through a fructoboronate  
 348 complex, followed by the second step characterized by the hydrolysis of the fructoboronate complex  
 349 to D-fructose and release of D-fructose in the aqueous receiving phase, and finally by the  
 350 dehydration of fructose to HMF by heterogeneous catalyst in the third step (Figure 14).

#### 351 2.4.1. D-glucose isomerization and D-fructose complexation/transportation

352 The aqueous feed phase contained Tris-HCl buffer 100 mM at pH 8.5, D-Glucose and immobilized  
 353 glucose isomerase at the required amount. The organic layer contained the system determined by  
 354 the previous study: [3,4-DCPBA] = 100 mM, [Aliquat336®] = 200 mM. The initial aqueous D-glucose  
 355 concentration was 100 mM. The system was mixed in a reactor where the enzyme was placed in a  
 356 basket to avoid any contact with the organic layer. Moreover, the use of MIBK, with a low solubility  
 357 in water, as an organic solvent contributed to bring the contact between MIBK and enzyme  
 358 negligible. The system was heated at 343 K for 3 hours. The amount of D-glucose and D-fructose  
 359 were chromatographically determined in the aqueous phase during the process. Table 5 shows the  
 360 performances of each step. An interesting isomerization yield of 74.5% was obtained showing the  
 361 actual shifting of the equilibrium of the enzymatic catalysis (55%) by fructose elimination from the  
 362 media through complexation with the boronic acid. Moreover, according to the extraction yield of  
 363 D-fructose and D-glucose, 56.5% and 1%, respectively, a very high selectivity of the complexation  
 364 reaction with the chosen boronic acid towards the D-fructose was actually demonstrated. The partial  
 365 D-fructose extraction yield of 56.5% showed that the reaction conditions could be still improved.



366

367 **Figure 14.** Illustration of the sequential process with isomerization and complexation/transportation  
 368 step, hydrolysis/release step and dehydration step.

369

370 **Table 5.** Results obtained from sequential process. The D-Fructose released yield was calculated  
 371 using the amount of D-Fructose in the organic phase at the end of the 1st transport. The fructose  
 372 conversion yield was calculated using the amount of fructose in the aqueous receiving phase at the  
 373 end of the 2nd transport. The HMF total amount was calculated using the initial amount of glucose  
 374 in the aqueous donor phase.

	1 <sup>st</sup> step		2 <sup>nd</sup> step		3 <sup>rd</sup> step		
	isomerization	D-Fructose extracted	D-Glucose extracted	D-Fructose released	Fructose conversion	HMF yield	HMF total
%	74.5	56.5	1.56	57.4	52	21.9	5.3
time		3 h		3 h	35 h		41 h

375

376

#### 377 2.4.2. Fructoboronate complex hydrolysis, release of D-fructose and dehydration to HMF


378 The basket of IGI was removed and the aqueous feed phase was discarded and replaced by the  
 379 receiving aqueous phase with a citrate buffer at 100 mM keeping a pH 3. The system was heated at

380 343 K for 3 hours. In these conditions, the hydrolysis step led to 63% of fructose release in the  
 381 aqueous phase. Then, after 3 hours at 343 K, the organic layer was discarded and the acidic resin was  
 382 introduced. The system was heated at 363 K. The dehydration step led to a HMF yield of 20% after  
 383 10 h. The total HMF yield was 5%.

384

### 385 2.5. Study of continuous process

386 In the optimal conditions previously determined, a continuous integrated process was tested [33].  
 387 The reactor, phases and matter flows are presented in Figure 15. The "coaxial" reactor is composed  
 388 of two cylinders coaxially placed, a cover enabling introduction of the stirring system and a  
 389 refrigerant for temperature control. The inner cylinder contains the donor aqueous phase and has a  
 390 height less than that of the outer cylinder which contains the receiving aqueous phase, which allows  
 391 the organic phase to be in contact with both aqueous phases. The experiment was conducted with a  
 392 donor aqueous phase formed by 50 mL of 100 mM Tris-HCl pH 8.5 containing 100 mM of D-glucose,  
 393 200 mg IGI, 20 mM MgSO<sub>4</sub>, 8 mM Na<sub>2</sub>SO<sub>3</sub>. The receiving aqueous phase was formed by 55 mL of 100  
 394 mM sodium citrate buffer, pH 3. The organic phase consisted of 100 mL of MIBK containing 100 mM  
 395 of 3,4-DCPBA and 200 mM of Aliquat336®. All the phases were maintained at 343 K. The stirring was  
 396 carried out at 180 rpm, which is the maximum rate to avoid the mixing of the phases, thanks to a  
 397 stirring blade immersed in the organic phase. This "coaxial" reactor was used to simultaneously  
 398 perform the isomerization, extraction, hydrolysis and dehydration steps. The monitoring of the  
 399 D-fructose concentration in the aqueous phases was carried out, making it possible to calculate the  
 400 efficiency of the continuous process for the simultaneous isomerization, fructose extraction and  
 401 dehydration. The results are shown in Table 6.



402

403 **Figure 15.** Illustration of the continuous process.

404

405 **Table 6.** Results obtained by continuous process.

	Isomerization	Fructose extraction	HMF	Selectivity
Yield (%)	70.1	50.2	4.1	70.4

406

407

408 The D-glucose isomerization yield was 70%, with 50% of the formed D-fructose transported and a  
 409 global HMF yield of 4% (table 8). Those results are comparable to those obtained by the sequential  
 410 process. Therefore, this process demonstrates the efficiency of using a cascade chemo-enzymatic  
 411 catalysis to continuously obtain HMF from D-glucose. However, insufficient agitation of each phase  
 412 seems to be the main limitation of this reactor configuration. Further improvements of this proof of  
 413 concept towards enhanced HMF yields and global performances will be reported in due course.

### 414 3. Materials and Methods

#### 415 3.1. Chemicals and reagents

416 D-Glucose ( $\geq 99.5\%$ ), D-Fructose ( $\geq 99\%$ ), N-methyl-N,N,N-trioctylammonium chloride  
 417 (Aliquat336<sup>®</sup>), 3,4-dichlorophenylboronic acid (3,4-DCPBA), 3,5-dichlorophenylboronic acid  
 418 (3,5-DCPBA) (98%), 2,4-dichlorophenylboronic acid (2,4-DCPBA), 2,3-dichlorophenylboronic acid  
 419 (2,3-DCPBA), 4-tert-butylphenylboronic acid (4-TBPBA) ( $\geq 95\%$ ), 4-(trifluoromethyl)phenylboronic  
 420 acid (4-TFMPBA), Trizma base ( $\geq 99.9\%$ ), hydrochloric acid (36.8-38%), sulfuric acid (95-97%),  
 421 Sweetzyme IT<sup>®</sup>Extra (IGI) ( $\geq 350$  U/g), Dowex monosphere 650C, levulinic acid (LA) (98%), formic  
 422 acid (FA) (95%), trisodium citrate ( $\geq 99\%$ ), 5-hydroxymethylfurfural (HMF) (99%) were purchased  
 423 from Sigma Aldrich Co. (St. Louis, MO). Phenylboronic acid (PBA) (95%) and 4-methyl-2-pentanone  
 424 (MIBK) ( $\geq 99\%$ ) were purchased from Sigma-Fluka. Citric acid ( $\geq 99\%$ ) was purchased from Merck.  
 425 Distilled-deionized water (Mili-Q) grad was used whenever necessary, obtained from a MilliQ water  
 426 purification system (Millipore, Academic, SA, France).

427

#### 428 3.2. Characterization

429 In the conversion/complexation experiments, the determination of reactants, intermediates and  
 430 products quantities (Table 7) was carried out using an HPLC Shimadzu equipped with a  
 431 LC-20ADXR pump, a DGU-20A5R degasser, a SIL-20ACXR autosampler, a SPD-MD20A diode array  
 432 detector, a CTO-20AC oven and a CBM-20A communicator module piloted by a LabSolution  
 433 software. The column was an Aminex HPX-87H (300 mm x 7.8 mm, 9  $\mu\text{m}$ ; Bio-Rad). The mobile  
 434 phase was 5 mM H<sub>2</sub>SO<sub>4</sub>, using isocratic mode with 0.4 mL.min<sup>-1</sup>. The chosen wavelength for HMF  
 435 and levulinic acid detection were 284 nm and 266 nm, respectively. The chosen wavelength for  
 436 detection of formic acid, D-Glucose and D-Fructose was 195 nm. In the case of monosaccharides, a  
 437 Refractive Index Detector was also used (analytical standards are listed in Table 7).

438

439 **Table 7.** Concentration range of products used for analytical standards.

Analytical method	RID				UV 284 nm	UV 195 nm		
	HMF	D-Glc	D-Fru	LA	HMF	D-Glc	D-Fru	FA
Concentration range (mM)	0.14-30	0.5-25	0.5-25	0.5-25	0.007-0.5	0.5-25	0.5-25	0.5-35

440

441

#### 442 3.3. Aqueous phase D-glucose isomerization (aqueous feed phase) and D-fructose complexation/transportation

##### 443 3.3.1. D-glucose isomerization in aqueous feed phase

444 We used one of commercial enzymes the most largely used for the conversion of starches to high  
 445 fructose syrup, namely immobilized glucose isomerase (IGI, Sweetzyme<sup>®</sup>IT extra). 31 1 mL of a  
 446 D-glucose solution was prepared by mixing D-glucose (100 mM), Na<sub>2</sub>SO<sub>3</sub> (8 mM) and MgSO<sub>4</sub> (20  
 447 mM) in a buffer solution (100 mM) at a selected pH (Tris-HCl buffer for pH 7, 7.5, 8, 8.5, 9, phosphate  
 448 buffer for pH 6.5, or citrate buffer for pH 4.5) and then introduced in an Eppendorf tube. The  
 449 solution was incubated at the desired temperature (323 to 363 K) in a Mixing Block (BIOER-102



450 Thermocell) at 900 rpm. After rehydration in water for 12 h, IGI (10 mg) was put in contact with the  
451 aforementioned prepared D-glucose solution. The reaction time was set at 90 min.

### 452 3.3.2. D-fructose transport in the organic phase

453 First, a D-fructose/D-glucose (25 to 600 mM) solution was prepared in Tris-HCl buffer (100 mM). 600  
454  $\mu\text{L}$  of this solution were introduced in an Eppendorf tube (solution 1). Aliquat336® (25-300 mM), and  
455 arylboronic acid (25-300 mM) were dissolved in MIBK as a solvent to prepare solution 2. 600  $\mu\text{L}$  of  
456 solution 2 were then introduced in a second Eppendorf tube. Solutions 1 and 2 were incubated at the  
457 desired temperature in a Mixing Block (BIOER-102 Thermocell) at 750 rpm. After 5 min, both  
458 solutions were mixed in an Eppendorf tube placed in the Mixing Block in the same conditions. After  
459 D-fructose transport, the aqueous and organic phases were separated by pipetting.

460

461

### 462 3.3.3. Aqueous phase D-fructose complex hydrolysis in receiving aqueous phase

463 600  $\mu\text{L}$  of citrate buffer were introduced in an Eppendorf tube and incubated at the required  
464 temperature (343 K) in a Mixing Block at 750 rpm. After 5 min, 600  $\mu\text{L}$  of the organic phase obtained  
465 after the extraction procedure and containing the fructoboronate complex were introduced in the  
466 tube. After a reaction time of 45 min, the aqueous and organic phases were separated by pipetting.

### 467 3.3.4. Study of sequential process from glucose isomerization to fructose dehydration

468 A 250 mL vessel (Reactor-Ready™ Lab Reactor, Radley, Figure 16) equipped with a temperature  
469 control system (Ministat, Huber temperature control system, HUBER, UK) was used for these  
470 experiments.

471 • Simultaneous D-glucose isomerization by enzymatic catalysis in the aqueous feed phase and  
472 D-fructose complexation/transportation in the organic phase

473 IGI (0.5 g) was loaded in a stirring basket introduced in a first reactor (Figure 16 – Left) with 100 mL  
474 of a aqueous solution containing D-Glucose (100 mM),  $\text{Na}_2\text{SO}_4$  (8 mM),  $\text{MgCl}_2$  (20 mM), in Tris-HCl  
475 buffer (100 mM, pH 8.5) at 343 K. Then, 100 mL of MIBK containing 3,4-DCPBA (100 mM) and  
476 Aliquat336® (200 mM) were introduced in the reactor. The mixture was stirred via the stirring basket  
477 at 200 rpm, with the temperature kept at 343 K for 180 min, before discarding the aqueous feed  
478 phase.

479 • D-fructose complex hydrolysis in the aqueous receiving phase

480 100 mL of the organic phase issued from the first reactor and containing the fructoboronate complex  
481 were transferred in a second reactor (Figure 16 – Middle) and heated at 343 K. Then, 100 mL of a  
482 citrate buffer solution (100 mM, pH 3) were introduced. The mixture was stirred with a turbine  
483 stirring shaft at 200 rpm and set up at 343 K for 180 min.

484

## 485 4. Conclusion

486 In this work, we have first provided the conditions of the key parameters (temperature, pH,  
487 arylboronic acid structure and concentrations, Aliquat336® concentrations) applied for a sequential  
488 process from glucose to HMF through fructose transportation in an organic liquid membrane. The

489 optimized conditions have been then applied to an unprecedented integrated process in a  
490 specifically designed reactor. Therein, enzymatic glucose isomerization and fructose dehydration to  
491 HMF, involving an intermediate fructose transportation by complexation with an aryboronic  
492 acid-Aliquat336® complex, could be carried out simultaneously in independent aqueous phases.  
493 These phases were both in contact with a single organic phase enabling the transportation of fructose  
494 from the first aqueous phase to the second. In this process, we thus succeeded in simultaneously  
495 enzymatically isomerizing glucose to fructose and dehydrating fructose to HMF by minimizing the  
496 number of reaction media and separation steps. Indeed, no intermediate product or phase isolation  
497 was necessary. The use of numerous organic solvents or ionic liquid, as previously described in the  
498 literature, is limited to a unique organic phase used for fructose transportation. Improvement of  
499 performances in terms of HMF yields implying better diffusion of species between the different  
500 phases by improving the conception of new reactor systems will be reported in due course.

501

502

503

## 504 Acknowledgements

505 The authors want to thank IFP Energies nouvelles for funding these studies. Chevreul Institute (FR  
506 2638), Ministère de l'Enseignement Supérieur et de la Recherche, Région Hauts-de-France (CPER  
507 ALIBIOTECH), REALCAT platform ('Future Investments' program (PIA), with the contractual  
508 reference 'ANR-11-EQPX-0037') and FEDER are acknowledged for supporting and funding partially  
509 this work.

510

## 511 References

- 512 1. Nanda, S.; Mohammad, J.; Reddy, S. N.; Kozinski, J. A.; Dalai, A. K. Pathways of lignocellulosic biomass  
513 conversion to renewable fuels. *Biomass Conv. Bioref.* **2014**, *4*, 157–191.
- 514 2. Besson, M.; Gallezot, P.; Pinel, C. Conversion of Biomass into Chemicals over Metal Catalysts. *Chem. Rev.*  
515 **2014**, *114*, 1827–1870.
- 516 3. Aresta, M.; Dibenedetto, A.; Dumeignil, F. *Biorefineries: From Biomass to Chemicals and Fuels*, 3<sup>rd</sup> ed.; De  
517 Gruyter: Berlin, Boston, 2012; ISBN 978-3-11-026023-6
- 518 4. Agbor, V.B.; Cicek, N.; Sparling, R.; Berlin, A.; Levin, D.B. Biomass pretreatment: Fundamentals toward  
519 application. *Biotechnol. Adv.* **2011**, *29*, 675–682.
- 520 5. Bozell, J. J., Petersen, G. R. Technology development for the production of biobased products from  
521 biorefinery carbohydrates—the US Department of Energy's 'Top 10' revisited. *Green Chem.* **2010**, *12*,  
522 539–545.
- 523 6. Van Heerden, C.D., Nicol, W. Continuous succinic acid fermentation by *Actinobacillus succinogenes*.  
524 *Biochem. Eng. J.* **2013**, *73*, 5–11.
- 525 7. Girard, E.; Delcroix, D.; Cabiac, A. Catalytic conversion of cellulose to C2–C3 glycols by dual association of  
526 a homogeneous metallic salt and a perovskite-supported platinum catalyst. *Catal. Sci.* **2016**, *6*, 5534–5542.
- 527 8. Shanks, B. H. Unleashing biocatalysis/chemical catalysis synergies for efficient biomass conversion. *ACS*  
528 *Chem. Biol.* **2007**, *2*, 533–535.
- 529 9. Vennestrøm, P. N. R., Taarning, E., Christensen, C.H., Pedersen, S., Grunwaldt, J.D., Woodley, J.M.  
530 Chemoenzymatic Combination of Glucose Oxidase with Titanium Silicalite-1. *ChemCatChem* **2010**, *2*, 943–  
531 945.
- 532 10. Denard, C. A., Hartwig, J. F., Zhao, H. Multistep One-Pot Reactions Combining Biocatalysts and Chemical  
533 Catalysts for Asymmetric Synthesis. *ACS Catal.* **2013**, *3*, 2856–2864.
- 534 11. Dumeignil, F. *Public Serv. Rev. Eur. Union* **2011**, *22*, 528.
- 535 12. F. Dumeignil, *Chemie Ing. Tech.*, 2014, *86*, 1496.

- 536 13. Van Putten, R., Van der Waal, J.C., de Jong, E., Rasrendra, C.B., Heeres, H.G., de Vries, J.G.  
537 Hydroxymethylfurfural, A versatile platform chemical made from renewable resources. *Chem. Rev.* **2013**,  
538 *113*, 1499–1597.
- 539 14. Moliner, M.; Román-Ishkov, Y., Davis, M. E. Tin-containing zeolites are highly active catalysts for the  
540 isomerization of glucose in water. *Proc Natl Acad Sci U S A.* **2010**, *107*, 6164–6168.
- 541 15. Pacheco, J. J., Davis, M. E. Synthesis of terephthalic acid via Diels-Alder reactions with ethylene and  
542 oxidized variants of 5-hydroxymethylfurfural. *Proc Natl Acad Sci U S A.* **2014**, *111*, 8363–8367.
- 543 16. Lin, Z., Nikolakis, V., Ierapetritou, M. Alternative Approaches for p-Xylene Production from Starch:  
544 Techno-Economic Analysis. *Ind. Eng. Chem. Res.* **2014**, *53*, 10688–10699.
- 545 17. E. Jong, M.A. Dam, L. Sipos, and G. Gruter. Furandicarboxylic Acid (FDCA), A Versatile Building Block  
546 for a Very Interesting Class of Polyesters *Am. Chem. Soc.* **2012**, *1*, 1-13.
- 547 18. Boisen, A., Christensen, T.B., Fu, W., Gorbanev, Y.Y., Hansen, T.S., Jensen, J.S., Klitgaard, S.K., Pedersen,  
548 S., Riisager, A., Stahlberg, T., Woodley, J.M. *Chem. Eng. Res. Des.* **2009**, *87*, 1318-1325.
- 549 19. Akien, G.R., Qi, L., Horvath, I.T. Molecular mapping of the acid catalysed dehydration of fructose. *Chem.*  
550 *Commun.* **2012**, *48*, 5850-5852.
- 551 20. Parker, K., Salas, M., Nwosu, V.C. High fructose corn syrup: Production, uses and public health concerns  
552 *Biotechnol. Mol. Biol. Rev.* **2010**, *5*, 71-78.
- 553 21. Li, H., Yang, S., Saravanamurugan, S., Riisager, A. Glucose Isomerization by Enzymes and  
554 Chemo-catalysts: Status and Current Advances. *ACS Catal* **2017**, *7*, 3010-3029.
- 555 22. Delidovich, I., Palkovits, R. Catalytic Isomerization of Biomass-Derived Aldoses: A Review. *ChemSusChem*  
556 **2016**, *9*, 547–561.
- 557 23. Choudhary, V., Pinar, A.B., Lobo, Vlachos, D.G., Sandler, S.I. Comparison of homogeneous and  
558 heterogeneous catalysts for glucose-to-fructose isomerization in aqueous media. *ChemSusChem* **2013**, *6*,  
559 2369-2376.
- 560 24. Marianou, A.A., Michailof, C.M., Pineda, A., Iliopoulou, E.F., Triantafyllidis, K.S., Lappas, A.A. Glucose to  
561 Fructose Isomerization in Aqueous Media over Homogeneous and Heterogeneous Catalysts.  
562 *ChemCatChem* **2016**, *8*, 1100-1110.
- 563 25. Zhao, S., Guo, X., Bai, P., Lv, L. Catalytic Isomerization of Biomass-Derived Aldoses: A Review. *Asian J.*  
564 *Chem.* **2014**, *26*, 4537-4542.
- 565 26. Demerdash, M., Attia, R. M. Equilibrium kinetics of D-glucose to D-fructose isomerization catalyzed by  
566 glucose isomerase enzyme from *Streptomyces phaeochromogenus*. *Zentralbl. Mikrobiol.* **1992**, *147*, 297–303.
- 567 27. Gaily, M.H., Elhassan, B.M., Abasaeed A.E., Al-shrhan, M. Kinetics of a Three-Step Isomerization of  
568 Glucose to Fructose Using Immobilized Enzyme. *Int. J. Eng. Technol.* **2010**, *10*, 1-6.
- 569 28. Takasaki, Y. Kinetic and Equilibrium Studies on D-Glucose-D-Fructose Isomerization Catalyzed by  
570 Glucose Isomerase from *Streptomyces* sp. *Agric. Biol. Chem.* **1967**, *31*, 309–313.
- 571 29. Huang, H., Denard, C.A., Alamillo, R., Crisci, A.J., Miao, Y., Dumesic, J.A., Scott, S.L., Zhao, H. Tandem  
572 catalytic conversion of glucose to 5-hydroxymethylfurfural with an immobilized enzyme and a solid acid.  
573 *ACS Catal.* **2014**, *4*, 2165-2172.
- 574 30. Huang, R., Qi, W., Su, R., He, Z. Integrating enzymatic and acid catalysis to convert glucose into  
575 5-hydroxymethylfurfural. *Chem. Commun. (Camb).* **2010**, *46*, 1115-1125.
- 576 31. Delidovich, I., Palkovits, P. Catalytic Isomerization of Biomass-Derived Aldoses: A Review. *Green Chem.*  
577 **2016**, *18*, 5822-5830.
- 578 32. Alipour, S., Relue, P.A., Viamajala, S., Varanasi, S. High yield 5-(hydroxymethyl)furfural production from  
579 biomass sugars at facile reaction conditions: a hybrid enzyme- and chemo-catalytic technology. *Green*  
580 *Chem.* **2016**, *18*, 4990-5002.
- 581 33. Gimbernat, A., Guehl, M., Capron, M., Lopes-Ferreira, N., Froidevaux, R., Girardon, J.S., Dhulster, P.,  
582 Delcroix, D., Dumegnil, F. Hybrid Catalysis: A Suitable Concept for the Valorization of Biosourced  
583 Saccharides to Value-Added Chemicals. *ChemCat Chem* **2017**, *9*, 2080-2888.
- 584 34. Bhosale, S.H., Rao, M.B., Deshpande, V.V. Molecular and industrial aspects of glucose isomerase.  
585 *Microbiol. Rev.* **1996**, *60*, 280-290.
- 586 35. McKay, G., Tavlarides, T. Enzymatic isomerization kinetics of D-Glucose to D-Fructose. *J. Mol. Catal.* **1979**,  
587 *6*, 57-65.

- 588 36. Morin, G.T., Paugam, M.F., Hughes, M.P., Smith, B.D. Boronic Acids Mediate Glycoside Transport  
589 through a Liquid Organic Membrane via Reversible Formation of Trigonal Boronate Esters. *J. Org. Chem.*  
590 **1994**, *59*, 2724-2730.
- 591 37. Karpa, M.J., Duggan, P.J., Griffin, G.J., Freudigmann, S.J. Competitive transport of reducing sugars  
592 through a lipophilic membrane facilitated by aryl boron acids. *Tetrahedron* **1997**, *53*, 3669-3676.
- 593 38. Westmark, P.R., Gardiner, S.J., Smith, B.D. Selective monosaccharide transport through lipid bilayers  
594 using boronic acid carriers. *J. Am. Chem. Soc.* **1996**, *118*, 11093-11100.
- 595 39. Shinbo, T., Nishimura, K., Yamaguchi, T., Sugiura, M. Transport of monosaccharides across an organic  
596 liquid membrane. *J. Chem. Soc. Chem. Commun.* **1986**, 349-358.
- 597 40. Paugam, M.; Riggs, J.A.; Smith, B.D. High fructose syrup production using fructose-selective liquid  
598 membranes. *ChemComm.* **1996**, *22*, 2539-2540.
- 599  
600