

1	<b><u>SUPPORTING INFORMATION SECTION 1: LIST OF EQUATIONS, SYMBOLS AND INDICES</u></b>	<b>2</b>
2	<b><u>SUPPORTING INFORMATION SECTION 2: IMPACT CATEGORIES DESCRIPTION</u></b>	<b>9</b>
3	CLIMATE CHANGE	9
4	MARINE ACIDIFICATION	10
5	MINERAL AND FOSSIL RESOURCES USE	10
6	ACIDIFICATION	11
7	TOXICITY IMPACTS	12
8	PARTICULATE MATTER FORMATION	14
9	WATER AVAILABILITY IMPACT	14
10	LAND USE	15
11	PHOTOCHEMICAL OXIDANT FORMATION, IONIZING RADIATION, AND OZONE LAYER DEPLETION	16
12	<b><u>SUPPORTING INFORMATION SECTION 3: INTERIM IMPACT CATEGORIES</u></b>	<b>18</b>
13	CLIMATE CHANGE IMPACTS ON RESOURCES & ECOSYSTEM SERVICES	18
14	LAND USE IMPACTS ON RESOURCES & ECOSYSTEM SERVICES	18
15	WATER AVAILABILITY IMPACTS	18
16	WATER AVAILABILITY IMPACTS ON RESOURCES & ECOSYSTEM SERVICES	19
17	MARINE AND TERRESTRIAL ECOTOXICITY	19
18	PHOTOCHEMICAL OXIDANT FORMATION	20
19	IONIZING RADIATION	20
20	OZONE LAYER DEPLETION	20
21	<b><u>SUPPORTING INFORMATION SECTION 4: DETAILED IMPACT WORLD+ FRAMEWORK</u></b>	<b>20</b>
22	<b><u>SUPPORTING INFORMATION SECTION 5: CHARACTERIZATION FACTORS DATABASE</u></b>	<b>22</b>
23	<b><u>SUPPORTING INFORMATION SECTION 6: OVERALL MODEL COHERENCE</u></b>	<b>23</b>
24	MODELLING ASSUMPTIONS COHERENCE.	23
25	TEMPORAL COHERENCE	24
26	SPATIAL COHERENCE	24
27	<b><u>SUPPORTING INFORMATION SECTION 7: RESIDENCE TIME IN AIR INFLUENCE ON THE INTAKE</u></b>	<b>26</b>
28	<b><u>FRACTIONS OF A SUBSTANCE</u></b>	<b>26</b>
29	<b><u>SUPPORTING INFORMATION SECTION 8: SPATIAL VARIABILITY OF TOXIC IMPACTS</u></b>	<b>27</b>
30	<b><u>REFERENCES</u></b>	<b>32</b>
31		

## 1 **Supporting information section 1: List of equations, symbols and indices**

---

### 2 **Acronyms in text**

3 AoP Area of Protection

4 AoC Area of Concern

5  $\text{Damage}_{\text{AOP}}$  Impact category unit for corresponding AoP

6

7 CF Characterization factor

8 FF Fate factor

9 XF Exposure factor

10 ERF Exposure-response factor

11 SF Severity factor

12 CSI Competition scarcity index, that expresses the quantity of resource that is going to  
13 deprive competing users (current or future) sharing the same resource per quantity of  
14 resource used in a dissipative manner

15

16  $\Delta_x$  Difference in metric x described in subscript

17  $t_1$  lower time limit for time integration of impacts;  $t_1 = 100$  yr for damage level long-term  
18 indicators,  $t_1 = 0$  for all other indicators.

19  $t_2$  higher time limit for time integration of impacts;  $t_2 = 100$  yr for shorter term indicators,  
20  $t_2 = 500$  yr for climate change, long-term indicators,  $t_2 = \infty$  for other indicators

21

### 22 **Indices - superscripts**

23 s = elementary flow (or environmental intervention)

24 k = impact category (midpoint or damage level)

25 AOP = Area of Protection

26 AOC = Area of Concern

27

### 28 **Indices - subscripts**

29 i = emitting environmental compartment

30 j = receiving environmental compartment

31

32 a = emitting region (or spatial unit) (native or aggregated at country, continental or global  
33 level)

34 n = native emitting region (or spatial unit) for a considered impact category

35 b = receiving region (or spatial unit)

36

37 p = exposure pathway per unit of active substance

38 r = response due to a change in exposure of human population or ecosystems

39 u = user affected by the competition for the resource

40

41 midpoint = midpoint level

42 damage = damage level

43

44 **Abbreviations and Symbols** (units will usually depend on the considered impact category)

1  $I^k$  Impact score at midpoint or damage levels for impact category k  
2  $m_i^s$  Inventory elementary flow s emitted to or extracted from an emitting compartment i  
3  $CF_i^{sk}$  Characterization factor for elementary flow s, impact category k and emitting  
4 compartment i  
5  $I_a^k$  Impact score at midpoint or damage levels for impact category k and an emitting region a  
6  $M_{ia}^s$  Inventory elementary flow s emitted to or extracted from an emitting compartment i  
7 and an emitting region a  
8  $CF_{ai}^{sk}$  Characterization factor for elementary flow s, impact category k and emitting  
9 compartment i and an emitting region a  
10  
11  $FF_{ji}^{sk}$  Fate factor for impact category k and for elementary flow s emitted into or extracted  
12 from emitting compartment i and transfer into receiving compartment j  
13  $XF_{pj}^{sk}$  Exposure factor for impact category k and for elementary flow s in the receiving  
14 compartment j through the exposure pathway p  
15  $ERF_{rp}^{sk}$  Exposure-response factor, for impact category k and elementary flow s, for the  
16 response r due to a change in exposure pathway p of human population or ecosystems  
17  $SF_r^{sk}$  Severity factor, for impact category k and elementary flow s, for a response r due to a  
18 change in exposure of human population or ecosystems  
19  $CSI_{ui}^{sk}$  Competition scarcity index factor for impact category k and elementary flow s, that  
20 expresses the quantity of resource in compartment i that is going to deprive competing user  
21 u (current or future) sharing the same resource per quantity of resource used in a  
22 dissipative manner  
23  $XF_{pu}^{sk}$  Exposure factor, for impact category k and elementary flow s, for the user u affected  
24 by the competition for the resource through the exposure pathway p  
25  
26  $CF_{ni}^{sk}$  Characterization factor for elementary flow s, impact category k and emitting  
27 compartment i and an emitting native region n  
28  $FF_{jni}^{sk}$  Fate factor for impact category k and for elementary flow s emitted into or extracted  
29 from emitting compartment i and emitting native region n and transfer into receiving  
30 compartment j and receiving region b  
31  $XF_{pbj}^{sk}$  Exposure factor for impact category k and for elementary flow s in the receiving  
32 compartment j and receiving region b through the exposure pathway p  
33  $ERF_{rbp}^{sk}$  Exposure-response factor, for impact category k and elementary flow s, for the  
34 response r in receiving region b due to a change in exposure pathway p of human  
35 population or ecosystems  
36  $CSI_{ubni}^{sk}$  Competition scarcity index factor for impact category k and elementary flow s, that  
37 expresses the quantity of resource in compartment i and in receiving region b that is going  
38 to deprive competing user u (current or future) sharing the same resource per quantity of  
39 resource used in a dissipative manner  
40  $XF_{pbu}^{sk}$  Exposure factor, for impact category k and elementary flow s, for the user u affected  
41 by the competition for the resource through the exposure pathway p in receiving region b  
42  
43  $SPF_{ni}^{sk}$  Spatial proportionality factor used to aggregate  $CF_{ni}^{sk}$  into a coarser resolution a (for  
44 instance national, continental, or global).

- 1  $F_{ni}^S$  overall annual environmental intervention for elementary flow  $s$  within the native  
2 spatial unit  $n$  for an emitting compartment  $i$   
3  $F_{ai}^S$  overall annual environmental intervention for elementary flow  $s$  within the region  $a$  for  
4 an emitting compartment  $i$   
5  $A_n$  surface area of the native spatial unit  $n$   
6  $A_a$  surface area of the emitting spatial unit  $a$   
7  $A_{n \cap a}$  surface area intersecting of the native spatial unit  $n$  and the emitting spatial unit  $a$   
8  
9  $S^{AOP}$  Aggregated impact score in AOP  
10  $S^{AOC}$  Aggregated impact score in AOC  
11  
12  $NF^{AOP}$  Normalization factors of IMPACT World+ for AOP  
13  $I_{world\ annual}^k$  Impact score at damage levels for impact category  $k$  due to the total world  
14 emissions and extractions for one year  
15  $N_{world\ pop}$  Total world population count  
16  $S_{world\ annual}^{AOP}$  Aggregated impact score in AOP due to all the total world emissions and  
17 extractions for one year  
18  
19

## 20 1. The impact score at midpoint or damage level for an impact category

$$22 I^k = \sum_s \overrightarrow{CF}^{sk} \cdot \overrightarrow{m}^s \text{ (Eq. SI1)}$$

23 With:

- 24 •  $Dim(x)$  is the number of elements along the  $x$  axis
- 25 •  $I^k$  a scalar
- 26 •  $\overrightarrow{CF}^{sk}$  a vector with  $(Dim(i))$  dimension and elements defined as  $CF_i^{sk}$
- 27 •  $\overrightarrow{m}^s$  a vector with  $(Dim(i))$  dimension and elements defined as  $m_i^s$
- 28 • The operator “.” is defined as the inner product (or dot product) between vector  
29 such as  $V = \overrightarrow{T} \cdot \overrightarrow{U}$  is equivalent to  $V = \sum_i T_i U_i$ .

31 And elements given by:

$$32 I^k = \sum_s \sum_i CF_i^{sk} m_i^s \text{ (Eq. SI1.1)}$$

### 34 1.1. For regionalized impact calculation

$$36 \mathbf{I}^k = \sum_s \overrightarrow{\mathbf{I}} \cdot \mathbf{CF}^{sk} \circ \mathbf{M}^s \text{ (Eq. SI1_regio)}$$

37 With:

- 38 •  $Dim(x)$  is the number of elements along the  $x$  axis
- 39 •  $\mathbf{I}^k$  a vector with  $(Dim(a))$  dimension and elements defined as  $I_a^k$
- 40 •  $\mathbf{CF}^{sk}$  a matrix with  $(Dim(a) \times Dim(i))$  dimensions and elements defined as  $CF_{ai}^{sk}$
- 41 •  $\mathbf{M}^s$  a matrix with  $(Dim(i) \times Dim(a))$  dimensions and elements defined as  $M_{ia}^s$
- 42 •  $\overrightarrow{\mathbf{I}}$  a vector with  $Dim(i)$  dimension and all elements equal to 1
- 43 • The operator “o” is defined as the Hadamard product such as  $\mathbf{V} = \mathbf{T} \circ \mathbf{U}$  with elements  
44 given by  $V_{ij} = T_{ij} U_{ij}$ .

- The operator “.” is defined as the inner product (or dot product) such as  $V = T \cdot U$  elements given by  $V_{ik} = \sum_j T_{ij} U_{jk}$ .

And elements given by:

$$I_a^k = \sum_s \sum_i CF_{ai}^{sk} M_{ia}^s \text{ (Eq. SI1_regio.1)}$$

## 2. Damage CF for an impact category and for an elementary flow

### 2.1. Damage CF for emission related impact category

#### 2.1.1. For non-regionalized impact category

$$\overrightarrow{CF}^{sk} = \overrightarrow{SF}^{sk} \cdot \mathbf{ERF}^{sk} \cdot \mathbf{XF}^{sk} \cdot \mathbf{FF}^{sk} \text{ (Eq. SI2)}$$

With:

- $Dim(x)$  is the number of elements along the  $x$  axis
- $\overrightarrow{CF}^{sk}$  a vector with  $(Dim(i))$  dimension and elements defined as  $CF_i^{sk}$
- $\mathbf{FF}^{sk}$  a matrix with  $(Dim(j) \times Dim(i))$  dimensions and elements defined as  $FF_{ji}^{sk}$
- $\mathbf{XF}^{sk}$  a matrix with  $(Dim(p) \times Dim(j))$  dimensions and elements defined as  $XF_{pj}^{sk}$
- $\mathbf{ERF}^{sk}$  a matrix with  $(Dim(r) \times Dim(p))$  dimensions and elements defined as  $ERF_{rp}^{sk}$
- $\overrightarrow{SF}^{sk}$  a vector with  $(Dim(r))$  dimension and elements defined as  $SF_r^{sk}$
- The operator “.” is defined as the inner product (or dot product) such as  $V = T \cdot U$  elements given by  $V_{ik} = \sum_j T_{ij} U_{jk}$ .

And elements given by:

$$CF_i^{sk} = \sum_r SF_r^{sk} \left( \sum_p ERF_{rp}^{sk} \left( \sum_j XF_{pj}^{sk} FF_{ji}^{sk} \right) \right) \text{ (Eq. SI2.1)}$$

Which is equivalent to the following equation:

$$CF_i^{sk} = \sum_r \sum_p \sum_j SF_r^{sk} ERF_{rp}^{sk} XF_{pj}^{sk} FF_{ji}^{sk} \text{ (Eq. SI2.2)}$$

#### 2.1.2. For regionalized impact category

$$\mathbf{CF}^{sk} = \vec{1} \cdot \overrightarrow{SF}^{sk} \cdot \mathbf{ERF}^{sk} \cdot \mathbf{XF}^{sk} \cdot \mathbf{FF}^{sk} \text{ (Eq. SI2_regio)}$$

With:

- $Dim(x)$  is the number of elements along the  $x$  axis
- $\mathbf{CF}^{sk}$  a matrix with  $(Dim(a) \times Dim(i))$  dimensions and elements defined as  $CF_{ni}^{sk}$
- $\mathbf{FF}^{sk}$  a matrix (or tensor) with  $(Dim(j) \times Dim(b) \times Dim(n) \times Dim(i))$  dimensions and elements defined as  $FF_{jbni}^{sk}$
- $\mathbf{XF}^{sk}$  a matrix (or tensor) with  $(Dim(p) \times Dim(b) \times Dim(j))$  dimensions and elements defined as  $XF_{pbj}^{sk}$

- $\mathbf{ERF}^{\text{sk}}$  a matrix (or tensor) with  $(Dim(r) \times Dim(b) \times Dim(p))$  dimensions and elements defined as  $ERF_{rbp}^{\text{sk}}$
- $\overrightarrow{\mathbf{SF}}^{\text{sk}}$  a vector with  $(Dim(r))$  dimension and elements defined as  $SF_r^{\text{sk}}$
- $\vec{\mathbf{1}}$  a vector with  $Dim(b)$  dimension and all elements equal to 1
- The operator “.” is defined as the inner product (or dot product) between tensors, matrices and vectors such as  $\mathbf{V} = \mathbf{T} \cdot \mathbf{U}$  elements given by  $V_{ijk} = \sum_l T_{ijl} U_{lk}$ .

And elements given by:

$$CF_{ni}^{\text{sk}} = \sum_b (\sum_r SF_r^{\text{sk}} (\sum_p ERF_{rbp}^{\text{sk}} (\sum_j XF_{pbj}^{\text{sk}} FF_{jbni}^{\text{sk}}))) \quad (\text{Eq. SI2\_regio.1})$$

Which is equivalent to the following equation:

$$CF_{ni}^{\text{sk}} = \sum_b \sum_r \sum_p \sum_j SF_r^{\text{sk}} ERF_{rbp}^{\text{sk}} XF_{pbj}^{\text{sk}} FF_{jbni}^{\text{sk}} \quad (\text{Eq. SI2\_regio.2})$$

## 2.2. Damage CF for resource related impact category

### 2.2.1. For non-regionalized impact category

$$\overrightarrow{\mathbf{CF}}^{\text{sk}} = \overrightarrow{\mathbf{SF}}^{\text{sk}} \cdot \mathbf{ERF}^{\text{sk}} \cdot \mathbf{XF}^{\text{sk}} \cdot \mathbf{CSI}^{\text{sk}} \quad (\text{Eq. SI2})$$

With:

- $Dim(x)$  is the number of elements along the  $x$  axis
- $\overrightarrow{\mathbf{CF}}^{\text{sk}}$  a vector with  $(Dim(i))$  dimension and elements defined as  $CF_i^{\text{sk}}$
- $\mathbf{CSI}^{\text{sk}}$  a matrix with  $(Dim(u) \times Dim(i))$  dimensions and elements defined as  $CSI_{ui}^{\text{sk}}$
- $\mathbf{XF}^{\text{sk}}$  a matrix with  $(Dim(p) \times Dim(u))$  dimensions and elements defined as  $XF_{pu}^{\text{sk}}$
- $\mathbf{ERF}^{\text{sk}}$  a matrix with  $(Dim(r) \times Dim(p))$  dimensions and elements defined as  $ERF_{rp}^{\text{sk}}$
- $\overrightarrow{\mathbf{SF}}^{\text{sk}}$  a vector with  $(Dim(r))$  dimension and elements defined as  $SF_r^{\text{sk}}$
- The operator “.” is defined as the inner product (or dot product) such as  $\mathbf{V} = \mathbf{T} \cdot \mathbf{U}$  elements given by  $V_{ik} = \sum_j T_{ij} U_{jk}$ .

And elements given by:

$$CF_i^{\text{sk}} = \sum_r SF_r^{\text{sk}} (\sum_p ERF_{rp}^{\text{sk}} (\sum_u XF_{pu}^{\text{sk}} CSI_{ui}^{\text{sk}})) \quad (\text{Eq. SI2.1})$$

Which is equivalent to the following equation:

$$CF_i^{\text{sk}} = \sum_r \sum_p \sum_u SF_r^{\text{sk}} ERF_{rp}^{\text{sk}} XF_{pu}^{\text{sk}} CSI_{ui}^{\text{sk}} \quad (\text{Eq. SI2.2})$$

### 2.2.2. For regionalized impact category

$$\mathbf{CF}^{\text{sk}} = \vec{\mathbf{1}} \cdot \overrightarrow{\mathbf{SF}}^{\text{sk}} \cdot \mathbf{ERF}^{\text{sk}} \cdot \mathbf{XF}^{\text{sk}} \cdot \mathbf{CSI}^{\text{sk}} \quad (\text{Eq. SI2\_regio})$$

With:

- $Dim(x)$  is the number of elements along the  $x$  axis
- $\mathbf{CF}^{\text{sk}}$  a matrix with  $(Dim(n) \times Dim(i))$  dimensions and elements defined as  $CF_{ni}^{\text{sk}}$

- 1 •  $\mathbf{CSI}^{sk}$  a matrix (or tensor) with  $(Dim(u) \times Dim(b) \times Dim(n) \times Dim(i))$  dimensions
- 2 and elements defined as  $CSI_{ubni}^{sk}$
- 3 •  $\mathbf{XF}^{sk}$  a matrix (or tensor) with  $(Dim(p) \times Dim(b) \times Dim(u))$  dimensions and
- 4 elements defined as  $XF_{pbu}^{sk}$
- 5 •  $\mathbf{ERF}^{sk}$  a matrix (or tensor) with  $(Dim(r) \times Dim(b) \times Dim(p))$  dimensions and
- 6 elements defined as  $ERF_{rbp}^{sk}$
- 7 •  $\overrightarrow{\mathbf{SF}}^{sk}$  a vector with  $(Dim(r))$  dimension and elements defined as  $SF_r^{sk}$
- 8 •  $\vec{\mathbf{1}}$  a vector with  $Dim(b)$  dimension and all elements equal to 1
- 9 • The operator “.” is defined as the inner product (or dot product) between tensors,
- 10 matrices and vectors such as  $\mathbf{V} = \mathbf{T} \cdot \mathbf{U}$  elements given by  $V_{ijk} = \sum_l T_{ijl} U_{lk}$ .

11 And elements given by:

$$12 \quad CF_{ni}^{sk} = \sum_b (\sum_r SF_r^{sk} (\sum_p ERF_{rbp}^{sk} (\sum_u XF_{pbu}^{sk} CSI_{ubni}^{sk}))) \quad (\text{Eq. SI2\_regio.1})$$

13 Which is equivalent to the following equation:

$$14 \quad CF_{ni}^{sk} = \sum_b \sum_r \sum_p \sum_u SF_r^{sk} ERF_{rbp}^{sk} XF_{pbu}^{sk} CSI_{ubni}^{sk} \quad (\text{Eq. SI2\_regio.2})$$

### 15 3. Spatial aggregation of CF for spatially differentiated impact categories

16 The aggregated CF for a given region a, for an elementary flow s, for an impact category k

17 and for an emitting compartment I is given by:

$$18 \quad CF_{ai}^{sk} = \sum_n CF_{ni}^{sk} \times SPF_{ni}^{sk} \quad (\text{Eq. SI3})$$

$$19 \quad SPF_{ni}^{sk} = \frac{F_{ni}^s \times \frac{A_{n\Omega a}}{A_a}}{\sum_n F_{ni}^s \times \frac{A_{n\Omega a}}{A_a}} = \frac{F_{ni}^s \times \frac{A_{n\Omega a}}{A_a}}{F_{ai}^s} \quad (\text{Eq. SI3.1})$$

20  $\frac{A_{n\Omega a}}{A_a} = 1$  when the spatial unit n is totally included in the spatial unit a.

### 21 4. Midpoint level equations

22 Depending on the position of the midpoint indicator along the cause-effect chain, midpoint

23 level characterization factors for emission-related impact categories follow Equation SI4.1

24 (freshwater eutrophication, water scarcity), SI4.2 (Climate change shorter term, climate

25 change long term, terrestrial acidification, freshwater acidification, SI4.3 (marine

26 eutrophication, freshwater ecotoxicity, human toxicity cancer, human toxicity non cancer,

27 particulate matter formation, photochemical oxidant formation, ionising radiation HH,

28 ionising radiation EQ) or SI4.4 (ozone layer depletion, water stream use and management,

29 land transformation biodiversity, land occupation biodiversity) with

30  $SF_r^{sk}, ERF_{rbp}^{sk}, XF_{pbj}^{sk}, FF_{jbni}^{sk}$  defined as in damage level equations:

$$31 \quad CF_{ni}^{sk} = \sum_b \sum_j FF_{jbni}^{sk} \quad (\text{Eq SI.4.1})$$

$$32 \quad CF_{ni}^{sk} = \sum_b (\sum_j XF_{pbj}^{sk} FF_{jbni}^{sk}) \quad (\text{Eq SI.4.2})$$

$$33 \quad CF_{ni}^{sk} = \sum_b (\sum_p ERF_{rbp}^{sk} (\sum_j XF_{pbj}^{sk} FF_{jbni}^{sk})) \quad (\text{Eq SI.4.3})$$

1  $CF_{ni}^{sk} = \sum_b (\sum_r SF_r^{sk} (\sum_p ERF_{rbp}^{sk} (\sum_j XF_{pbj}^{sk} FF_{jbni}^{sk})))$  (Eq SI.4.4)

2  
3 For both resource-related impact categories (mineral resources use, fossil energy use),  
4 midpoint level characterization factor follows Equation SI4.5 with  $XF_{pu}^{sk}$ ,  $CSI_{ui}^{sk}$  defined as in  
5 damage level equations:

6  $CF_i^{sk} = \sum_p \sum_u XF_{pu}^{sk} CSI_{ui}^{sk}$  (Eq SI4.5)

7  
8 **5. Overall damage on the AoPs considering all the different impact categories at**  
9 **damage level**

10  
11  $S^{AOP} = \sum_{k \in AOP} I^k$  (Eq. SI5)

12  
13  
14 **6. Overall damage contributing to the same AoC**

15  
16  $S^{AOC} = \sum_{k \in AOC} I^k$  (Eq. SI6)

17  
18 **7. Normalization factors**

19  $NF^{AOP} = \frac{\sum_{k \in AOP} I_{world\ annual}^k}{N_{world\ pop}} = \frac{S_{world\ annual}^{AOP}}{N_{world\ pop}}$  (Eq. SI7)

20



## 1 **Supporting information section 2: impact categories description**

---

### 2 **Climate change**

3 A greenhouse gas emission first leads to an increase in atmospheric concentration, which  
4 then leads to positive radiative forcing. This positive radiative forcing may cause different  
5 climate effects such as an increase in temperature or precipitation changes. IMPACT  
6 World+ focuses on the temperature increase pathway, not considering the other types of  
7 climate effects due to a lack of data and knowledge. An increase in the Earth average  
8 temperature leads to potential impacts on humans and ecosystems through several  
9 pathways.

10  
11 The fate factor corresponds to the time-integrated mass of the greenhouse gas for a given  
12 mass emitted in the atmosphere. It is determined using the impulse response function  
13 proposed by Joos *et al.* (2013) and adopted by the IPCC (Myhre 2013). The exposure factor  
14 gives the time-integrated increase in temperature due to this time integrated mass in the  
15 atmosphere. This exposure factor can be decomposed in two sub-indicators: the time-  
16 integrated radiative forcing due to the time-integrated mass in the atmosphere, and the  
17 time-integrated increase in temperature due to the time-integrated radiative forcing.

18  
19 The Global Warming Potential for a 100-year time horizon (GWP100) as adopted by the  
20 IPCC (Myhre 2013) is used as a midpoint indicator for shorter-term climate change,  
21 characterizing the cumulative radiative forcing per kg greenhouse gas emitted. The Global  
22 Temperature Potential for a 100-year time horizon (GTP100), as also proposed in the latest  
23 IPCC report (Myhre 2013), is our second midpoint indicator for long-term climate change  
24 impacts and represents a change in global mean surface temperature at a chosen point in  
25 time. It is therefore not a cumulative indicator, but it is consensually considered as an  
26 appropriate proxy to represent climate change long-term impacts (Levasseur *et al.* 2016).  
27 Those two indicators are needed because they express different impacts: GTP100 (climate  
28 change long-term) are impacts related to long-term cumulative warming (e.g. sea level rise),  
29 while GWP100 (climate change shorter-term) are impacts related to a rapid increase in  
30 temperature to which humans and species must adapt very quickly.

31  
32 At the damage level, cumulative metrics need to be considered in LCIA to ensure additivity  
33 of impacts (Frischknecht *et al.* 2016). Hence the GTP100 cannot be used. To model the  
34 impact up to the damage, we therefore use the time-integrated temperature increase  
35 calculated from absolute GTP (aGTP) equations, as proposed by the IPCC (Myhre 2013). In  
36 compliance with what is proposed by the IPCC, the effect of CO<sub>2</sub> that is formed from the  
37 oxidation of CH<sub>4</sub> and CO is considered, but not the oxidation products of other VOCs (see  
38 supporting info, Section xxx for details). This time-integrated temperature increase is thus a  
39 combination of fate and exposure factors of humans and ecosystems to climate change.  
40 Human health effect factors are calculated based on the increase in risk of dying associated  
41 with a time-integrated temperature increase, as proposed by de Schryver *et al.* (2009),  
42 building on a study from the World Health Organization (2003). It provides the relative risk  
43 of dying from five different causes (cardiovascular diseases, malaria, diarrhoea, floods and  
44 malnutrition). This represents what is feasible today, and it is likely to represent only a  
45 fraction of the actual DALYs caused by climate change – the majority of which will likely  
46 results from conflicts in an unstable world (Barnett and Adger 2007). This is thus a proxy of

1 the lower bond of human health damage from climate change. The severity corresponding  
2 to each cause of risk (DALY/case) are taken from the Global Burden-of-Disease report  
3 (Mathers et al. 2008) for cardiovascular diseases, malaria, diarrhoea and malnutrition,  
4 while the International Disaster Database (2009) is used in combination with the Global  
5 Burden-of-Disease data for unintentional injuries to estimate DALY/case for floods. For  
6 ecosystem quality, as proposed by de Schryver *et al.* (2009), effect factors are calculated  
7 from a study compiling a number of regional studies that aim at predicting the extinction of  
8 species related to an increase in temperature (Thomas et al. 2004) considering a global  
9 surface of semi-natural terrestrial areas of the world of  $2.29 \cdot 10^{13} \text{ m}^2$ . Both damage level  
10 impact indicators on ecosystems and human health are temporally resolved: shorter-term  
11 cumulative impacts are calculated for a time horizon within 100 years from the emission,  
12 while long-term cumulative impacts are calculated between 100 years and 500 years.

13  
14 IMPACT World+ also considers additional interim impact pathways related to climate  
15 change affecting the “resources and ecosystem services” AoP, as described the supporting  
16 information Section 4.

17

### 18 **Marine acidification**

19 Carbon dioxide released into the atmosphere partly partitions into the oceans, and reacts  
20 with the water to form carbonic acid. Some of these carbonic acid molecules dissociate to  
21 give bicarbonate and hydronium ions, thus increasing the ocean's acidity ( $\text{H}^+$  ion  
22 concentration). Other chemical reactions are also triggered (shift in the carbonate system  
23 toward lower pH), which results in an actual net decrease in the amount of carbonate ions  
24 available. In the oceans, this makes it more difficult for marine calcifying organisms, such as  
25 coral and some plankton, to form biogenic calcium carbonate, and such existing structures  
26 become vulnerable to dissolution.

27

28 To ensure consistency with climate change modelling, a) the fate model for  $\text{CO}_2$  emissions is  
29 the same as for climate change (Myhre 2013); b)  $\text{CH}_4$  and  $\text{CO}$  are also classified within this  
30 impact category to consider the effect of  $\text{CO}_2$  that is formed from the oxidation of  $\text{CH}_4$  and  
31  $\text{CO}$ ; c) the model is temporally resolved, differentiating between impacts in the first 100  
32 years and long-term impacts occurring between 100 and 500 years. The exposure factor  
33 considers the decrease in ocean pH due to an increase in  $\text{CO}_2$  in the atmosphere (Azevedo et  
34 al. 2015). The effect factor is based on a species sensitivity distribution (SSD) curve using  
35 the  $\text{HC50}_{\text{EC50}}$  (i.e. the  $\text{H}^+$  concentration affecting 50% of the population of 50% of the  
36 considered species) due to pH modification on marine ecosystems (Azevedo et al. 2015).  
37 The effect model uses a linearity assumption of the SSD curve between 0 and the  $\text{HC50}_{\text{EC50}}$ .  
38 This is consistent with the way SSD curves are used in USEtox to generate effect factors for  
39 ecotoxicity.

40

### 41 **Mineral and fossil resources use**

42 For mineral resources use impact, IMPACT World+ uses the material competition scarcity  
43 index from de Bruille (2014) as a midpoint indicator. This factor represents the fraction of  
44 material needed by future users that are not able to adapt to a full dissipation of the easily  
45 available stock. It is expressed in terms of kg of deprived resource per kg of dissipated  
46 resource.

1  
2 For fossil energy use impact, IMPACT World+ uses the primary energy content  
3 (Frischknecht 2003) as a midpoint indicator considering that it is a reasonable proxy to  
4 assess the MJ deprived per MJ consumed, under the assumption that fossil resources are  
5 mainly functional for energy purposes. A functionality specific effect factor from Fatemi  
6 (2012) for fossil energy use and from De Bruille (2014) for mineral resources use is then  
7 applied, giving the adaptation price to fulfil the need of non-adapted users using a backup  
8 technology (\$/kg deprived) to obtain the interim damage level CF.

9  
10 These models are operational to assess mineral and fossil resources both at midpoint and  
11 damage levels, but have only yet been applied at damage level to coal and petroleum for  
12 fossil energy use and to aluminium, cadmium, cobalt, lithium, manganese, nickel and rare  
13 earths for mineral resources, assuming today's technology for backup. These damage level  
14 CFs are therefore considered as interim.

## 16 Acidification

17 When emitted to the atmosphere, acidifying substances will disperse, react with other  
18 substances in the atmosphere and travel potentially long distances before depositing on soil  
19 and/or water. These deposits may change the soil and water acidity levels. For every  
20 species, there is an optimum pH range, and a deviation from this optimum may be harmful  
21 for that specific specie. Consequently, a change in pH may decrease the species distribution  
22 in an ecosystem.

23  
24 The characterization factors presented by Roy *et al.* (Roy et al. 2014; Roy et al. 2012a; Roy  
25 et al. 2012b) are used for terrestrial and freshwater acidification respectively, both at  
26 midpoint and damage levels. The midpoint characterization factors express the change in  
27 pH in receiving environments (soil and freshwater, respectively) due to an emission of  
28 nitrogen oxides (NO<sub>x</sub>), ammonia (NH<sub>3</sub>) and sulphur dioxide (SO<sub>2</sub>). It combines fate  
29 assessment addressing atmospheric source-deposition relationships using the GEOS Chem  
30 model and soil sensitivity assessment for terrestrial acidification. An additional soil fate  
31 assessment gives the transfer of H<sup>+</sup> to freshwater ecosystems for freshwater acidification.  
32 Damages on ecosystem quality are expressed in terms of PDF·m<sup>2</sup>·yr per kg of substance  
33 emitted and computed combining the midpoint CF with an effect factor (Roy et al. 2014).  
34 For terrestrial acidification, the latter determines the change in vascular plant species per  
35 change in the soil pH. For freshwater acidification, it measures the change of potentially  
36 disappeared fraction of fish species per change in the water pH.

## 38 Eutrophication

39 Eutrophication is the result of increased nutrient loading to a surface water body, driving  
40 growth of primary producers, changing species abundance and diversity, ultimately leading  
41 to decreased levels of oxygen that affect freshwater or coastal ecosystems. As the  
42 underlying (simplifying) assumption, phosphorous is considered the only limiting nutrient  
43 causing freshwater eutrophication, whereas nitrogen is modelled as the limiting nutrient  
44 for marine coastal zones.

1 For freshwater eutrophication, the work of Helmes *et al.* (2012) is used to determine the  
 2 fate factor of phosphorous in freshwater at a 0.5° x 0.5° resolution. This fate factor  
 3 expresses the increase in phosphorus mass per kg<sub>P</sub> discharged to freshwater and is used as  
 4 the midpoint CF for freshwater eutrophication. A stoichiometric ratio between substances  
 5 is used to generate CFs for other P containing substances (phosphate, phosphoric acid,  
 6 phosphorus pentoxide) and an equivalency factor of 0.022 kg PO<sub>4</sub><sup>3-</sup>eq/kg was considered for  
 7 COD and BOD as recommended in CML (Guinée *et al.* 2002). The damage factor is obtained  
 8 multiplying the midpoint CF by an effect factor of 11.4 PDF·m<sup>2</sup>·yr/kg PO<sub>4</sub><sup>3-</sup>eq from Tirado-  
 9 Seco (2005).

10  
 11 For marine eutrophication, the coastal zone considered is the zone where the ocean depth  
 12 is less than 200 m, consistently with the (sub-)continental parameterisation of USEtox  
 13 (Kounina *et al.* 2014). The same atmospheric fate model as used by Roy *et al.* (2012b) for  
 14 acidification (GEOS Chem) is used to determine the source-to-deposition relationship of  
 15 Ammonia (NH<sub>3</sub>) and (NO<sub>x</sub>) atmospheric emissions on coastal zones. For emissions to  
 16 freshwater, 70% of the N containing substances discharged is assumed to reach the coastal  
 17 zone as done in ReCiPe and EDIP (Goedkoop *et al.* 2013; Hauschild and Potting 2005;  
 18 Hauschild and Wenzel 1998). This reflects the fact that elimination due to denitrification in  
 19 anaerobic zones in freshwater is treated as a constant with a generic removal of 30 % in the  
 20 CARMEN European model used in both LCIA methods. Hence, 70 % of the nitrogen input  
 21 transports to sea. An empirically-based EF of 12.5 PDF·m<sup>2</sup>·yr/kg<sub>deposited</sub> has been  
 22 determined based on the ratio between observed eutrophied areas in highly eutrophied  
 23 regions (Gulf of Mexico, Baltic sea, Chesapeake Bay) and nitrogen load.  
 24 Eutrophication in the Gulf of Mexico, Baltic Sea and Chesapeake Bay are primarily caused  
 25 by Nitrogen flux into the water system. To analyse this relationship and to get a correlation  
 26 between the nutrient fluxes into the aquatic ecosystem in consideration and the extent of  
 27 hypoxia, the metrics PDF m<sup>2</sup> year/kg N is used (see results in table S3.1). The average value  
 28 of 12.5 PDF m<sup>2</sup> yr/ kgN is used as the marine eutrophication effect factor in IMPACT  
 29 World+.

30  
 31 **Table SI 5.1 : Hypoxic areas/kg nutrients for various affected water systems**

Water Body	Analysis Years	Total Nitrogen/ Phosphorus (t/year)	Average Hypoxia Area (km <sup>2</sup> )	Hypoxia Area/ kgN (PDF m <sup>2</sup> year/ kg N)
Gulf of Mexico	<b>1985-2010</b>	<b>1 419 760</b>	<b>13 810</b>	<b>9,7</b>
Gulf of Mexico	2005-2010	1 368 200	17 300	<b>12,6</b>
Chesapeake Bay	1985-2011	91 330,45	1 183,20	<b>13,0</b>
Chesapeake Bay	1970-2011	94 823,95	1 105,40	<b>11,7</b>
Baltic Sea	1995-2009	3 729 000,00	48 000,00	<b>12,9</b>

32  
 33 An update of the Marine eutrophication indicator is to be expected in the near future to  
 34 account for recent research developments done by Cosme (Cosme and Hauschild 2016).  
 35

### 36 Toxicity impacts

37 Outdoor and indoor emissions of chemical substances may cause toxic effects to human  
 38 health and to ecosystems. Once emitted into air, freshwater or soil, the substances may

1 reach and affect freshwater, terrestrial and marine ecosystems. Humans may also be  
2 exposed to these substances through different pathways (e.g. inhalation of air, ingestion of  
3 food and water), which may cause multiple health outcomes and diseases in different  
4 human and/or ecosystem populations.

5  
6 The UNEP/SETAC scientific consensus model USEtox for characterizing human toxicity and  
7 ecosystem toxicity impacts (Hauschild et al. 2008; Rosenbaum et al. 2008), that accounts  
8 for fate, exposure and effects of chemicals, is used and adapted to generate the IMPACT  
9 World+ (eco)toxicity CFs. Version 2.0 of USEtox is used to determine the global default CFs,  
10 including continental versions based on the work of Kounina et al. (2014). Characterization  
11 factors for human toxicity modeled with USEtox are expressed at the midpoint level in  
12 comparative toxic units ( $CTU_h$ ) per unit mass of a chemical emitted, providing the estimated  
13 increase in morbidity in the global human population per unit mass of a chemical emitted  
14 into a specific environmental compartment (disease cases per kg emitted) both for cancer  
15 and non-cancer diseases. Since these two indicators should not be directly summed up  
16 without severity assessment (Rosenbaum et al. 2011), damage level CFs are calculated  
17 using severity factors of 11.5 and 2.7 DALY (Huijbregts et al. 2005) per disease case for  
18 cancer and non-cancer, respectively. Human toxicity CFs also include toxicity impacts from  
19 indoor emissions – using the USEtox indoor CFs for household and industrial indoor  
20 emissions – considering different archetypes for OECD countries and non OECD airtight and  
21 non airtight buildings (Hellweg et al. 2009; Rosenbaum et al. 2015; Wenger et al. 2012) and  
22 pesticide residues ingestion from crops (Fantke et al. 2011; Fantke and Jolliet 2016; Fantke  
23 et al. 2012) for both cancer and non-cancer effects.

24  
25 Of special interest is the case of metals, which have typically contributed significantly to  
26 toxic impacts. There is a need to account for the essentiality of zinc, since most of the world  
27 population is reported to be deficient in Zn. We therefore only applied the toxicity non-  
28 cancer characterization factor of Zn to the small fraction of the population that may be  
29 exposed at potentially toxic levels – 2% according to expert judgement – (Nriagu 2014).

30  
31 For freshwater ecotoxicity, USEtox midpoint level CFs are expressed as comparative toxic  
32 units ( $CTU_e$ ) per per unit mass of a chemical emitted, providing an estimation of the  
33 potentially affected fraction (PAF) of the exposed ecosystem species integrated over time  
34 and water volume per unit mass of a chemical emitted ( $PAF \cdot m^3 \cdot day/kg$ ). A generic severity  
35 factor of 0.5 is applied to convert PAF to PDF – the potentially disappeared fraction of  
36 species – based on the assumption that 50% of the affected species will disappear from the  
37 ecosystem after exposure (Jolliet et al. 2003a). An average surface water depth of 2.5 m is  
38 used to yield a damage expressed in  $PDF \cdot m^2 \cdot day/kg$ .

39 *We acknowledge that the continental resolution with population density archetypes used to*  
40 *determine the fate in USEtox is meaningful for human toxicity, but not as relevant for*  
41 *ecotoxicity, because the target organisms are not distributed in the same way. Moreover,*  
42 *USEtox doesn't attempt to represent the spatial variability of (eco)toxicity. Further research is*  
43 *needed to adequately model the spatial variability of this impact category in a manageable*  
44 *way in an LCA context, knowing that thousands of substances have to be characterized for*  
45 *each native spatial unit.*

1 USEtox CFs are differentiated between shorter-term impacts taking place over the first 100  
2 years and long-term impacts from 100 years to infinity,, of which the latter are only  
3 substantial for very persistent substances, such as metals.

4  
5 Interim CFs are also proposed for marine and terrestrial ecotoxicity impacts and are  
6 described in supporting information, Section 4.  
7

## 8 **Particulate matter formation**

9 Inhalation of fine particulate matter (PM<sub>2.5</sub>), i.e. particles with diameter less than 2.5 µm, is  
10 known to cause a number of health related issues and reduction in life expectancy,  
11 including chronic and acute respiratory and cardiovascular morbidity, chronic and acute  
12 mortality, lung cancer, diabetes, and adverse birth outcomes (Fantke et al. 2015). PM<sub>2.5</sub> is  
13 composed of primary and secondary particles. The latter originates from the oxidation of  
14 primary gases such as sulfur oxides and nitrogen oxides into ammonium sulfates and  
15 ammonium nitrates PM<sub>2.5</sub>.

16  
17 Characterization factors are modelled using epidemiologically derived factors from  
18 Humbert et al. (2011) and Gronlund et al. (2015). PM<sub>10</sub> is considered in those publications  
19 1.67 times less toxic than PM<sub>2.5</sub> and converted into PM<sub>2.5</sub> equivalents via this factor. The  
20 midpoint CFs account for fate, exposure, and effect. Intake fractions for primary PM<sub>2.5</sub> are  
21 defined as the fractions of the emission taken in (inhaled) by the overall population  
22 (Bennett et al. 2002; Hodas et al. 2015) and are consistent with USEtox fate factors using  
23 archetypes for remote, rural and urban outdoor environments. The intake fraction for  
24 secondary PM<sub>2.5</sub> is the inhaled mass of PM<sub>2.5</sub> attributable to (i.e. formed from) a specific  
25 precursor substance per mass emitted of this precursor. midpoint CFs are expressed in  
26 PM<sub>2.5-eq</sub> per kg, and correspond to the number of deaths per kilogram emitted normalised  
27 using PM<sub>2.5</sub> as a reference substance. Damage level CFs are calculated assuming 0.0083  
28 DALY/kg PM<sub>2.5-eq</sub>, which corresponds to an average severity factor of 19 DALY per death for  
29 cardiopulmonary disease and lung cancer.  
30

## 31 **Water availability impact**

32 IMPACT World+ uses the water scarcity AWARE model (Boulay et al. 2016) at the midpoint  
33 level as a proxy midpoint for all the water scarcity impacts. It is not directly on the cause  
34 effect chain leading to impacts on human health nor on the one leading to impacts on  
35 ecosystems, as no common midpoint exists between both cause effect chains, but it  
36 combines both users water needs (humans and ecosystems) to assess the water scarcity.  
37 This index is based on the remaining water available per area after human and aquatic  
38 ecosystem demand has been met, relative to the world average. It can be interpreted as the  
39 hypothetical surface-time equivalent necessary to generate an unused volume of water in a  
40 specific watershed, compared to the world average. AWARE is recommended by the  
41 UNEP/SETAC Life Cycle Initiative and the European Commission.  
42

43 The approach of Boulay *et al.* (2011) is used to model the water availability impacts on  
44 human health. It includes a CSI (expressed in m<sup>3</sup> deprived per m<sup>3</sup> dissipated), an XF, which  
45 characterizes exposure of competing users to deprivation and accounts for adaptation  
46 capacity and water functionality (i.e. only competing users unable to adapt will suffer

1 human health impacts) and some function-specific EF. The latter are applied to obtain the  
2 impacts on human health per m<sup>3</sup> deprived, focusing on the irrigation, domestic use and  
3 fisheries functions that are directly affecting human health. The model allows to accounts  
4 for the fact that consuming bad quality water affects less competing users (quality specific  
5 interim CFs are available) but the recommended default CFs do not account for water  
6 quality as LCA tools and databases are not mature enough to integrate water quality (see  
7 supporting information, Section 4 for details).

8  
9 For damages on ecosystem quality, several methods are combined to model the cause-effect  
10 chain for water scarcity impacts, as recommended by Kounina *et al.* (2013). The work of  
11 Hanafiah *et al.* (2011) is used to model impacts of freshwater consumption on freshwater  
12 ecosystems and the model of van Zelm *et al.* (2011) is used to assess water scarcity impacts  
13 on terrestrial ecosystems from groundwater. A CF of 0 is considered for deep groundwater.  
14 Despite Kounina *et al.* (2013) suggestion to use it in combination with the other models  
15 assessing water scarcity impact on ecosystem quality, the model from Pfister *et al.* (2009)  
16 has not been integrated in IMPACT World+. This was done in order to keep the overall  
17 coherence of the method and to avoid double counting of the impact of groundwater  
18 consumption on terrestrial ecosystems (see supporting info, Section 4 for details). The  
19 model from Verones *et al.* (2010) is used to assess the impact of thermally polluted water,  
20 with the assumption that the “Water, cooling, unspecified natural origin” currently found in  
21 life cycle inventory databases with no further specification is released in a 3 m deep river  
22 with a 4 °C temperature increase.

23  
24 IMPACT World+ also considers as interim indicators the impact of water availability on the  
25 resources & ecosystem services AoP, and water stream use and management impacts on  
26 ecosystem quality AoP, modelled using the Humbert & Maendly model (2009), both being  
27 described in supporting information, Section 4 and not yet being peer reviewed.

## 29 Land use

30 Human activities cause impacts to lands, which are either converted from natural state  
31 (land transformation) or occupied, i.e. maintained in a certain non-natural state (land  
32 occupation). Impacts on land have consequences in terms of terrestrial biodiversity but also  
33 in terms of fundamental ecosystem services for the human society such as biotic production,  
34 water regulation, freshwater recharge and filtration, climate regulation and erosion  
35 resistance.

36  
37 Potential impacts of land occupation and land transformation on ecosystem quality are  
38 characterized using local empirical CFs at the biome level. Those CFs, from de Baan *et al.*  
39 (2013) were preferred over the regional and global CFs from Chaudhary *et al.* (2015) –  
40 despite the latter being preliminary recommended by the UNEP/SETAC Life Cycle Initiative.  
41 This choice was done to ensure coherence with the other impact categories on the physical  
42 meaning of impact scores at the damage level. *Regional and global impacts as modelled by*  
43 *Chaudhary et al. (2015) are meaningful as they give important complementary information*  
44 *on the (semi)-irreversible disappearance of species in a region or at the global scale. However,*  
45 *they are not consistent with the other ecosystem quality impact indicators of IMPACT World+*  
46 *(acidification, eutrophication, ecotoxicity, water scarcity, etc.) or any other existing LCIA*

1 *method, because their indicator has a different meaning. The regional PDF indicator from*  
2 *Chaudhary et al. (2015) quantifies the fraction of species disappeared forever from a region of*  
3 *the world, whereas the PDF.m2.yr indicators of the IMPACT World+ framework quantify the*  
4 *temporary disappearance of species (PDF) over a given surface (m2) during a certain time*  
5 *(yr). To our understanding, it is hardly possible to convert the regional PDF from Chaudhary et*  
6 *al. (2015) into PDF.m2.yr, because the former corresponds to a permanent disappearance and*  
7 *would therefore lead to an infinite impact. Using the Curran (2010) regeneration time to*  
8 *convert those PDFs in PDF.m2.yr – as proposed by Chaudhary et al. (2015) when calculating*  
9 *land transformation impact – seems inconsistent as this regeneration time is supposed to*  
10 *apply to an ecosystem and not to the irreversibly disappeared species. However, by doing so,*  
11 *the land use impact scores would be orders of magnitude lower than the local impacts of de*  
12 *Baan et al. (2013) as only a very small fraction of the species affected by land occupation or*  
13 *transformation will permanently disappear from the affected region. We therefore decided to*  
14 *use the de Baan et al. (2013) model, which quantifies damage indicators in PDF.m2.yr that are*  
15 *consistent with the damage indicators used in IMPACT World+. Nevertheless, we acknowledge*  
16 *the need to further develop regional CFs for the other impact categories in line with the*  
17 *approach proposed by Chaudhary et al. (2015) as complementary information to the*  
18 *assessment of local impacts as currently done in life cycle impact assessment.*

19  
20 For the case of land occupation impacts on terrestrial biodiversity, there is no need to  
21 consider fate and exposure since effects occur directly in the environmental compartment  
22 where the environmental intervention is reported. The characterization factor therefore  
23 directly relates the land occupation (in m<sup>2</sup>·yr) to the biodiversity loss (in PDF·m<sup>2</sup>·yr).  
24 Midpoint indicators are thus determined by dividing damage values by the global average  
25 CF for arable land, used as a reference, expressing these normalized damage values in  
26 m<sup>2</sup>·yr<sub>eq</sub> of arable land. Land transformation impacts are considered as long-term impacts  
27 only, as they represent the remaining impacts once the occupation ends and the ecosystem  
28 recovers which, in most cases, will occur in the far future. They were calculated using the  
29 land occupation CFs and considering the recovery times from Curran (2010) as proposed  
30 by Chaudhary *et al.* (2015).

31  
32 IMPACT World+ also considers, as interim indicators, the potential impact of land  
33 occupation and land transformation on the resources and ecosystem services AoP  
34 (including the following ecological soil functions: erosion resistance, freshwater recharge,  
35 physical and chemical filtration). Similarly, indirect land use impacts on human health due  
36 to the modification of freshwater recharge, or physical and chemical filtration are  
37 considered interim and computed consistently with the model used to assess water  
38 availability impacts on human health. Such indirect impacts from land transformation on  
39 the ecosystem quality AoP are not considered in IMPACT World+ as this may lead to double  
40 counting: the empirical CFs from de Baan et al. (2013) include all the impact pathways  
41 affecting local ecosystems, including the modification of its access to water. All those  
42 interim impact indicators are further described in supporting information, Section 4.

#### 44 **Photochemical oxidant formation, ionizing radiation, and ozone layer depletion**

45 The ILCD handbook recommendations (European Commission 2011) were followed for  
46 these three impact categories. Model calculations were updated to account for the most up-



- 1 to-date World Meteorological Organization (WMO (World Meteorological Organization)
- 2 2014) values of ozone depletion potential.

## 1 Supporting information section 3: Interim impact categories

### 2 Climate change impacts on resources & ecosystem services

3 As proposed by Cao et al (2015) for land use impacts on carbon sequestration potential of  
 4 soils, the social cost of carbon proposed by Ackerman and Stanton (2010) is used as an  
 5 estimation of the cost due to the loss of service due to climate change. This effect factor is  
 6 applied directly to the midpoint (and not to the temperature increase) as this value was  
 7 initially determined based on GWP100.

### 8 Land use impacts on resources & ecosystem services

9 IMPACT World+ also consider as interim categories the impact of land transformation as  
 10 well as the potential impact of land occupation on the following soil ecological functions:  
 11 erosion prevention, groundwater recharge, physical and chemical filtration and are  
 12 described in section xxx of the supporting information. The fate factors (Saad et al. 2011)  
 13 describes the modification of biophysical indicators per quantity of land occupation (in m<sup>2</sup>-  
 14 yr). Biophysical indicators are further modelled with respect to the loss of ecosystem  
 15 services using the model of Cao et al. (2015). Damage oriented CFs express the social cost to  
 16 compensate (or adapt) to the loss of ecosystem service related to the change of soil  
 17 ecological function due to land use ( $\$/\text{Service}_{\text{lost}}/\text{m}^2\text{-yr}$ ). These are calculated as the  
 18 multiplication of a dimensionless exposure factor defining the loss of service per change of  
 19 the biophysical indicator ( $\text{Service}_{\text{lost}}/\Delta \text{Biophys. Indicator } i$ ) and an effect factor ( $\$/\text{service}$   
 20  $\text{lost}$ ) that monetizes the value of a given ecosystem service. The exposure factor accounts  
 21 for the fraction of service loss that is affecting the population and for the adaptation  
 22 capacity of the society.

23 Given the still exploratory state of these models, we consider the biophysical midpoints and  
 24 damage CFs as interim. Thus, only land occupation impacts on biodiversity are considered  
 25 in the recommended version of IMPACT World+.

### 26 Water availability impacts

27 Water availability impacts on Human health CFs are provided for the different level of  
 28 water quality detailed in the table SI2 to complement the default water availability impact  
 29 on human health CFs which are not accounting for the water quality.

30

31 *Table SI 4.1: Description of the water quality classes*

	Excellent	Good	Average	Average-Tox	Average-Bio	Poor	Very poor	Unusable
<b>S Surface</b>	1	2a	2b	2c	2d	3	4	5
<b>G Groundwater</b>								
<b>Coliforms</b>	low	low	medium	low	high	high	high	Unusable
<b>Toxics</b>	low	medium	medium	high	low	medium	high	

32

33 Those CFs are to be used with care, acknowledging a lack of coherence between human  
 34 toxicity impacts modeled with USEtox and this water availability model, which may lead to  
 35 an overlap and a potential double counting.

36 Those water quality classes were originally developed to acknowledge the fact that a bad  
 37 quality water is not functional for as many users as a good quality water, hence using some  
 38 non conventional sources of water (such as for example treated wastewater) would have  
 39 less impact in term of users deprivation than using very high quality water.

1 However, this assumption lead to a bias in the operationalization of the model, as a water  
2 balance is made to account only for the quantity of water that is consumed by a process.  
3 The quantity of water withdrawn and the quantity of water released are quantified and  
4 characterized to obtain the net water availability impact. If the water is released with a  
5 lower quality than the water originally withdrawn, the model mathematically considers  
6 that some usages of the water are “lost” due to quality degradation, leading to water  
7 availability impacts.

8 1) This is debatable due to the fate in surface water of the released emission which may  
9 lead to acceptable level of contamination quickly after the release, hence not depriving any  
10 competing users;

11 2) This is not consistent with the assumptions done in USEtox, both in term of toxic  
12 contaminant fate when emitted to water and in term of human exposure to toxic emission  
13 through drinking water. The water availability model considers that domestic users will be  
14 deprived of good quality water and will turn toward other bad quality water sources or  
15 adopt bad hygiene habits leading to diseases, ie will not consumed the degraded water,  
16 whereas USEtox assumes that a fraction of the toxic emission to be consumed through  
17 drinking water. Both assumptions are inconsistent.

18  
19 The model from Pfister et al. (2009) has not been integrated in IMPACT World+ in order to  
20 keep the overall coherence and to avoid double counting the impact of groundwater  
21 consumption on terrestrial ecosystems. This model assesses the impact of water  
22 consumption (surface and groundwater) on plants, hence partly overlapping the impact  
23 pathway covered by the van Zelm *et al.* (2011) model which is focusing on the plant water  
24 deprivation due to water table lowering when pumping groundwater. We consider that van  
25 Zelm *et al.* (2011) has a more robust fate model based on hydrogeology to determine the  
26 fraction of water pumped which is going to deprive the plants from soil moisture. Pfister et  
27 al. (2009) considers this fraction as being 100%, which is very conservative . The other  
28 impact pathway covered by Pfister et al. (2009) is the water deprivation of terrestrial  
29 plants due to surface water consumption, which is not covered in IMPACT World+. Pfister  
30 et al. (2009) use again the very conservative assumption that 100% of the water pumped in  
31 surface water will deprive terrestrial plants in the watershed.

### 32 **Water availability impacts on resources & ecosystem services**

33 The same CSI scarcity index is used at midpoint for water use impacts on resources and  
34 ecosystem services. The exposure factor is complementary to the exposure factor  
35 considered for human health impacts: only competing users able to adapt will have to pay  
36 for this adaptation. The effect factor is the same as the one considered in land use impacts  
37 for groundwater recharge and filtration (in both cases, the same “service” is lost for the  
38 society and the adaptation costs to access to good quality water is the same)(Cao et al.  
39 2015). At this stage, these unpublished characterization factors of water use impacts on  
40 resources and ecosystem services are considered interim.

41

### 42 **Marine and terrestrial ecotoxicity**

43 Interim CFs are also proposed for marine and terrestrial ecotoxicity based on the USEtox  
44 fate factors for the coastal zone and for the natural soil environmental compartments. The  
45 marine effect factors are considered equal as per freshwater ecotoxicity. As a side remark,  
46 no differentiation is currently done currently in the freshwater ecotox EFs determination in

1 USEtox between marine and freshwater species as both are mixed in the aquatic ecotox  
2 databases used to determine USEtox EFs, meaning that the assumption of similar ecosystem  
3 sensitivity is already implicitly done. The terrestrial ecosystem EF are extrapolated from  
4 the freshwater EF using the soil-water partition coefficient proposed by Hauschild and  
5 Wenzel (1998), as already done in IMPACT 2002+ (Jolliet et al. 2003b).

#### 6 **Photochemical oxidant formation**

7 This category is related to the impacts of ozone and other reactive oxygen compounds  
8 formed as secondary contaminants in the troposphere (the region in the atmosphere  
9 closest to the surface) by the oxidation of the primary contaminants (Non Methane Volatile  
10 Organic Compounds, NMVOC, or carbon monoxide) in the presence of nitrogen oxides  
11 (NO<sub>x</sub>) and under the influence of light. Ozone concentrations in the troposphere lead to  
12 increased frequency and severity of respiratory diseases, such as asthma and Chronic  
13 Obstructive Pulmonary Diseases. IMPACT World+ uses the method used in ReCiPe  
14 (Goedkoop et al. 2013) and recommended by the ILCD handbook (Margni et al. 2008) for  
15 both mid-points and end-points.

#### 16 **Ionizing radiation**

17 The routine releases of radioactive material to the environment is responsible of both  
18 human health and ecosystem effects. Human health characterization factors are taken from  
19 Frischknecht et al. (Frischknecht et al. 2000) and characterization factors for ecosystems  
20 are based on the approach of Garnier-Laplace et al. (Garnier-Laplace et al. 2008), which is  
21 used consistently with the ecotoxicity assessment in USEtox as per the recommendations of  
22 the ILCD Handbook (Margni et al. 2008).

#### 23 **Ozone Layer depletion**

24 Ozone depleting substances emitted by human activity destroy the ozone layer in the  
25 stratosphere, which blocks UVB, by breaking ozone molecules into molecular oxygen  
26 through heterogeneous catalysis. Exposure to UVB radiations increases the risk of skin  
27 cancer and cataract. It may also cause premature aging and suppression of the immune  
28 system. As recommended by the ILCD handbook, midpoint characterization factors are  
29 based on the ozone depletion potentials produced by the World and Global  
30 Meteorological Organisation (WMO (World Meteorological Organization) 2014) using the  
31 infinite time perspective and human health severity factors developed by Struijs et al.  
32 (Struijs et al. 2009). Model calculations were updated to account for the most up to date  
33 WMO2014 values of ozone depletion potential.


34  
35  
36

#### 37 **Supporting information section 4: Detailed IMPACT World+ framework**

---



**IMPACT World+ detailed framework**



## 1 **Supporting information section 5: Characterization factors database**

---

2 The database joined as supporting information section 5 includes the characterization  
3 factors for all the recommended impact categories of IMPACT World+.

4  
5 Alternatively, it can be found in the following dropbox file:

6 <https://www.dropbox.com/sh/2sdgbqf08yn91bc/AAA-mnN7YxkQxfyFx2LYK0PCa?dl=0>

7  
8 For the regionalized impact categories, the characterization factors are available at four  
9 different resolution scales: at the native resolution scale and aggregated at the country,  
10 continent and global level. *Regionalized CFs are available for all 197 countries considered in*  
11 *the ISO 3166-2 norm (2013), which nomenclature is used to name the regionalized*  
12 *environmental intervention*. For the aggregated characterization factors, the uncertainty due  
13 to spatial variability within the geographical cell considered is also documented in the  
14 database with information on the minimum, the maximum, the 5<sup>th</sup>, 25<sup>th</sup>, 75<sup>th</sup> and 95<sup>th</sup>  
15 centiles, the mean and the weighted average using the probability of emission in each of the  
16 aggregated native resolution cells.

17 *The compartments considered in the recommended impact categories are the following:*

- 18 • *Air*
  - 19 ○ *With sub-compartments based on population density archetypes (high/low) for*  
20 *toxic and ecotoxic impacts as well as particulate matter formation and ionizing*  
21 *radiation*
  - 22 ○ *And an additional indoor sub-compartment for toxic impact*
- 23 • *Soil*
  - 24 ○ *With a sub compartment agricultural soil for human toxicological and*  
25 *ecotoxicological impacts as well as ionizing radiation*
- 26 • *Water*
  - 27 ○ *With sub compartments groundwater, lake, river, ocean for water use impacts*
  - 28 ○ *With sub compartment ocean for human toxicological, ecotoxicological,*  
29 *ionizing radiation impacts*
  - 30 ○ *With sub compartments groundwater and ocean for eutrophication impacts*
- 31 • *Raw material*
  - 32 ○ *With sub compartment land for land use impacts*
  - 33 ○ *With sub compartment water for water use impact*

34  
35  
36  
37

## 1 Supporting information section 6: Overall model coherence

---

### 2 Modelling assumptions coherence.

3 The same models are used to model similar environmental mechanisms for different impact  
4 categories:

- 5 • The same coherent framework was used across all the impact categories  
6 considering fate (or competition in the case of resources), exposure and effect
- 7 • Resources are all modelled using a functionality based approach, ie considering the  
8 loss of resources functional value for the humans (and not any intrinsic value loss)
- 9 • Water scarcity/availability and land use impact categories are strongly inter-related  
10 and modelled coherently: when land use is influencing the quality or the quantity of  
11 groundwater (through impacts on water recharge or water filtration), the water  
12 availability impacts are applied to complement the cause effect chain toward human  
13 health and resources & ecosystem services areas of protection.
- 14 • Climate change impacts and land use impacts on soil carbon sequestration are  
15 strongly inter-related and the loss of carbon sequestration potential is directly  
16 modelled under the climate change impact category.
- 17 • The method proposed by Cao et al to assess the impacts of land transformation on  
18 resources & ecosystem services area of protection has been coherently applied to  
19 other climate change contributing emission in the interim version of the  
20 methodology.
- 21 • The atmospheric fate of CO<sub>2</sub>, CO and CH<sub>4</sub>, contributing to both climate change and  
22 marine acidification impact categories, is computed with the same model from the  
23 IPCC.
- 24 • For both climate change and marine acidification, the CO<sub>2</sub> resulting from the quick  
25 oxidation of CO and CH<sub>4</sub> was considered in the model.
- 26 • The effect model for marine acidification uses a linearity proxy assumption of the  
27 SSD curve of H<sup>+</sup> concentration between 0 and the HC50<sub>EC50</sub>. This is consistent with  
28 the way SSD curves are used in USEtox to generate effect factors for ecotoxicity.
- 29 • Cancer, non cancer, marine, terrestrial and freshwater ecotoxicity models use the  
30 same fate factors, from USEtox
- 31 • The exposure and effect modelling assumptions across all those toxicity related  
32 impact categories are all coherent.
- 33 • The fate models for indoor exposure and pesticide residues that are integrated in  
34 the cancer and non-cancer impact categories are also coherent in term of fate,  
35 exposure and effect with the USEtox model.
- 36 • The atmospheric fate of NO<sub>x</sub> and ammonia contributing to terrestrial acidification,  
37 aquatic acidification and marine eutrophication was computed with the same model  
38 (GEOS-Chem).
- 39 • The warming effect of CO<sub>2</sub> that is formed from the oxidation of methane is now  
40 included in the GWP value of methane published in the IPCC report (Myhre 2013).  
41 For consistency purposes, we decided to add a GWP value for CO to account for the  
42 warming effect of CO<sub>2</sub> that is formed from its oxidation. Since the lifetime of CO is  
43 very short (a few months), we consider that a molecule of CO instantly becomes a  
44 molecule of CO<sub>2</sub>, leading to a GWP value equal to the ratio of their molar mass

1 (44/28 = 1.57) since 1 kg of CO will oxidize into 1.57 kg CO<sub>2</sub>. For consistency  
2 purpose, the formed CO<sub>2</sub> is also considered in marine acidification impact category.  
3 *Remark : Should this be extended to other non-GHG substances containing carbon and*  
4 *nitrogen that could ultimately be degraded into CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O, depending on the*  
5 *aerobic or anaerobic conditions in which they are degraded? This is less straight*  
6 *forward as on the one hand the degradation products depend on the degradation*  
7 *conditions, and on the other hand, some chemicals can take years to achieve full*  
8 *degradation, leading to delayed GHG emissions. The half-life of a chemical in a given*  
9 *environmental compartment should be put into perspective with the 100 years time*  
10 *frame of the mid-point characterization factors to keep consistency. In the case of*  
11 *methane and CO, oxidation is occurring quite rapidly. An attempt was made by*  
12 *(Muñoz et al. 2013) to calculate the potential GHG emissions (CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O) from*  
13 *the degradation of a few organic chemicals, showing that it may not be negligible.*  
14 *However, the climate change effect of GHGs formed from the mineralization of organic*  
15 *compounds was considered not mature enough to be integrated in the present version*  
16 *of IMPACT World+ and should be studied further.*

### 17 **Temporal coherence**

18 Temporal coherence is respected across impact categories:

- 19 • For all long term the impact categories, a “shorter term” time horizon has been set  
20 coherently at 100 yrs after the emission occurs. This allows to express separately
- 21 • Long-term impacts are the remaining impacts after 100 yrs.
- 22 • Long-term impacts are integrated to the infinite when there is a full recovery.
- 23 • When some permanent remaining impacts occur, a 500 yrs time horizon has been  
24 chosen consistently across impact categories (this is the case for marine  
25 acidification and climate change) as an infinite time horizon would lead in that case  
26 to an almost infinite impact.
- 27 • Remark: to ensure temporal coherence between inventory and impact assessment,  
28 long-term emissions in the inventory (i.e. emitted in more than 100 yrs) should be  
29 considered to only have long-term impact.

### 30 **Spatial coherence**


31 Spatial coherence is also respected across impact categories:

- 32 • The same geographical parameterisation was used across all the regionalized  
33 impact categories. For example the coastal zone considered for impacts on marine  
34 ecosystems is the same for marine acidification, marine ecotoxicity and marine  
35 eutrophication and is defined as the zone where the depth of water is less than 200  
36 meters.
- 37 • The same maps and projections are used to define countries and continents, the  
38 same geographical data (population density, etc) are used to model native  
39 resolution CFs and to aggregate them at the country / continental / global level.
- 40 • The same approach has been used across all the regionalized impact categories to  
41 generate CFs at the country / continent / global level based on the probability of  
42 emission (or resource consumption) in each of the native resolution scale  
43 geographical cell.



- 1 • For all the regionalized impact categories, when no CF is available for a specific  
2 region of the world, the global default CF is used and should be considered with the  
3 corresponding spatial variability.
- 4 • The modelling assumptions used for respiratory inorganic are coherent with the  
5 USEtox modelling assumption in term of definition of the urban archetype  
6 (population density)  
7  
8  
9

1 **Supporting information section 7: Residence time in air influence on the intake fractions of a**  
 2 **substance**




3  
 4 *Figure SI 7.1: Indoor (OECD country average archetype), urban and rural intake fraction as a*  
 5 *function of the fate factor (persistence) in rural continental air*  
 6


1 **Supporting information section 8: Spatial variability of toxic impacts**

2 Carcinogen characterization factors for an emission to soil and water, non-carcinogen  
3 characterization factors for an emission to air, soil and water and freshwater  
4 ecotoxicological characterization factors for an emission to soil, air and water are shown in  
5 log<sub>10</sub> scale in Figures SI 8.1 to SI 8.8, differentiating min, max and generic values across  
6 continents and archetypes for air emissions.


7 *Those figures have to be interpreted with care: the population density archetype is meaningful*  
8 *to assess toxic impacts of airborne emissions, but are less relevant for emissions into water or*  
9 *soil and in general for ecotoxicity impact category. It is likely that intercontinental spatial*  
10 *variability presented here underestimates the spatial variability within each continent.*




11




1




2  
3




1



2  
3




1  
2



3

1



2  
3  
4

## 1 References

---

- 2 Ackerman F, Stanton EA (2010) The social cost of carbon Economics for Equity and  
3 Environment (E3 Network) Available online at  
4 [http://www.e3network.org/papers/SocialCostOfCarbon\\_SEI\\_20100401.pdf](http://www.e3network.org/papers/SocialCostOfCarbon_SEI_20100401.pdf)  
5 Azevedo LB, De Schryver AM, Hendriks AJ, Huijbregts MA (2015) Calcifying species  
6 sensitivity distributions for ocean acidification Environ Sci Technol 49:1495-1500  
7 Boulay A-M et al. (2016) Consensus-based water scarcity footprint method from WULCA:  
8 the AWARE model Environ Sci Technol submitted  
9 Boulay AM, Bulle, C., Bayart, J.B., Deschenes, L., Margni, M (2011) Regional characterization  
10 of freshwater use in LCA: Modeling direct impacts on human health Environ Sci  
11 Technol 5:8948-8957  
12 Cao V, Margni M, Favis BD, Deschênes L (2015) Aggregated indicator to assess land use  
13 impacts in life cycle assessment (LCA) based on the economic value of ecosystem  
14 services J Clean Prod 94:56-66  
15 Cosme N, Hauschild MZ (2016) Effect factors for marine eutrophication in LCIA based on  
16 species sensitivity to hypoxia Ecol Indic 69:453-462  
17 Curran M et al. (2010) Toward meaningful end points of biodiversity in life cycle  
18 assessment† Environ Sci Technol 45:70-79  
19 de Baan L, Alkemade R, Koellner T (2013) Land use impacts on biodiversity in LCA: a global  
20 approach Int J Life Cycle Assess 18:1216-1230 doi:10.1007/s11367-012-0412-0  
21 de Bruille V (2014) Impact de l'utilisation des ressources minérales et métalliques dans un  
22 contexte cycle de vie : une approche fonctionnelle. Polytechnique Montreal  
23 De Schryver A, Brakkee K, Goedkoop M, Huijbregts M (2009) Characterization factors for  
24 global warming in life cycle assessment based on damages to humans and  
25 ecosystems Environ Sci Technol 43:1689-1695  
26 Fantke P, Charles R, de Alencastro LF, Friedrich R, Jolliet O (2011) Plant uptake of pesticides  
27 and human health: dynamic modeling of residues in wheat and ingestion intake  
28 Chemosphere 85:1639-1647  
29 Fantke P, Jolliet O (2016) Life cycle human health impacts of 875 pesticides Int J Life Cycle  
30 Assess 21:722-733  
31 Fantke P et al. (2015) Health effects of fine particulate matter in life cycle impact  
32 assessment: findings from the Basel Guidance Workshop Int J Life Cycle Assess  
33 20:276-288  
34 Fantke P, Wieland P, Juraske R, Shaddick G, Itoiz ES, Friedrich R, Jolliet O (2012)  
35 Parameterization models for pesticide exposure via crop consumption Environ Sci  
36 Technol 46:12864-12872  
37 Fatemi F (2012) A novel methodology for the assessment of the direct and indirect impacts  
38 associated with the depletion of fossil resources in life cycle assessment.  
39 Polytechnique Montreal  
40ecoinvent database (2003) <http://www.ecoinvent.ch/>.  
41 Frischknecht R, Braunschweig A, Hofstetter P, Suter P (2000) Human health damages due to  
42 ionising radiation in life cycle impact assessment Environ Impact Assess Rev 20:159-  
43 189  
44 Garnier-Laplace J et al. (2008) Issues and practices in the use of effects data from  
45 FREDERICA in the ERICA Integrated Approach J Environ Radioact 99:1474-1483  
46 doi:10.1016/j.jenvrad.2008.04.012



- 1 Goedkoop M, Heijungs R, Huijbregts M, De Schryver A, Struijs J, Van Zelm R (2013) ReCiPe  
2 2008, A life cycle impact assessment method which comprises harmonised category  
3 indicators at the midpoint and the endpoint level. First edition (version 1.08). Report  
4 I: Characterisation. The Hague, Ministry of Housing, Spatial Planning and  
5 Environment.,
- 6 Gronlund CJ, Humbert S, Shaked S, O'Neill MS, Jolliet O (2015) Characterizing the burden of  
7 disease of particulate matter for life cycle impact assessment *Air Qual Atmos* 8:29-46
- 8 Guinée JB et al. (2002) Handbook on life cycle assessment. Operational guide to the ISO  
9 standards. I: LCA in perspective. IIa: Guide. IIb: Operational annex. III: Scientific  
10 background. Kluwer Academic Publishers, Dordrecht
- 11 Hanafiah M, Xenopoulos M, Pfister S, Leuven RS, Huijbregts M (2011) Characterization  
12 factors for water consumption and greenhouse gas emissions based on freshwater  
13 fish species extinction *Environ Sci Technol* 45:5272-5278
- 14 Hauschild M et al. (2008) Building a model based on scientific consensus for life cycle  
15 impact assessment of chemicals: the search for harmony and parsimony *Environ Sci*  
16 *Technol* 42:7032-7037
- 17 Hauschild M, Potting J (2005) Spatial differentiation in life cycle impact assessment: The  
18 EDIP2003 methodology *Environmental News* 80
- 19 Hauschild MZ, Wenzel H (1998) Environmental assessment of products. Vol. 2 - Scientific  
20 background. Kluwer Academic Publishers, Hingham, MA. USA
- 21 Hellweg S, Demou E, Bruzzi R, Meijer A, Rosenbaum RK, Huijbregts MA, McKone TE (2009)  
22 Integrating human indoor air pollutant exposure within life cycle impact assessment  
23 *Environ Sci Technol* 43:1670-1679
- 24 Helmes RJ, Huijbregts MA, Henderson AD, Jolliet O (2012) Spatially explicit fate factors of  
25 phosphorous emissions to freshwater at the global scale *Int J Life Cycle Assess*  
26 17:646-654
- 27 Huijbregts MAJ, Rombouts LJA, Ragas AMJ, van de Meent D (2005) Human-toxicological  
28 effect and damage factors of carcinogenic and noncarcinogenic chemicals for life  
29 cycle impact assessment *Integr Environ Assess Manage* 1:181-244
- 30 Humbert S, Maendly R (2009) Characterization factors for damage to aquatic biodiversity  
31 caused by water use especially from dams used for hydropower *LCA IX*, Boston
- 32 Humbert S et al. (2011) Intake fraction for particulate matter: recommendations for life  
33 cycle impact assessment *Environ Sci Technol* 45:4808-4816
- 34 Jolliet O, Margni M, Charles R, Humbert S, Payet J, Rebitzer G, Rosenbaum R (2003a)  
35 IMPACT 2002+: A New Life Cycle Impact Assessment Methodology *Int J Life Cycle*  
36 *Assess* 8:324-330
- 37 Joos F et al. (2013) Carbon dioxide and climate impulse response functions for the  
38 computation of greenhouse gas metrics: a multi-model analysis *Atmos Chem Phys*  
39 13:2793-2825
- 40 Kounina A, Margni M, Shaked S, Bulle C, Jolliet O (2014) Spatial analysis of toxic emissions  
41 in LCA: A sub-continental nested USEtox model with freshwater archetypes *Environ*  
42 *Int* 69:67-89
- 43 Margni M et al. (2008) Guidance on how to move from current practice to recommended  
44 practice in Life Cycle Impact Assessment. UNEP-SETAC Life Cycle Initiative,  
45 Mathers C, Doris MF, Jan TB (2008) The global burden of disease: 2004 update.

- 1 Muñoz I, Rigarlsford G, i Canals LM, King H (2013) Accounting for greenhouse gas emissions  
2 from the degradation of chemicals in the environment *Int J Life Cycle Assess* 18:252-  
3 262
- 4 Myhre G (2013) Anthropogenic and natural radiative forcing *Climate Change 2013: The*  
5 *Physical Science Basis. Contribution of Working Group I to the Fifth Assessment*  
6 *Report 576 of the Intergovernmental Panel on Climate Change.*
- 7 Rosenbaum R et al. (2008) USEtox—the UNEP-SETAC toxicity model: recommended  
8 characterisation factors for human toxicity and freshwater ecotoxicity in life cycle  
9 impact assessment *Int J Life Cycle Assess* 13:532-546
- 10 Rosenbaum RK et al. (2015) Indoor air pollutant exposure for life cycle assessment:  
11 regional health impact factors for households *Environ Sci Technol* 49:12823-12831
- 12 Roy P-O, Azevedo LB, Margni M, van Zelm R, Deschênes L, Huijbregts MA (2014)  
13 Characterization factors for terrestrial acidification at the global scale: A systematic  
14 analysis of spatial variability and uncertainty *Sci Total Environ* 500:270-276
- 15 Roy P-O, Deschênes L, Margni M (2012a) Life cycle impact assessment of terrestrial  
16 acidification: modeling spatially explicit soil sensitivity at the global scale *Environ*  
17 *Sci Technol* 46:8270-8278
- 18 Roy P-O, Huijbregts M, Deschênes L, Margni M (2012b) Spatially-differentiated atmospheric  
19 source–receptor relationships for nitrogen oxides, sulfur oxides and ammonia  
20 emissions at the global scale for life cycle impact assessment *Atmos Environ* 62:74-  
21 81
- 22 Saad R, Margni M, Koellner T, Wittstock B, Deschênes L (2011) Assessment of land use  
23 impacts on soil ecological functions: development of spatially differentiated  
24 characterization factors within a Canadian context *Int J Life Cycle Assess* 16:198-211
- 25 Struijs J, van Dijk A, Slaper H, van Wijnen HJ, Velders GJ, Chaplin G, Huijbregts MA (2009)  
26 Spatial-and time-explicit human damage modeling of ozone depleting substances in  
27 life cycle impact assessment *Environ Sci Technol* 44:204-209
- 28 Thomas C et al. (2004) Extinction risk from climate change *Nature* 427:145-148
- 29 Tirado-Seco P (2005) Development of damage functions for aquatic eutrophication in *Life*  
30 *Cycle Assessment. Université de Genève*
- 31 van Zelm R, Schipper A, Rombouts M, Snepvangers J, Huijbregts M (2011) Implementing  
32 Groundwater Extraction in Life Cycle Impact Assessment: Characterization Factors  
33 Based on Plant Species Richness for the Netherlands *Environ Sci Technol* 45:629-  
34 635
- 35 Verones F, Hanafiah MM, Pfister S, Huijbregts MAJ, Pelletier GJ, Koehler A (2010)  
36 Characterization Factors for Thermal Pollution in Freshwater Aquatic Environments  
37 *Environ Sci Technol* 44:9364-9369
- 38 Wenger Y, Li D, Jolliet O (2012) Indoor intake fraction considering surface sorption of air  
39 organic compounds for life cycle assessment *Int J Life Cycle Assess* 17:919-931
- 40 WHO. (2003) *Climate change and human health: Risks and Responses.* World Health  
41 Organization, Geneva
- 42 WMO (World Meteorological Organization), 88 pp. (2014) *Assessment for Decision-Makers:*  
43 *Scientific Assessment of Ozone Depletion - Global Ozone Research and Monitoring*  
44 *Project—Report No. 56.* WMO (World Meteorological Organization), Geneva,  
45 Switzerland
- 46  
47

1 Ackerman F, Stanton EA (2010) The social cost of carbon Economics for Equity and  
2 Environment (E3 Network) Available online at [http://www](http://www.e3network.org/papers/SocialCostOfCarbon_SEI_20100401.pdf)  
3 [e3network](http://www.e3network.org/papers/SocialCostOfCarbon_SEI_20100401.pdf)  
4 Azevedo LB, De Schryver AM, Hendriks AJ, Huijbregts MA (2015) Calcifying species  
5 sensitivity distributions for ocean acidification Environmental Science & Technology  
6 49:1495-1500  
7 Barnett J, Adger WN (2007) Climate change, human security and violent conflict Political  
8 Geography 26:639-655  
9 Bennett DH, McKone TE, Evans JS, Nazaroff WW, Margni MD, Jolliet O, Smith KR (2002)  
10 Defining intake fraction Environmental Science & Technology 36:207A-211A  
11 Boulay A-M et al. (2016) Consensus-based water scarcity footprint method from WULCA:  
12 the AWARE model Environmental Science & Technology submitted  
13 Boulay AM, Bouchard C, Bulle C, Deschênes L, Margni M (2011) Categorizing water for LCA  
14 inventory The International Journal of Life Cycle Assessment 16:639-651  
15 Cao V, Margni M, Favis BD, Deschênes L (2015) Aggregated indicator to assess land use  
16 impacts in life cycle assessment (LCA) based on the economic value of ecosystem  
17 services Journal of Cleaner Production 94:56-66  
18 Chaudhary A, Verones F, de Baan L, Hellweg S (2015) Quantifying Land Use Impacts on  
19 Biodiversity: Combining Species–Area Models and Vulnerability Indicators  
20 Environmental Science & Technology 49:9987-9995 doi:10.1021/acs.est.5b02507  
21 Cosme N, Hauschild MZ (2016) Effect factors for marine eutrophication in LCIA based on  
22 species sensitivity to hypoxia Ecological Indicators 69:453-462  
23 Curran M et al. (2010) Toward meaningful end points of biodiversity in life cycle  
24 assessment† Environmental Science & Technology 45:70-79  
25 de Baan L, Alkemade R, Koellner T (2013) Land use impacts on biodiversity in LCA: a global  
26 approach. The International Journal of Life Cycle Assessment 18:1216-1230  
27 doi:10.1007/s11367-012-0412-0  
28 de Bruille V (2014) Impact de l'utilisation des ressources minérales et métalliques dans un  
29 contexte cycle de vie : une approche fonctionnelle. Polytechnique Montreal  
30 De Schryver A, Brakkee K, Goedkoop M, Huijbregts M (2009) Characterization factors for  
31 global warming in life cycle assessment based on damages to humans and  
32 ecosystems Environmental science & technology 43:1689-1695  
33 European Commission (2011) International Reference Life Cycle Data System (ILCD)  
34 Handbook- Recommendations for Life Cycle Impact Assessment in the European  
35 context., First edition edn. European commission - Joint Research Centre - Institute  
36 for Environment and Sustainability, Luxemburg  
37 Fantke P, Charles R, de Alencastro LF, Friedrich R, Jolliet O (2011) Plant uptake of pesticides  
38 and human health: dynamic modeling of residues in wheat and ingestion intake  
39 Chemosphere 85:1639-1647  
40 Fantke P, Jolliet O (2016) Life cycle human health impacts of 875 pesticides The  
41 International Journal of Life Cycle Assessment 21:722-733  
42 Fantke P et al. (2015) Health effects of fine particulate matter in life cycle impact  
43 assessment: findings from the Basel Guidance Workshop The International Journal  
44 of Life Cycle Assessment 20:276-288  
45 Fantke P, Wieland P, Juraske R, Shaddick G, Itoiz ES, Friedrich R, Jolliet O (2012)  
46 Parameterization models for pesticide exposure via crop consumption  
47 Environmental Science & Technology 46:12864-12872

1 Fatemi F (2012) A novel methodology for the assessment of the direct and indirect impacts  
2 associated with the depletion of fossil resources in life cycle assessment.  
3 Polytechnique Montreal  
4 ecoinvent database (2003) <http://www.ecoinvent.ch/>.  
5 Frischknecht R, Braunschweig A, Hofstetter P, Suter P (2000) Human health damages due to  
6 ionising radiation in life cycle impact assessment Environmental impact assessment  
7 Review 20:159-189  
8 Frischknecht R et al. (2016) Global guidance on environmental life cycle impact assessment  
9 indicators: progress and case study The International Journal of Life Cycle  
10 Assessment 21:429-442  
11 Garnier-Laplace J et al. (2008) Issues and practices in the use of effects data from  
12 FREDERICA in the ERICA Integrated Approach J Environ Radioact 99:1474-1483  
13 doi:10.1016/j.jenvrad.2008.04.012  
14 Goedkoop M, Heijungs R, Huijbregts M, De Schryver A, Struijs J, Van Zelm R (2013) ReCiPe  
15 2008, A life cycle impact assessment method which comprises harmonised category  
16 indicators at the midpoint and the endpoint level. First edition (version 1.08). Report  
17 I: Characterisation. The Hague, Ministry of Housing, Spatial Planning and  
18 Environment,  
19 Gronlund CJ, Humbert S, Shaked S, O'Neill MS, Jolliet O (2015) Characterizing the burden of  
20 disease of particulate matter for life cycle impact assessment Air Quality,  
21 Atmosphere & Health 8:29-46  
22 Guinée JB et al. (2002) Handbook on life cycle assessment. Operational guide to the ISO  
23 standards. I: LCA in perspective. IIa: Guide. IIb: Operational annex. III: Scientific  
24 background. Kluwer Academic Publishers, Dordrecht  
25 Hanafiah M, Xenopoulos M, Pfister S, Leuven RS, Huijbregts M (2011) Characterization  
26 factors for water consumption and greenhouse gas emissions based on freshwater  
27 fish species extinction Environment Science and Technology 45:5272-5278  
28 Hauschild M et al. (2008) Building a model based on scientific consensus for life cycle  
29 impact assessment of chemicals: the search for harmony and parsimony  
30 Environmental Science & Technology 42:7032-7037  
31 Hauschild M, Potting J (2005) Spatial differentiation in life cycle impact assessment: The  
32 EDIP2003 methodology. Environmental News 80  
33 Hauschild MZ, Wenzel H (1998) Environmental assessment of products. Vol. 2 - Scientific  
34 background. Kluwer Academic Publishers, Hingham, MA. USA  
35 Hellweg S, Demou E, Bruzzi R, Meijer A, Rosenbaum RK, Huijbregts MA, McKone TE (2009)  
36 Integrating human indoor air pollutant exposure within life cycle impact assessment  
37 Environmental Science & Technology 43:1670-1679  
38 Helmes RJ, Huijbregts MA, Henderson AD, Jolliet O (2012) Spatially explicit fate factors of  
39 phosphorous emissions to freshwater at the global scale The International Journal of  
40 Life Cycle Assessment 17:646-654  
41 Hodas N et al. (2015) Indoor inhalation intake fractions of fine particulate matter: Review  
42 of influencing factors Indoor Air  
43 Huijbregts MAJ, Rombouts LJA, Ragas AMJ, van de Meent D (2005) Human-toxicological  
44 effect and damage factors of carcinogenic and noncarcinogenic chemicals for life  
45 cycle impact assessment Integrated Environmental Assessment and Management  
46 1:181-244

1 Humbert S, Maendly R (2009) Characterization factors for damage to aquatic biodiversity  
2 caused by water use especially from dams used for hydropower LCA IX, Boston 1  
3 Humbert S et al. (2011) Intake fraction for particulate matter: recommendations for life  
4 cycle impact assessment Environmental Science & Technology 45:4808-4816  
5 International disaster database (2009).  
6 ISO (2013) ISO 3166-2 - Codes for the representation of names of countries and their  
7 subdivisions -- Part 2: Country subdivision code. International Organization for  
8 Standardization,  
9 Jolliet O, Margni M, Charles R, Humbert S, Payet J, Rebitzer G, Rosenbaum R (2003a)  
10 IMPACT 2002+: A New Life Cycle Impact Assessment Methodology The International  
11 Journal of Life Cycle Assessment 8:324-330  
12 Jolliet O, Margni M, Charles R, Humbert S, Payet J, Rebitzer G, Rosenbaum R (2003b)  
13 IMPACT 2002+: A New Life Cycle Impact Assessment Methodology International  
14 journal of LCA 8:324-330  
15 Joos F et al. (2013) Carbon dioxide and climate impulse response functions for the  
16 computation of greenhouse gas metrics: a multi-model analysis Atmospheric  
17 Chemistry and Physics 13:2793-2825  
18 Kounina A et al. (2013) Review of methods addressing freshwater use in life cycle inventory  
19 and impact assessment The International Journal of Life Cycle Assessment 18:707-  
20 721 doi:doi:10.1007/s11367-012-0519-3  
21 Kounina A, Margni M, Shaked S, Bulle C, Jolliet O (2014) Spatial analysis of toxic emissions  
22 in LCA: A sub-continental nested USEtox model with freshwater archetypes  
23 Environment International 69:67-89  
24 Levasseur A, de Schryver A, Hauschild M, Kabe Y, Sahnoune A, Tanaka K, Cherubini F (2016)  
25 Greenhouse gas emissions and climate change impacts. In: Frischknecht R, Jolliet O  
26 (eds) Global Guidance for Life Cycle Impact Assessment Indicators, vol 1. United  
27 Nations Environment Programme,  
28 Margni M et al. (2008) Guidance on how to move from current practice to recommended  
29 practice in Life Cycle Impact Assessment. UNEP-SETAC Life Cycle Initiative,  
30 Mathers C, Doris MF, Jan TB (2008) *The global burden of disease: 2004 update*.  
31 Muñoz I, Rigarlsford G, i Canals LM, King H (2013) Accounting for greenhouse gas emissions  
32 from the degradation of chemicals in the environment The International Journal of  
33 Life Cycle Assessment 18:252-262  
34 Myhre G (2013) Anthropogenic and natural radiative forcing Climate Change 2013: The  
35 Physical Science Basis. Contribution of Working Group I to the Fifth Assessment  
36 Report 576 of the Intergovernmental Panel on Climate Change.  
37 Nriagu J (2014) Expert judgement based estimation of global human exposure to toxic zinc  
38 levels.  
39 Pfister S, Koehler A, Hellweg S (2009) Assessing the environmental impacts of freshwater  
40 consumption in LCA Environmental Science & Technology 43:4098-4104  
41 Rosenbaum R et al. (2008) USEtox—the UNEP-SETAC toxicity model: recommended  
42 characterisation factors for human toxicity and freshwater ecotoxicity in life cycle  
43 impact assessment The International Journal of Life Cycle Assessment 13:532-546  
44 Rosenbaum R et al. (2011) USEtox human exposure and toxicity factors for comparative  
45 assessment of toxic emissions in life cycle analysis: sensitivity to key chemical  
46 properties The International Journal of Life Cycle Assessment 16:710-727

1 Rosenbaum RK et al. (2015) Indoor air pollutant exposure for life cycle assessment:  
2 regional health impact factors for households *Environmental Science & Technology*  
3 49:12823-12831

4 Roy P-O, Azevedo LB, Margni M, van Zelm R, Deschênes L, Huijbregts MA (2014)  
5 Characterization factors for terrestrial acidification at the global scale: A systematic  
6 analysis of spatial variability and uncertainty *Science of the Total Environment*  
7 500:270-276

8 Roy P-O, Deschênes L, Margni M (2012a) Life cycle impact assessment of terrestrial  
9 acidification: modeling spatially explicit soil sensitivity at the global scale  
10 *Environmental Science & Technology* 46:8270-8278

11 Roy P-O, Huijbregts M, Deschênes L, Margni M (2012b) Spatially-differentiated atmospheric  
12 source–receptor relationships for nitrogen oxides, sulfur oxides and ammonia  
13 emissions at the global scale for life cycle impact assessment *Atmospheric*  
14 *Environment* 62:74-81

15 Saad R, Margni M, Koellner T, Wittstock B, Deschênes L (2011) Assessment of land use  
16 impacts on soil ecological functions: development of spatially differentiated  
17 characterization factors within a Canadian context. *The International Journal of Life*  
18 *Cycle Assessment* 16:198-211

19 Struijs J, van Dijk A, Slaper H, van Wijnen HJ, Velders GJ, Chaplin G, Huijbregts MA (2009)  
20 Spatial-and time-explicit human damage modeling of ozone depleting substances in  
21 life cycle impact assessment *Environmental science & technology* 44:204-209

22 Thomas C et al. (2004) Extinction risk from climate change *Nature* 427:145-148

23 Tirado-Seco P (2005) Development of damage functions for aquatic eutrophication in *Life*  
24 *Cycle Assessment*. Université de Genève

25 van Zelm R, Schipper A, Rombouts M, Snepvangers J, Huijbregts M (2011) Implementing  
26 Groundwater Extraction in Life Cycle Impact Assessment: Characterization Factors  
27 Based on Plant Species Richness for the Netherlands *Environmental Science &*  
28 *Technology* 45:629-635

29 Verones F, Hanafiah MM, Pfister S, Huijbregts MAJ, Pelletier GJ, Koehler A (2010)  
30 Characterization Factors for Thermal Pollution in Freshwater Aquatic Environments  
31 *Environmental Science & Technology* 44:9364-9369

32 Wenger Y, Li D, Jolliet O (2012) Indoor intake fraction considering surface sorption of air  
33 organic compounds for life cycle assessment *The International Journal of Life Cycle*  
34 *Assessment* 17:919-931

35 WHO. (2003) *Climate change and human health: Risks and Responses*. World Health  
36 Organization, Geneva

37 WMO (World Meteorological Organization), 88 pp., , 2014. (2014) *Assessment for*  
38 *Decision-Makers: Scientific Assessment of Ozone Depletion - Global Ozone Research*  
39 *and Monitoring Project—Report No. 56*. WMO (World Meteorological Organization),  
40 Geneva, Switzerland

41