



**HAL**  
open science

## **A Review of Pesticides Sorption in Biochar from Maize, Rice, and Wheat Residues: current status and challenges for soil application**

Allan Pretti Ogura, Jacqueline Zanin Lima, Jéssica Pelinsom Marques, Lucas Massaro Sousa, Valéria Guimarães Silvestre Rodrigues, Evaldo Luiz Gaeta Espíndola

### ► To cite this version:

Allan Pretti Ogura, Jacqueline Zanin Lima, Jéssica Pelinsom Marques, Lucas Massaro Sousa, Valéria Guimarães Silvestre Rodrigues, et al.. A Review of Pesticides Sorption in Biochar from Maize, Rice, and Wheat Residues: current status and challenges for soil application. *Journal of Environmental Management*, 2021, 300, pp.113753. 10.1016/j.jenvman.2021.113753 . hal-03462367

**HAL Id: hal-03462367**

**<https://ifp.hal.science/hal-03462367v1>**

Submitted on 1 Dec 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **A review of pesticides sorption in biochar from maize, rice, and wheat residues:**

2 **current status and challenges for soil application**

3 Allan Pretti Ogura <sup>1\*</sup>, Jacqueline Zanin Lima <sup>2</sup>, Jéssica Pelinsom Marques <sup>2</sup>, Lucas Massaro

4 Sousa <sup>3</sup>, Valéria Guimarães Silvestre Rodrigues <sup>2</sup>, and Evaldo Luiz Gaeta Espíndola <sup>1</sup>

5 <sup>1</sup> PPG-SEA and NEEA/CRHEA/SHS, São Carlos School of Engineering, University of São

6 Paulo, São Carlos, SP, 13560-970, Brazil

7 <sup>2</sup> Department of Geotechnical Engineering, São Carlos School of Engineering, University of

8 São Paulo, São Carlos, SP, 13560-970, Brazil

9 <sup>3</sup> Process Design and Modeling Division, IFP Energies Nouvelles, Rond-Point Échangeur de

10 Solaize, 69360, Solaize, France

11  
12 \* Corresponding author

13 Allan Pretti Ogura

14 ORCID 0000-0001-8549-5820

15 e-mail: allan.ogura@usp.br

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 23 ABSTRACT

1  
2  
3 24 The use of pesticides has been increasing in recent years for maintaining traditional agricultural  
4  
5 25 practices. However, these chemicals are associated with several environmental impacts,  
6  
7  
8 26 demanding urgent remediation techniques. Biochar is a carbonaceous material produced by  
9  
10 27 pyrolysis that has the potential for pesticide sorption and remediation. In this context, this  
11  
12  
13 28 interdisciplinary review systematically assessed the state of the knowledge of crop residues to  
14  
15 29 produce biochar for pesticide sorption. We focused on maize, rice, and wheat residues since  
16  
17  
18 30 these are the three most-produced grains worldwide. Besides, we evaluated different biochar  
19  
20 31 handling, storage, and soil dispersion techniques to ease its implementation in agriculture. In  
21  
22 32 general, pyrolysis temperature influences biochar characteristics and its potential for pesticide  
23  
24  
25 33 sorption. Furthermore, biochar amended soils had greater pesticide sorption capacity, limiting  
26  
27 34 potential leaching and runoff. Most studies showed that the feedstock and specific surface area  
28  
29  
30 35 influence the biochar sorption properties, among other factors. Also, biochar reduces  
31  
32 36 pesticides' bioavailability, decreasing their toxicity to soil organisms and improving soil  
33  
34  
35 37 fertility and crop yields. Nonetheless, the retrieved papers assessed only 21 pesticides, mainly  
36  
37 38 consisting of lab-scale batch experiments. Therefore, there is still a gap in studies evaluating  
38  
39  
40 39 biochar aging, its potential desorption, pesticide co-contaminations, the associated  
41  
42 40 microbiological processes, and field applications. Determining flow properties for biochars of  
43  
44 41 different sizes and pellets is vital for reliable handling equipment design, and performing  
45  
46 42 techno-economic assessment under different farm contexts is encouraged. Ultimately, coupling  
47  
48  
49 43 biochar production with residue management could address this challenge on sustainable  
50  
51  
52 44 agricultural systems.

53  
54  
55 45 **KEYWORDS:** Biochar amendment, Biochar handling, Crop residues, Contamination, Soil  
56  
57 46 remediation, Waste management.  
58  
59  
60  
61  
62  
63  
64  
65

## 47 1. Introduction

48 Synthetic chemicals used in conventional agriculture have increased production by controlling  
49 pests, weeds, and diseases. Pesticides target living species, as they are biologically active  
50 substances with a structure capable of imitating specific molecules (Das, 2013; Rani et al.,  
51 2021). Over the past decade, pesticide use surpassed 4 million tons per year worldwide (FAO,  
52 2020). However, pesticides' extensive use has drastically affected the soil, water, and air  
53 quality, causing undesirable impacts (e.g., toxicity, carcinogenicity, and mutagenicity) on non-  
54 target organisms, including humans (Khalid et al., 2020; Rani et al., 2021; Varjani et al., 2019).  
55 In agricultural systems, pesticide contamination can negatively affect soil quality and crop  
56 production, jeopardizing ecosystem services, nutrient cycling, enzyme activity, soil biota, and  
57 biodiversity (Liu et al., 2018; Yu et al., 2019). Pesticides can also reach the surface and  
58 groundwater through diffuse pollution (e.g., run-off and leaching) (Damalas and  
59 Eleftherohorinos, 2011; Khalid et al., 2020).

60 In this scenario, research trends focused on different approaches to mitigate pesticide risks in  
61 soil and water, including chemical remediation, containment or immobilization, and  
62 bioremediation (Ganie et al., 2021; Morillo and Villaverde, 2017; Rani et al., 2021; Saleh et  
63 al., 2020). However, biochar stands out as a cost-effective and environment-friendly alternative  
64 for remediation due to its sorption potential (Kwon et al., 2020; Lehmann and Joseph, 2015;  
65 Liu et al., 2018; Varjani et al., 2019). This carbon-enriched product results from the thermal  
66 decomposition of biomass or organic material, including crop residues (Ali et al., 2019; N. Liu  
67 et al., 2015), animal manure (N. Liu et al., 2015; Ren et al., 2018), woody materials (You et  
68 al., 2020; Zhu et al., 2020), and biosolids (Ali et al., 2019; Regkouzas and Diamadopoulos,  
69 2019). The pyrolysis process generally occurs at high temperatures (between 350 and 1200 °C)  
70 and in an oxygen-limited environment (Lehmann and Joseph, 2015). Biochar application

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

71 contributes to the immobilization of organic and inorganic pollutants due to its large pore  
72 structure (micro and mesoporous), rich surface functional groups (e.g., carbonyl, hydroxyl,  
73 phenolic hydroxyl, and carboxyl), large specific surface area (SSA), high pH, and high cation  
74 exchange capacity (CEC) (Beesley et al., 2011; Liu et al., 2018; Waqas et al., 2020; Yu et al.,  
75 2019). Several studies have demonstrated the potential sorption of various pesticides (e.g.,  
76 atrazine, fipronil, diuron, 1,3-D, and 2,4-D) onto biochar solutions (Y. Liu et al., 2015;  
77 Regkouzas and Diamadopoulou, 2019; Zhu et al., 2020), and soils amended with biochar (Ren  
78 et al., 2018; Wu et al., 2019; Yavari et al., 2020; You et al., 2020).

79 Biochar amendments on agricultural soils can reduce pesticide mobility, transport, and  
80 bioavailability or microbial uptake (Yu et al., 2019). Moreover, it can stimulate soil microbiota  
81 while enhancing pesticide degradation (Varjani et al., 2019; Waqas et al., 2020), increase  
82 nutrient content and water-holding capacity, ameliorate acidic soils and aeration properties,  
83 improving soil fertility and crop yields (Khalid et al., 2020; Liu et al., 2018; Palansooriya et  
84 al., 2019). Thus, biochar amendments minimize the environmental risks of pesticides in soil  
85 and water.

86 Millions of tons of agricultural waste are produced globally every year from various crop  
87 cultivation and processing (Duque-Acevedo et al., 2020). Thus, crop residues are potential low-  
88 cost feedstocks for biochar production and an alternative for agricultural waste management.  
89 This approach has sustainable and economic benefits by converting wastes into value-added  
90 products. Cereals are the most important staple foods for humans and animals worldwide  
91 (FAO, 2015), with maize, rice, and wheat figuring among the leading global productions (FAO,  
92 2019); hence, biochar production from these crop residues is of great interest.

93 With increasing publications on biochar, some review articles summarized the aspects of  
94 biochar production and applications (Khalid et al., 2020; Kwon et al., 2020; Li et al., 2019; Liu

1 95 et al., 2018; Palansooriya et al., 2019; Varjani et al., 2019; Waqas et al., 2020; Wei et al., 2018;  
2 96 Yaashikaa et al., 2019). However, these reviews focused mainly on biochar's physicochemical  
3  
4 97 properties, its influence on soil fertility, removing metals on laboratory experiments or using  
5  
6  
7 98 biochar in aqueous solutions with pesticides. Despite this, few papers highlighted recent trends  
8  
9  
10 99 and implications of using biochar from crop residues on pesticide-contaminated soils (Khalid,  
11  
12 100 2019; Varjani et al., 2019). Therefore, given the potential of crop residues-based biochar for  
13  
14 101 amendment of pesticides-contaminated agricultural soils, it would be of great interest to have:  
15  
16  
17 102 i) an up-to-date systematic review on biochar amendment in soils, showing current state-of-  
18  
19 103 knowledge and future challenges and perspectives for its advancement; ii) a discussion and  
20  
21  
22 104 comparison regarding biochar studies from different crop residues as feedstocks, focusing on  
23  
24 105 the most produced ones (i.e., maize, rice, and wheat); iii) an interdisciplinary review that  
25  
26  
27 106 highlights and proposes solutions for typical biochar field application challenges, such as  
28  
29 107 equipment for proper transport, storage, and handling, besides methods for dispersion in farm  
30  
31 108 soils.

32  
33  
34  
35 109 In this scenario, this research's general objective was to systematically review the current  
36  
37 110 knowledge about crop residues as feedstocks to biochar production for pesticide sorption. The  
38  
39  
40 111 specific aims were to i) gather research data on agricultural residues (i.e., maize, rice, and wheat  
41  
42 112 crops) and pesticide use; ii) provide an overview of the main properties of crop residues-based  
43  
44  
45 113 biochar and their effects on the sorption of pesticides in the soil; iii) conclude the potential of  
46  
47 114 biochar amendments to improve the pesticide sorption capacity; iv) investigate some options  
48  
49  
50 115 for biochar handling, storage, and dispersion in soil applied to agriculture; v) establish limiting  
51  
52 116 factors, uncertainties, and gaps in current research while proposing future approaches.

53  
54  
55 117  
56  
57  
58  
59 118  
60  
61  
62  
63  
64  
65

## 119 2. Systematic review

1  
2  
3 120 The systematic review (Figure 1) performed in this study followed PRISMA guidelines (Moher  
4  
5  
6 121 et al., 2009), and used the databases from Web of Science, ScienceDirect, SpringerLink, Taylor  
7  
8 122 & Francis, and Scopus. The research gathered papers published between January 1<sup>st</sup>, 2009, and  
9  
10 123 January 17<sup>th</sup>, 2021. The adopted string was “biochar AND (adsorption OR sorption) AND  
11  
12 124 (water OR soil) AND (pesticide OR herbicide OR insecticide OR fungicide)”, considering only  
13  
14  
15 125 documents in English.

16  
17  
18 126 From the selected databases, we identified a total of 2746 papers and 2385 after removing  
19  
20  
21 127 duplicates. The first step consisted of retrieving the following information on these papers: title,  
22  
23 128 first author, publication year, database(s), author(s) country(ies), keywords, environmental  
24  
25  
26 129 matrix, biochar feedstock, and pesticide. The results were then screened on Microsoft Excel<sup>®</sup>.  
27  
28 130 There has been an increasing number of papers published recently, as shown in Figure 1. Most  
29  
30  
31 131 articles per year were observed in 2020 (n = 749), and 68 documents from 2021 were available  
32  
33 132 by January 17<sup>th</sup>.

34  
35  
36 133

37  
38  
39  
40 134

41  
42  
43 135

44  
45  
46 136

47  
48  
49  
50 137


51  
52  
53 138

54  
55  
56 139

57  
58  
59  
60 140

61  
62  
63  
64  
65

141 Figure 1 - Systematic review summary: search process and selection of the studies focused on maize, rice, and  
 142 wheat residues biochar for pesticide sorption.



143



144 The second step consisted of narrowing papers relevant to this study aim, which means biochar  
1 associated with pesticide sorption. Thus, non-experimental papers (e.g., review and  
2 145 simulations) were excluded, as well as research focusing on other sorbents (e.g., activated  
3 146 carbon and ash) and chemicals besides pesticides (e.g., pharmaceuticals and polychlorinated  
4 147 biphenyls). After, 310 papers were selected, which represents approximately 13% from the first  
5 148 step. A word map was created from the bibliometric extension on the Microsoft Word® to  
6 149 observe the most publishing countries and used keywords (Figure 2). China is the country that  
7 150 published the most (n = 145), followed by the USA (n = 65), and India (n = 29). Furthermore,  
8 151 the main used keywords were biochar (n = 207), sorption (n = 102), and soil (n = 82).  
9 152

153 By focusing on agricultural crop residues among different retrieved feedstocks, papers were  
154 reduced to 143. The papers were lastly screened to studies that i) focused on sorption; ii)  
155 focused on soil application or experimental research; iii) the biochar was produced from the  
156 pyrolysis, and iv) used rice, maize, wheat residues for biochar production as these are the most  
157 produced grains worldwide as detailed in Section 3 (FAO, 2019). A total of 32 papers were  
158 studied (Table 1), and the Supplementary Material presents full information regarding the  
159 biochar production, feedstock type, pyrolysis conditions, and whether pre- or post-treatment  
160 were considered. Biochar characterization is also reported: pH, CEC, SSA, particle sizes (PS),  
161 pore volume (PV), % ash, % C, % N, % H, and % of other elements. Other processes besides  
162 sorption are described (e.g., electrochemical and bioremediation) when evidenced by the  
163 authors. This paper reviewed the recovered studies (especially in Section 5), and the efficiency  
164 of pesticide sorption and soil characterization were gathered to compare and discuss the selected  
165 biochars.

166

167

168 Table 1 - Systematic review summary: studies from the final selection on pesticide sorption into crop residue  
 169 biochars.

Crop Feedstock	Pyrolysis	pH	CEC (cmol <sub>c</sub> kg <sup>-1</sup> )	SSA (m <sup>2</sup> g <sup>-1</sup> )	Studied pesticides	References	
Maize	corn cob	450 °C 4 h	n.a.	n.a.	1.5 - 356.0	atrazine	Ouyang (2016a)
	stalk	200 - 850 °C n.a.	9.4	n.a.	13.2 - 386.0	2,4-D, atrazine, flubendiamide	Clay (2016); Ouyang (2016a); Das and Mukherjee (2020); Tao (2020)
	straw	300 - 850 °C 4 - 8 h	6.8 - 10.6	459.0 - 468.0	1.7 - 196.0	1,3-D, atrazine, carbaryl, clothianidin, imidacloprid, oxyfluorfen, thiacloprid, topramezone	Graber (2011), Ouyang (2016b), Qin (2019), Ren (2016), Uwamungu (2019), Wu (2019), Zhang (2018, 2020)
Rice	hull	500 - 600 °C 3 - 4 h	6.7 - 9.6	n.a.	10.7 - 95.7	acetochlor, fomesafen, oxyfluorfen	Khorram (2015, 2017, 2018); Li (2018); Wu (2019)
	husk	300 - 700 °C 1 - 3 h	6.0 - 9.2	5.0 - 70.7	2.00 - 202.1	atrazine, diuron, imazapic, imazapyr, oxytetracycline	Aldana (2020); Yavari (2016, 2017a, 2017b, 2020)
	straw	300 - 700 °C 1 - 4 h	7.2 - 11.1	45.3 and 86.5	8.0 - 188.0	2,4-D, acetochlor, atrazine, carbaryl, fomesafen, pyrazosulfuron-ethyl	Lü (2012); Manna, Singh (2015); Ren (2016); Khorram (2018); Zhao (2019); Manna (2019)
Wheat	mids	550 °C n.a.	8.9	n.a.	24.7	1,3-D	Wang (2016)
	straw	250 - 600 °C 1 - 6 h	5.4 - 10.6	22.0 - 92.0	2.7 - 62.6	atrazine, chlorpyrifos, hexachlorobenzene, MCPA, pyrazosulfuron-ethyl, simazine	Song (2012, 2016); Tatarkova (2013); Manna and Singh (2015); Manna and Singh (2019); Wang (2016); Cheng (2017); Humera Aziz (2018); Ren (2018)

170 Notes: n.a. = not available

171

172 The recovered papers are further discussed in the following sections. First, primary crop  
173 residues to be used as feedstock for biochar production are presented (Section 3). Then, the  
174 pesticides from the retrieved papers are discussed, considering their primary characteristics  
175 influencing biochar sorption (Section 4). Next, the main factors that govern pesticide sorption  
176 are assessed based on the recovered studies, considering biochar properties and soil applications  
177 (Section 5.1). Besides, we focused on the studies addressing maize, rice, and wheat residues  
178 biochar (Section 5.2, 5.3, and 5.4, respectively). Further discussions are also provided for the  
179 retrieved studies with some key aspects, gaps, and uncertainties (Section 6). Next, we reviewed  
180 equipment and techniques for biochar transport, storage, and dispersion (Section 7), considering  
181 the lack of field studies to optimize soil application. Finally, a synthesis of the systematic review  
182 is presented, highlighting the advantages, limitations, and perspectives of pesticide sorption  
183 with residue-based biochar (Section 8).

184

### 185 **3. Crop residues as feedstock for biochar production**

186 Crop production generates large amounts of residues throughout its harvesting, including stalks,  
187 husks, and straws (Duque-Acevedo et al., 2020). Although residues are widely studied as  
188 feedstock for bioethanol (Kim and Dale, 2004), they could be used as soil amendments for  
189 reducing soil degradation (Lal, 2008). Therefore, the most appropriate crop residue uses should  
190 enhance, maintain, and sustain soil properties, increasing the soil organic carbon and  
191 minimizing soil pollution (Lal, 2005). Thus, residue management is crucial for achieving  
192 sustainable agricultural systems, recovering soil properties while contributing to carbon  
193 sequestration (Ding et al., 2017; Song et al., 2019).

194 The largest cereal crops worldwide are maize, wheat, and rice (FAO, 2020) (Figure SM1).

195 Maize (*Zea mays* L.) is an annual plant production, and the USA, China, and Brazil are the

196 biggest world producers; China and India are the major rice (*Oryza sativa* L.) producers; and  
197 wheat (*Triticum* spp.) cultivation is more prominent in China, India, and Russia (FAO, 2019).  
198 As of 2001, cereals corresponded to 74 % of the world's annual residue production, considering  
199 the estimated maize, rice, and wheat residue production of 609, 890, and 875 million Mg,  
200 respectively (Lal, 2005). This proportion indicates that maize, rice, and wheat produced 1148,  
201 1133, and 1148 million Mg residues in 2019, respectively (FAO, 2020). In terms of nitrogen,  
202 these three crops correspond to 75 % of the world's current residue production, considering  
203 9.36, 9.24, and 9.43 million Mg for maize, rice, and wheat, respectively (FAO, 2019). Maize  
204 residues (e.g., stalks) are commonly used for cattle livestock and tillage due to their high  
205 nutrient content (Villamil et al., 2015). On the other hand, rice straw has a high silica content,  
206 and it is not recommended for feeding animals (Agbagla-Dohnani et al., 2003). Rice hulls and  
207 straws can increase crop yields when incorporated into the soil (Sistani et al., 1998). Moreover,  
208 corn stover is a potential feedstock for bioethanol production (Kadam and McMillan, 2003),  
209 while corn cobs are not feasible for energy production (Erickson et al., 2011). Wheat straw and  
210 midds can be included in cattle feed and energy sources (Nguyen et al., 2013; ZoBell et al.,  
211 2005).

212

#### 213 **4. Studied pesticides uses and environmental concern**

214 Pesticides are widely applied to agricultural production, among other chemicals (i.e., fertilizers  
215 and pharmaceuticals) (L. Zhang et al., 2018). For instance, the commercial product consists of  
216 a combination of the active ingredient and other substances. Although the manufacturers'  
217 recommended dose varies according to each target species and crop, overdoses are mostly the  
218 cause of severe environmental effects (Meena et al., 2020). Asia and the Americas have a total  
219 pesticide use per area of cropland (3.67 and 3.52 kg ha<sup>-1</sup>, respectively) higher than the world's

220 average (2.63 kg ha<sup>-1</sup>) (FAO, 2020). Moreover, China, the USA, and Brazil are the leading  
221 pesticide consumers worldwide (FAO, 2020).

222 Understanding the pesticides' fate in the environment is essential. After application, these  
223 contaminants may leach and runoff and then reach aquatic habitats (Wang et al., 2019). These  
224 processes depend on the local climate conditions (Lammoglia et al., 2018), soil type (Vryzas,  
225 2018), and landscape features (Rouzies et al., 2019). Additionally, spray drift and accidental  
226 spills may increase their soil and water concentrations (Jemec et al., 2007). The soil type is  
227 crucial for pesticide's environmental behavior, and higher Koc (organic carbon-water partition  
228 coefficient) values indicate that pesticides are expected to have low mobility. In addition, the  
229 soil granulometry, pH, CEC, and carbon content can contribute to pesticide mobility  
230 (Lunagariya et al., 2020). Nonetheless, organic matter and clay particles increase some  
231 pesticides' sorption in soil (Sadegh-Zadeh et al., 2017; Spark and Swift, 2002). Although soil  
232 microbiota influences pesticide fate, biodegradation is not relevant when contaminants are not  
233 bioavailable (Scow and Johnson, 1996). Our research identified 21 pesticides in the final  
234 selection (Table 2), and information was presented concerning their density, solubility, log Kow  
235 (n-octanol-water partition coefficient), Koc, and half-life (Kim et al., 2021; USEPA, 2021).

236

237

238

239

240

241

242

Table 2. Summary of pesticides characteristics (adapted from Kim et al. 2021, and USEPA 2021)

Group	Pesticide	Density 20-25 °C (g cm <sup>-3</sup> )	Solubility 20-25 °C (mg L <sup>-1</sup> )	log Kow	Koc	Soil half-live (days)
Fungicide	hexachlorobenzene	1.21-2.04	0.0047	5.73	3.6-5.5	970-2100
	oxytetracycline	1.63	313	-0.90	195-93,317	>180
Herbicide	2,4-D	1.42	540-677	2.81	20-136	1.6-32.3
	acetochlor	1.11	223	4.14	98.5-335	7-203
	atrazine	1.23	33	2.61	26-1,164	1.4-108
	diuron	1.48	37.4-42	2.68	55.3-962	330
	fomesafen	1.28	50	2.90	34-1,200	21-360
	imazapic	0.24	220	0.39	3	31-410
	imazapyr	0.34	11,300	0.22	8.81	17.7-63.1
	MCPA	1.56	270,000	3.25	50-62	7-41
	oxyfluorfen	1.49	0.116	4.73	8,900	12
	pyrazosulfuron-ethyl	1.44	14.5	1.74	284.2	16-27
Insecticide	simazine	1.30	6.2	2.18	78-3,559	27-102
	topramezone	1.13	510	1.44	140	>125
	carbaryl	1.23	110	2.36	230-390	10
	chlorpyrifos	1.40	1.12-1.4	4.96	995-31,000	4-139
	clothianidin	1.61	327	0.70	60	148-1,155
	flubendiamide	1.66	0.0299	4.20	1,076-3,318	210-770
Nematicide	imidacloprid	1.54	610	0.57	156-800	34-190
	thiacloprid	1.46	185	1.26	1,100	0.6-3.8
	1,3-D	1.22	2,000-2,180	2.03	20-42	2-54

243

244 Most of the pesticides from the selected studies are herbicides (12 out of 21). Atrazine was the  
245 most studied pesticide on the retrieved papers (n = 80 after step 2, and n = 9 in the final  
246 selection) and the only pesticide on the top 10 keywords after step 2 (Figure 1). Atrazine has  
247 high to slight mobility in soil (Koc from 26 to 1164) and a half-life of more than 1,000 days  
248 (Kim et al., 2021). This herbicide modifies the growth, enzymatic processes, and photosynthesis  
249 in grassy and broadleaf plants (Singh et al., 2018). In addition, the potential atrazine sorption  
250 was investigated for several agricultural residues, including cassava waste (Deng et al., 2017;  
251 Li et al., 2018), peanut husk (Saha et al., 2017), sawdust (Gao et al., 2019), and sugarcane tops  
252 (Huang et al., 2018). Although glyphosate was one of the most studied pesticides after step 2  
253 (n = 15), several studies investigated the sorption on wood biochar (Hall et al., 2018; Junqueira  
254 et al., 2020). However, no paper in the final selection reported glyphosate sorption on biochar  
255 from our studied residues (i.e., maize, rice, and wheat).

256 Nonetheless, the sorption of other herbicides was studied for maize, rice, and wheat residues-  
257 based biochar. The herbicide 2,4-dichlorophenoxyacetic acid (2,4-D) is a synthetic auxin that  
258 targets broadleaf species, and it has high soil mobility (Koc from 20 to 136) and low persistence  
259 (1.6 to 32.3 days) (Gan et al., 2003). Acetochlor has high to moderate mobility in soil (Koc of  
260 98.5 to 335), while diuron sorption is strongly influenced by organic matter presence (Kim et  
261 al., 2021; Spurlock and Biggar, 1994). Fomesafen mobility depends on the soil type, with a  
262 wide range of Koc values (from 34 to 1200) (Kim et al., 2021). Imazapic and imazapyr are very  
263 mobile imidazolinone herbicides (Koc of 3 and 8.81, respectively), as the herbicide 2-methyl-  
264 4-chlorophenoxyacetic acid (MCPA) has high mobility (Koc from 50 to 62) and half-life from  
265 7 to 41 days (Kim et al., 2021). Pyrazosulfuron-ethyl has half-life values from 16 to 27 days in  
266 soils, and it is moderately sorbed in soils (Manna and Singh, 2015). Simazine has high to slight  
267 mobility (Koc from 78 to 3,559) with higher sorption under low pH, and topramezone has high  
268 mobility (Koc of 140) and low biodegradation potential (Kim et al., 2021).

269 Insecticides corresponded to 6 out of 21 pesticides from the selected studies. Carbaryl is a  
270 carbamate insecticide weakly sorbed and readily biodegradable in soil (half-life of 10 days)  
271 (Ahmad et al., 2004). Chlorpyrifos has low to no mobility in various soils (Koc of 995 to  
272 31,000), characterizing a strong sorption potential (Racke, 1993). Clothianidin has high soil  
273 mobility (Koc of 60) and persistence (half-life 148 to 1,155 days), a challenge for sorption  
274 techniques (Kim et al., 2021). Flubendiamide has slight to low soil mobility (Koc of 1,076 to  
275 3,318), and the half-life ranges from 210 to 770 days (Kim et al., 2021). For imidacloprid soil  
276 sorption, organic matter is more relevant than clay content and CEC (Liu et al., 2006).  
277 Thiacloprid has low mobility in soil (Koc of 1,100), and it is readily biodegradable (half-life of  
278 0.6 to 3.8 days) (Kim et al. 2020).

279 Two fungicides were found in the selected studies, hexachlorobenzene (HCB) and  
280 oxytetracycline (OTC). HCB is expected to be immobile on soils (Koc 3.6 to 5.5), it is  
281 considered a persistent organic pollutant (half-life from 970 to 2,100 days), and sorption might  
282 reduce its volatilization (Barber et al., 2005). OTC is an antimicrobial drug, also applied for  
283 fungi control, and its sorption depends mostly on soil texture, CEC, and iron oxide content  
284 (Jones et al., 2005). Although we did not include “nematicide” in our keywords, one study was  
285 found with 1,3-Dichloropropene (1,3-D). This pesticide is commonly applied with chloropicrin  
286 as an alternative to methyl bromide to control soilborne diseases and nematodes (Santos et al.,  
287 2006).

288 Residual pesticides can remain for long periods in the environment or agricultural origin  
289 products, posing a risk to human health (Kim et al., 2017) and establishing the maximum  
290 acceptable intake. Although these limits are less restrictive than those proposed for fauna and  
291 flora protection, the legislation is usually guided by human health protection. For this purpose,  
292 human exposure pathways are assessed and analyzed, including direct contact (e.g., dermal),  
293 ingestion (e.g., drinking water, food, and soil), and inhalation (e.g., soil particles and air).



294 However, most environmental agencies do not determine the maximum dose through all  
1  
2  
3 295 significant exposures, and some limits are still inefficient for health protection (Li and A.  
4  
5 296 Jennings, 2017). Also, pesticides might cause harmful effects to the biota even within limits  
6  
7 297 established by legislation and environmental agencies (Brovini et al., 2021).  
8  
9

10 298 Overall, pesticide pollution endangers the environment, considering adverse effects on water  
11  
12 299 quality, biodiversity, and human health, as highlighted by Tang et al. (2021). These authors  
13  
14 300 identified high-risk areas worldwide, and 34 % of them were in high-biodiversity regions, 5 %  
15  
16 301 in areas with water scarcity, and 19 % located in low and lower-middle-income nations. Also,  
17  
18 302 China was considered the top country susceptible to high pesticide pollution risk, emphasizing  
19  
20 303 the Huang He watershed. Since China is the leading country on studies of biochar application  
21  
22 304 for pesticide sorption, researchers and the government should perform these alternatives in  
23  
24 305 priority areas. Also, pesticides can cause effects on several non-target species (de Figueirêdo et  
25  
26 306 al., 2020; Sanches et al., 2017; Triques et al., 2021), and the ecotoxicological assessment of the  
27  
28 307 pesticides' impacts on representative species is a pathway to understanding their biota damages.  
29  
30 308 Moreover, the ecological risk of their presence in soil and water endangers the proper ecosystem  
31  
32 309 functioning and services (Schäfer et al., 2012; Schiesari and Grillitsch, 2011).  
33  
34  
35  
36  
37  
38  
39  
40  
41 310

## 42 43 311 **5. Biochar for pesticide-contaminated soils remediation**

44  
45  
46 312 Remediation techniques are applied to remove, reduce, degrade, or retain contaminants,  
47  
48 313 considering the ecological risks of pesticides in soils. The most appropriate method relies on  
49  
50 314 the environmental characterization of the area, financial resources, predicted execution time,  
51  
52 315 sustainability considerations, and remediation goals according to the future land use (Fernández  
53  
54 316 Rodríguez et al., 2014; Hou and Al-Tabbaa, 2014). Pesticide sorption processes stand out  
55  
56 317 among containment or immobilization technologies by limiting the contaminant potential  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 318 infiltration and leaching by applying carbonaceous materials as sorbents (Morillo and  
2 319 Villaverde, 2017).

3  
4  
5 320 Although biochar and activated carbon are carbon-rich solids with a similar porous structure,  
6  
7 321 the former has lower production and energy costs (Dai et al., 2019; Lehmann and Joseph, 2015).

8  
9  
10 322 Thus, biochar has been evaluated as an alternative to the commercial activated carbon for  
11  
12 323 contaminants sorption (Dai et al., 2019; Yaashikaa et al., 2019). The production of both sorbents

13  
14  
15 324 is through pyrolysis, but activated carbon can also be produced by chemical or steam activation  
16  
17 325 (Kazemi Shariat Panahi et al., 2020). Regarding the source material, activated carbon can come

18  
19  
20 326 from biomass or any carbonaceous substance (e.g., coal), although biochar is made exclusively  
21  
22 327 from biomass. Biochar as soil amendment aims to favor the contaminated or degraded soil's

23  
24  
25 328 physical, chemical, and biological properties (Kazemi Shariat Panahi et al., 2020). Biochar's  
26  
27 329 sorption capacity is not always as high as activated carbon (Liu et al., 2018), so the use of

28  
29  
30 330 engineered biochars resulting from activation and modification processes may be relevant in  
31  
32 331 some cases.

33  
34  
35  
36 332

### 37 38 39 333 *5.1. Primary factors for pesticide sorption on biochar*

40  
41  
42 334 The critical factors determining the biochar sorption efficiency are the feedstock, pyrolysis  
43  
44 335 temperature, solution pH, dosage of biochar, competition with co-existed ions and other

45  
46  
47 336 contaminants, the aging process, and modification/activation (Abbas et al., 2018). Higher  
48  
49 337 pyrolysis temperature increases the biochar's SSA and micropores, improving the organic

50  
51  
52 338 contaminants sorption (Abbas et al., 2018); this variation is related to how volatile compounds  
53  
54 339 are released during carbonization (Yu et al., 2019). The pyrolysis temperature also significantly

55  
56  
57 340 affects the functional chemistry of biochars, as plant-based feedstocks (e.g., crop residues)  
58  
59 341 undergo dehydration and depolymerization of lignin and cellulose in high temperatures.

60  
61  
62  
63  
64  
65

342 On the other hand, poultry manure and sewage sludge biochar do not contain lignocellulosic  
1  
2 343 compounds and do not suffer depolymerization (Ahmad et al., 2014). Pyrolysis residence time  
3  
4  
5 344 also increases SSA (Yu et al., 2019) and the ratio of O/C and H/C (Khalid et al., 2020). For  
6  
7 345 maize straw biochar, Zhang et al. (2011) indicate that increasing the pyrolysis temperature (100  
8  
9 346 to 600 °C) improves the C content (47.46 to 84.29 %), reducing the H and O contents (6.23 to  
10  
11 347 2.60 %, 45.95 to 11.95 %, respectively). Thereby, O/C and H/C ratios in biochars decrease with  
12  
13 348 high pyrolysis temperature, indicating the dehydration and deoxygenation of feedstock (Ahmad  
14  
15 349 et al., 2014). Aromatic structures are usually on lower H/C ratios, indicating biochar stability  
16  
17 350 and resistance to degradation (Varjani et al., 2019). It is worth noting that the functional group  
18  
19 351 abundance on the biochar surface reduces as the pyrolysis temperature increases (Khalid et al.,  
20  
21 352 2020).

22  
23  
24  
25  
26  
27 353 In general, the variation in the pyrolysis temperature also causes different influences in the  
28  
29 354 sorption efficiency. Liu et al. (2018) exemplified that biochar produced at high temperatures  
30  
31 355 has more active sites (i.e., greater SSA), although reducing functional groups can decrease the  
32  
33 356 pesticide sorption capacity. Thus, it is necessary to have balanced and holistic knowledge so  
34  
35 357 that biochar with the desirable characteristics is produced and applied to the soil. Additionally,  
36  
37 358 Biswas et al. (2017) compared pyrolysis behaviors (from 300 to 450 °C) of corn cob, rice husks,  
38  
39 359 rice straw, and wheat straw biochars. The optimal temperature associated with the maximum  
40  
41 360 bio-oil yield for corn cob and rice husks is 450 °C, while 400 °C is better for rice and wheat  
42  
43 361 straws. Among these biochars, rice husk presented the highest organic carbon conversion (56.62  
44  
45 362 %).

46  
47  
48  
49  
50  
51  
52 363 Crop residues are rich in minerals and, consequently, biochars have high mineral ash content  
53  
54 364 and high pH (Khalid et al., 2020). Despite biochar from higher pyrolysis temperature having  
55  
56 365 higher pH values, the  $pH_{PZC}$  and solution pH also influences sorption capacity, mainly if the  
57  
58 366 pesticide is ionizable (Khalid et al., 2020; Liu et al., 2018). Biochar has a variable charge, and  
59  
60  
61  
62  
63  
64  
65

367 high pH values increase the OH dissociation from the functional groups (e.g., carboxyl and  
368 hydroxyl) in particles' surface (Ahmad et al., 2014; Yu et al., 2019). Thus, the net negative  
369 charge increases, and there is an electrostatic attraction between negative charges and cationic  
370 species (Yaashikaa et al., 2019). Herath et al. (2016) reported the optimum pH at 4.0 for  
371 maximum glyphosate sorption (82 %) in steam-activated biochar produced from rice husk,  
372 while higher solution pH decreased the sorption capacity to 56 % (pH 10). Considering the  
373  $pH_{PZC}$  (at pH 4.0), the glyphosate dissociates, and the biochar surface charges positively (pH <  
374  $pH_{PZC} = 6.65$ ). Therefore, in acidic conditions, a strong electrostatic interaction is favored  
375 between the anionic glyphosate species (negatively charged) and the surface of the positively  
376 charged activated biochar.

377 It is essential to verify the biochar application rate and frequency to achieve ideal soil conditions  
378 for pesticide remediation (Liu et al., 2018). Different dosages can increase or decrease the pH  
379 and modify the soil electrical conductivity (EC) in varied proportions (Palansooriya et al.,  
380 2019). The efficiency of pesticide sorption in biochar-amended soils also involves competition,  
381 as the soil is a complex system with different organic and inorganic contaminants coexisting.  
382 The biochar interaction with inorganic and organic particles in the soil and other pollutants  
383 might be competitive and lead to pores blockage, decreasing the sorption capacity of targeted  
384 pesticides (Liu et al., 2018).


385 Biochar aging is a crucial factor that interferes with biochar properties and soil interactions.  
386 After application to the soil, synergistic changes of physical, chemical, and biological aging  
387 occur, and, especially in highly aromatic biochars, microbial metabolism is favored. As a result,  
388 the outer portions of the biochar particles are more altered and protect their interior from the  
389 action of the microorganisms. As biochar is a carbon-rich organic material and has relative  
390 stability and resistance to degradation, the labile carbon fraction is first degraded; hence, aged  
391 biochars become less susceptible to degradation than fresh biochars (Lehmann and Joseph,

392 2015). Ren et al. (2018) observed that wheat straw biochar's surface area and chemical  
1  
2  
3 393 composition changed with time after addition to soil. Consequently, aged biochars may have  
4  
5 394 different sorption performance than fresh biochars. Gámiz et al. (2019) indicated improved  
6  
7 395 sorption of aged biochar for removing herbicides (e.g., imazamox, picloram, and  
8  
9 396 terbuthylazine) than fresh biochar. However, Liu et al. (2018) showed that biochar aging  
10  
11 397 generally reduces its sorption potential. Therefore, the effects of aging on the fate of pesticides  
12  
13 398 in biochar-amended soils still need further research.  
14  
15  
16

17  
18 399 Biochar modified for particular soils' needs might enhance the pesticide's sorption (Kazemi  
19  
20 400 Shariat Panahi et al., 2020; Yu et al., 2019). This process occurs due to the increasing porosity,  
21  
22 401 micropore volume, SSA, and functional groups, eliminating potentially harmful elements (Liu  
23  
24 402 et al., 2018; Wei et al., 2018). Biochar modification methods include chemical reagents, iron  
25  
26 403 ions, metal oxide, and functional nanoparticles (Wei et al., 2018; Yaashikaa et al., 2019). Each  
27  
28 404 engineered biochar can be prepared by pretreating feedstock before pyrolysis or post-treating  
29  
30 405 onto untreated biochar (Liu et al., 2018). The chemical modification involves adding bases and  
31  
32 406 acids to the biochar, and generally, acid activation changes surface properties (e.g., functional  
33  
34 407 groups, charges, porosity, and SSA) (Wei et al., 2018). For example, feedstocks treated with  
35  
36 408 nitric acid can incorporate carboxylic groups (Yaashikaa et al., 2019). The primary activation  
37  
38 409 (usually using NaOH or KOH) induces the basicity and increases the O content on the surface  
39  
40 410 (Wei et al., 2018). Biochar modification also includes the physical activation using steam  
41  
42 411 (Yaashikaa et al., 2019) and the impregnation of functional nanomaterials (e.g., chitosan,  
43  
44 412 carbon nanotubes, ZnS nanocrystals, and graphene oxide). This process improves the SSA,  
45  
46 413 porosity, and thermal stability, creating a newly engineered compound used as an alternative  
47  
48 414 and innovative sorbent (Kazemi Shariat Panahi et al., 2020; Wei et al., 2018). Moreover,  
49  
50 415 modifying biochar for soil amendment must consider the costs of alteration methods and  
51  
52 416 potential risks of environmental pollution by the added elements (Liu et al., 2018).  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

417 In synthesis, several critical factors determine amending agricultural soils for pesticide sorption  
 418 (Figure 2). First, crop management should address biochar production residues, which depend  
 419 on the feedstock type and pyrolysis conditions (e.g., temperature, heating rate, and time). The  
 420 biochar characteristics related to pesticide sorption involve ash content, pH and  $pH_{PZC}$ , CEC,  
 421 functional groups, elemental ratios, porosity, and SSA. In agricultural soils, biochar  
 422 amendments contribute to pesticide sorption, leaching control, protecting the pesticide from  
 423 degradation, and avoiding microbial uptake. Moreover, the potential effects of the aging process  
 424 are not yet fully documented.

425 Figure 2 – Key aspects of biochar application for pesticide sorption on agricultural soils.



## 427 5.2. Biochar from maize crop residues

428 Twelve studies from the final selection evaluated 24 different maize residues biochars for  
 429 pesticide sorption (Table 1). Among these biochars, 15 used straw, 7 used stalk, and 2 used

1 430 corn cob as feedstock; 1 biochar was produced with pre-and post-treatment, 1 biochar with  
2 431 only pre-treatment, 1 biochar with post-treatment, and the others were not mentioned. All the  
3  
4 432 studies were conducted on a laboratory scale using spiked soils, and most of them involved  
5  
6  
7 433 agricultural soils. Also, pyrolysis temperatures ranged from 200 to 850 °C. Several studies  
8  
9 434 stated that biochar-amended soils were more effective in the pesticide sorption compared to the  
10  
11 435 biochar-free soil for atrazine (Clay et al., 2016; Ouyang et al., 2016a, 2016b; Qin et al., 2019;  
12  
13 436 Tao et al., 2020), 2,4-D (Clay et al., 2016), carbaryl (Ren et al., 2016) and thiacloprid (P. Zhang  
14  
15 437 et al., 2018). For instance, Clay et al. (2016) reported an increase from 4.5 to 6-fold on the  
16  
17 438 sorption efficiency for atrazine after soil amendment with 10 % of maize stalk biochar.  
18  
19  
20  
21

22  
23 439 The pesticide sorption on biochar amended soil depends on both pesticides and biochar  
24  
25 440 properties, besides soil characteristics. In our review, biochars from the same feedstock (e.g.,  
26  
27 441 maize crop residues) presented a significant variation in their properties: pH values varied from  
28  
29 442 6.8 to 10.6; the SSA was between 1.5 and 386 m<sup>2</sup> g<sup>-1</sup>, and ash content ranged from 3.01 to 32  
30  
31 443 %. Also, biochar properties are strongly influenced by the feedstock material characteristics  
32  
33 444 and pyrolysis conditions. For instance, maize straw biochar had the largest SSA (44.966 m<sup>2</sup> g<sup>-1</sup>  
34  
35 445 <sup>1</sup>), total pore volume (0.0345 cm<sup>3</sup> g<sup>-1</sup>), and ash content (9.03 %) than biochar from corn cob  
36  
37 446 (1.506 m<sup>2</sup> g<sup>-1</sup>, 0.0032 cm<sup>3</sup> g<sup>-1</sup> and 3.01 %, respectively). Consequently, this was probably one  
38  
39 447 of the factors that favored the better sorption performance of maize straw biochar (when the  
40  
41 448 initial concentration of atrazine solution was 1 mg L<sup>-1</sup>, the removal rate was 27.02 and 76.65  
42  
43 449 % for soil mixed with 5 % of corn cob and straw biochars, respectively) (Ouyang et al., 2016b).  
44  
45  
46  
47  
48  
49

50  
51 450 Several studies confirmed the temperature interference on the maize straw biochars SSA  
52  
53 451 among the pyrolysis conditions, improving from 6.71 to 265 m<sup>2</sup> g<sup>-1</sup>, as the pyrolysis  
54  
55 452 temperature increased from 350 to 700 °C. Similarly, the pH increased from 7.9 to 10.6.  
56  
57 453 Consequently, the highest carbaryl sorption affinity was evidenced for the biochar produced at  
58  
59  
60  
61  
62  
63  
64  
65

1 454 the highest temperature (700 °C) and with the highest rate of addition to the soil (5 %), with a  
2  
3 455 sorption affinity ( $K_f$ , Freundlich isotherm) of  $83.2 \text{ (mg kg}^{-1}) \text{ (mg L}^{-1})^{-n}$  (Ren et al., 2016).  
4  
5 456 Accordingly, Tao et al. (2020) evaluated maize stalk biochar, and batch sorption experiments  
6  
7 457 revealed an improvement in maximum sorption capacity from 21.96 to 35.88  $\text{mg g}^{-1}$ ,  
8  
9  
10 458 respectively, by increasing pyrolysis temperature from 250 to 850 °C. In the same way, biochar  
11  
12 459 application increased the  $K_f$  of imidacloprid, and clothianidin (Zhang et al., 2020) into maize  
13  
14 460 straw biochar-amended soil with charring temperature (300 - 700 °C).  
15  
16

17  
18 461 Besides, Zhang et al. (2020) reported that the amendment with low-pyrolyzing temperature  
19  
20 462 biochar (300 °C) benefited the imidacloprid and clothianidin biodegradation due to the supply  
21  
22 463 of organic carbon and available nitrogen for microorganisms, although inhibiting the chemical  
23  
24 464 degradation. On the other hand, the high-pyrolyzing temperature biochars (500 to 700 °C)  
25  
26 465 favored chemical degradation, inhibiting biodegradation. For thiacloprid, the observations  
27  
28 466 were similar (P. Zhang et al., 2018). Biochar particle size range influences the sorption  
29  
30 467 efficiency of pesticides, as observed by Clay et al. (2016). Atrazine and 2,4-D sorption  
31  
32 468 increased significantly with the reduction of the biochar particle size. For 2,4-D,  $K_f$  was risen  
33  
34 469 from 190 (for biochar mean size of 2 to 4 mm) to  $398 \mu\text{mol}^{1-1/n} \text{ L}^{1/n} \text{ kg}^{-1}$  (for size below 2 mm);  
35  
36 470 while for atrazine, the increase in  $K_f$  was more prominent from 209 to  $912 \mu\text{mol}^{1-1/n} \text{ L}^{1/n} \text{ kg}^{-1}$   
37  
38 471 in the same biochar size range (Clay et al., 2016).  
39  
40  
41  
42  
43  
44  
45

46 472 Some studies use biochar modification or pre- and post-treatment to develop biochars with  
47  
48 473 enhanced pesticide removal capacity from the soil. The maize straw biomass pretreatment with  
49  
50 474 ammonium dihydrogen phosphate (ADP) increased the atrazine removal from soil amended  
51  
52 475 due to the increase in biochar SSA ( $44.966$  for  $356.010 \text{ m}^2 \text{ g}^{-1}$ ) and total pore volume ( $0.0345$   
53  
54 476 for  $0.221 \text{ cm}^3 \text{ g}^{-1}$ ) (Ouyang et al., 2016b). Clay et al. (2016) evaluated the influence of post-  
55  
56 477 process handling. The first biochar sample was cooled to room temperature, while a second  
57  
58  
59  
60  
61  
62  
63  
64  
65



1 478 sample was immersed in water immediately after pyrolysis to prevent slow combustion. The  
2 479 atrazine sorption data for water-cooled biochar showed a  $K_f$  of  $151 \mu\text{mol}^{1-1/n} \text{L}^{1/n} \text{kg}^{-1}$ , a lower  
3  
4 480 value than the results for air-dried biochar (209 to  $912 \mu\text{mol}^{1-1/n} \text{L}^{1/n} \text{kg}^{-1}$ ).  
5  
6

7  
8 481 The biochar addition alters soil properties. For example, Zhang et al. (2020) observed that a  
9  
10 482 typical Chinese soil pH increased from 4.4 to 6.8 after a 2 % application of maize straw biochar  
11  
12 483 (500 °C). Also, pH, EC, available nutrients, organic carbon, hydrogen contents, and SSA of  
13  
14 484 the soil were improved by adding biochar from maize crop residues. The dissolved organic  
15  
16 485 carbon increased upon the addition of biochars pyrolyzed at 300 °C, while the biochars  
17  
18 486 produced at 700 °C decreased their concentrations (P. Zhang et al., 2018). Pesticide sorption  
19  
20 487 capacities are expected to vary on soils with different properties. Clay et al. (2016) pointed out  
21  
22 488 that a 1 % addition of maize stover biochar increased atrazine sorption affinity for the fine  
23  
24 489 loamy sand by about 45 % and for the silty clay loam by 25 %. The biochar-amended soil  
25  
26 490 constituents become heavily dependent on each other and with specific behaviors for each  
27  
28 491 environmental condition. At the initial atrazine solution of  $1 \text{ mg L}^{-1}$ , the soil amended with  
29  
30 492 corn cob- and stalk-based biochar (0.5 and 1.0 % addition rate) had better performance than  
31  
32 493 the pure biochar. On the contrary, at  $20 \text{ mg L}^{-1}$ , the pure biochar derived from maize stalk had  
33  
34 494 a better removal rate than the biochar-amended soil (Ouyang et al., 2016b).  
35  
36  
37  
38  
39  
40  
41  
42

43 495 Previous literature showed that the maize biochar application rate in the soil is fundamental for  
44  
45 496 optimizing pesticide sorption. Among the 12 included articles (Table 1), 11 studies involved  
46  
47 497 maximum doses of 5 %, and one study evaluated doses up to 10 %. Overall, it generated  
48  
49 498 significant improvements in the pesticide sorption capacity. The atrazine's removal rate in  
50  
51 499 column experiments increased from 38.7 to 58.3 % by increasing the application rate of the  
52  
53 500 maize straw-derived biochar from 0.5 to 1.0 % (Ouyang et al., 2016a). However, some studies  
54  
55 501 indicated that high application rates could result in negative impacts. Clay et al. (2016) reported  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 502 that 1 % addition of maize stalk biochar to two different soil types (silty clay loam and fine  
2 503 loamy sand) did not affect the pH or EC. Adding 10 % of biochar has already increased the  
3  
4 504 EC, especially in the sandy soil (less buffering capacity). However, large and frequent  
5  
6  
7 505 applications must be associated with monitoring EC to avoid saline conditions and,  
8  
9 506 consequently, losses in agricultural productivity.

11  
12  
13 507 Graber et al. (2011) investigated the interaction of 1,3-D in a solid-liquid-gaseous system of a  
14  
15 508 biochar-amended soil. The addition of 1 % of maize straw biochar (500 °C) promoted strong  
16  
17  
18 509 1,3-D sorption with solid particles, causing substantial reductions in the 1,3-D concentrations  
19  
20 510 in the gaseous phase. Still, there were also reductions in the aqueous phase, precisely the active  
21  
22 511 portion for nematode control. Thus, there was adequate control of the nematodes in the studied  
23  
24 512 system. However, the authors emphasized that biochars with greater sorption capacities (e.g.,  
25  
26 513 those with higher pyrolysis temperatures or large doses) should be used cautiously. Otherwise,  
27  
28  
29 514 depending on sorption conditions, it can negatively impact pest control and agronomic systems.

30  
31  
32  
33 515 Overall, the pesticide sorption onto the soils amended with maize residue biochars was  
34  
35 516 accurately described by the Freundlich model, such as for atrazine (Clay et al., 2016; Ouyang  
36  
37 517 et al., 2016b), 2,4-D (Clay et al., 2016), flubendiamide (Das and Mukherjee, 2020), 1,3-D  
38  
39 518 (Graber et al., 2011), carbaryl (Ren et al., 2016), thiacloprid (P. Zhang et al., 2018),  
40  
41 519 imidacloprid, and clothianidin (Zhang et al., 2020). For instance, the isothermal sorption fitted  
42  
43 520 well with the Freundlich model ( $R^2 > 0.96$ ) on soil amended with different biochar additions  
44  
45  
46 521 from corn cob and stalk (Ouyang et al., 2016b). Ren et al. (2016) also reported experimental  
47  
48 522 data adjustments ( $R^2 > 0.98$ ) of carbaryl sorption to maize straw biochar-amended soil. It  
49  
50  
51 523 suggests that the pesticide sorption occurred predominantly on the heterogeneous surfaces of  
52  
53  
54 524 the particles in these studies. On the other hand, the Langmuir model better described the  
55  
56  
57 525 topramezone's sorption process, which means the monolayer sorption played a dominating role

1 526 (Uwamungu et al., 2019). Tao et al. (2020) reported that the Freundlich model is better suited  
2 527 to fit the atrazine sorption onto biochar produced from maize stalk at 250 °C, while the  
3  
4 528 Langmuir model matched well with the biochars made at 550 and 850 °C. The kinetic  
5  
6  
7 529 experiments results showed that pesticide sorption was faster in the first phase by the larger  
8  
9  
10 530 number of active sorption sites. Thus, the sorption equilibrium was reached after 8 to 12 h,  
11  
12 531 when 0.5 g of soil and 0.02 g of biochar were added with topramezone (Uwamungu et al.,  
13  
14 532 2019), and 10 h for atrazine mixed with 15 mg of biochar (Tao et al., 2020).

16  
17  
18 533 In addition to increasing sorption efficiency, Das and Mukherjee (2020) pointed out that the  
19  
20 534 biochar addition in soil reduced the desorption process. A considerable portion of the  
21  
22 535 flubendiamide sorbed in the soil matrix remained retained during the desorption test. The  
23  
24  
25 536 biochar addition to the soil reduces diffuse pollution from leaching, as simulated by Ouyang et  
26  
27  
28 537 al. (2016a) in rainfall-leaching experiments with different soil column configurations. The  
29  
30 538 lowest leaching atrazine contents appeared in the smaller rainfall intensities and in the columns  
31  
32 539 where biochar was applied to the soil. The soil with a biochar and gravel combination provided  
33  
34  
35 540 the highest efficiency of atrazine control (on average 87.85 %) when the biochar addition rate  
36  
37 541 was 3 %.

38  
39  
40  
41 542 Previous studies have highlighted that biochar application in the soil could affect pesticide  
42  
43 543 biodegradation. For example, the biochar amendment in the soil at 5 % decreased the half-life  
44  
45 544 values of flubendiamide from 165.3 and 178.6 to 103.5 and 117.4 days, respectively (Das and  
46  
47  
48 545 Mukherjee, 2020). For carbaryl, the decrease with 5 % biochar addition was 34.6 to 27.7 (350  
49  
50 546 °C) and 33 days (700 °C) (Ren et al., 2016). For imidacloprid, the decrease with 2 % biochar  
51  
52 547 addition was 86.7 to 73.0 (300 °C), 77.6 (500 °C), and 80.6 days (700 °C) (Zhang et al., 2020).  
53  
54  
55 548 Conversely, for clothianidin, these authors reported an increase in half-life with 2 % addition  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 549 from 52.4 to 93.0 (300 °C), 72.0 (500 °C), and 62.8 days (700 °C). Thus, the degradation rate  
2 550 increased with biochar addition on soil, occurring via biotic and abiotic processes.  
3  
4

5 551 Most researchers investigate pesticide sorption individually, despite the complexity of actual  
6  
7 552 contamination and the need for combinations of treatments. Tao et al. (2020) evaluated the  
8  
9 553 combined application between biochar and co-cultured functional microorganisms. The results  
10  
11 554 showed a 49 % higher potential to degrade 100 mg L<sup>-1</sup> atrazine within 24 h than pure degrading  
12  
13 555 bacteria DNS32 and increase the soil's total and available phosphorus content. Qin et al. (2019)  
14  
15 556 assessed the maize straw biochar and sepiolite to remediate the combined pesticide and metal  
16  
17 557 pollution. While biochar targeted atrazine, cadmium was mostly immobilized by sepiolite;  
18  
19 558 thus, the authors concluded that combined application was efficient for immobilizing both  
20  
21 559 contaminants.  
22  
23  
24  
25  
26

27 560

### 31 561 *5.3. Biochar from rice crop residues*

32  
33  
34 562 Table 1 shows 15 articles that assessed 25 biochars produced from rice crop residues, most  
35  
36 563 from rice straw (n = 13). Rice husk and rice hull were also used. Pyrolysis temperature varied  
37  
38 564 from 300 to 700 °C, and just 3 of the 25 biochars passed for some treatment after pyrolysis. The  
39  
40 565 most studied pesticides were imazapic (n = 4), imazapyr (n = 3) e fomesafen (n = 3). Although  
41  
42 566 all biochars were made from rice crop residues, they have a variety of characteristics. For  
43  
44 567 instance, pH varied in the range of 6.0 to 11.1, and SSA went from 1.99 to 202.11 m<sup>2</sup> g<sup>-1</sup>. Ash  
45  
46 568 content was between 9 and 50.3 %. All these variations are related to the fact that, besides  
47  
48 569 feedstock, pyrolysis conditions are also determinants for biochar properties.  
49  
50  
51  
52  
53

54  
55 570 In general, for rice-derived biochar, higher pyrolysis temperatures resulted in materials with  
56  
57 571 higher pH, ash content, SSA, and porosity, but with lower CEC (Lü et al., 2012; Manna and  
58  
59 572 Singh, 2015; Ren et al., 2016; Yavari et al., 2016; Zhao et al., 2019). Apart from temperature,  
60  
61  
62  
63  
64  
65

573 other pyrolysis conditions influence biochar characteristics, and consequently, its sorption  
1  
2 574 capacity. Yavari et al. (2017) studied the influence of temperature (from 300 to 700 °C), heating  
3  
4 575 rate (3 to 10 °C min<sup>-1</sup>), and residence time (1 to 3 h) in rice husk biochar characteristics. The  
5  
6  
7 576 maximum organic carbon content was found in the biochar with the highest pyrolysis  
8  
9  
10 577 temperature and the slowest heating rate. CEC was negatively correlated with temperature and  
11  
12 578 heating rate, while the SSA was positively correlated with temperature and residence time.

13  
14  
15 579 Overall, rice residues-derived biochar application in the soil can improve pesticide sorption  
16  
17  
18 580 capacity (Aldana et al., 2020; Khorram et al., 2015; Lü et al., 2012; Manna and Singh, 2015;  
19  
20  
21 581 Ren et al., 2016; Wu et al., 2019). This improvement happens even with small biochar doses,  
22  
23 582 as the maximum amount used was 5 % (w/w) (Table 1). Khorram et al. (2015) investigated the  
24  
25 583 sorption of fomesafen in soils amended with 0.5, 1, and 2 % rice hull biochar produced at 600  
26  
27  
28 584 °C. Biochar-amended soils enhanced fomesafen sorption, as  $K_f$  increased from 0.69 to 14.44  
29  
30 585  $\text{mg}^{1-1/nf} \text{L}^{1/nf} \text{kg}^{-1}$  in the soil with 2 % biochar. Similarly, for carbaryl sorption, Ren et al. (2016)  
31  
32  
33 586 showed an increase of  $K_f$  from 15.5 to 125.9  $\text{mg kg}^{-1} (\text{mg L}^{-1})^{-n}$  after applying rice biochar  
34  
35 587 (700 °C) at 5 %. In this case, the authors compared biochars produced at two different  
36  
37  
38 588 temperatures and showed that biochars made at higher temperatures had the most satisfactory  
39  
40 589 results. Several studies have also examined the biochar effects in the leaching of pesticides,  
41  
42  
43 590 and it has been demonstrated that biochar helps retain the substances at the top layer. For  
44  
45 591 example, Khorram et al. (2018) assessed the leaching of fomesafen in rice biochar-amended  
46  
47  
48 592 soil in columns, and the retention increased from 13.45 to 76 % with 2 % (w/w) biochar-  
49  
50 593 amended soil; the pesticide was retained in the top 5 cm of the columns.

51  
52  
53 594 Manna and Singh (2015) also reported the temperature effects for pyrazosulfuron-ethyl, as  
54  
55  
56 595 biochars from higher temperatures achieved higher specific surface and porosity. In this  
57  
58 596 research, the control soil (sandy loam) had 5.3 to 8.6 % of pesticide sorption, whereas biochar  
59  
60  
61  
62  
63  
64  
65

1 597 produced at 400 °C had 7.5 to 50.4 %, and 600 °C biochar sorption was 55.9 to 91.8 %.  
2  
3 598 However, Yavari et al. (2017b) reported different effects for imazapic and imazapyr sorption  
4  
5 599 by soils with 1 % of rice husk biochar at different pyrolysis conditions. In this case, sorption  
6  
7 600 capacities for both herbicides were higher for the biochar produced at the lowest temperature  
8  
9 601 (300 °C) and a slow heating rate (3 °C min<sup>-1</sup>).  
10

11  
12  
13 602 Few studies addressed biochars after treatments. Yavari et al. (2020) produced a rice husk  
14  
15 603 biochar (pyrolysis at 300 °C, 3 h) and modified this material by mixing it with a chitosan  
16  
17 604 solution diluted in acid acetic for 30 min, and by dripping NaOH solution into it. After 12 h,  
18  
19 605 deionized water washed the modified biochar. The comparative analysis of biochar sorption  
20  
21 606 performance before and after treatment is indicative of the changes resulting from the action of  
22  
23 607 chitosan. The authors reported that chitosan-modified biochars had remarkably higher moisture  
24  
25 608 content, pH, and CEC than non-modified biochars. Chitosan-modified biochar also showed a  
26  
27 609 higher sorption capacity of imazapic and imazapyr. The value of  $K_f$  for imazapic varied from  
28  
29 610 1.744 in the pure soil to 3.090 in the biochar amended soil with 1 % biochar and to 4.391 for  
30  
31 611 chitosan-modified biochar amended soil. However, it would also be relevant to test only  
32  
33 612 chitosan to validate its effects in the modified biochar.  
34  
35  
36  
37  
38  
39  
40

41 613 Finally, most researchers also investigated the pesticides' sorption individually. A recent study  
42  
43 614 by Aldana et al. (2020) examined the sorption of atrazine, diuron, and OTC on mixtures of  
44  
45 615 tropical soils and rice biochar at 1, 2.5, and 5%. Higher doses increased the sorption of all  
46  
47 616 compounds, and atrazine was the only substance detected in the equilibrium solution, even in  
48  
49 617 the mixture with 5 % biochar. According to these authors, diuron was sorbed first and competed  
50  
51 618 with atrazine for the sorption sites. Besides, soil dissolved organic matter might cover sorption  
52  
53 619 sites from biochar. Despite sorption, the degradation of the pesticides by biochar amended-  
54  
55 620 soils has also been addressed, and it depends on the pesticide's type. Manna and Singh (2019)  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

621 showed that applying 400 and 600 °C rice straw biochar in sandy loam soil at a biochar dose  
1  
2  
3 622 of 0.02 and 0.05 g kg<sup>-1</sup> increased pyrazosulfuron-ethyl persistence under flooded and non-  
4  
5 623 flooded conditions. An increasing amount of biochar decreased pyrazosulfuron-ethyl  
6  
7 624 degradation. Conversely, for carbaryl, Ren et al. (2016) reported that biochars as soil  
8  
9  
10 625 amendments promoted chemical degradation and affected the pesticide's microbial  
11  
12 626 degradation.

13  
14  
15  
16 627

#### 19 628 *5.4. Biochar from wheat crop residues*

22 629 Table 1 shows 13 different biochars produced from wheat crop residues. Most of them are  
23  
24  
25 630 derived from wheat straw, and only one from wheat midds. The pyrolysis temperature in these  
26  
27 631 studies ranged from 250 to 600 °C. Among the materials (Table 1), only one was submitted to  
28  
29 632 treatment before pyrolysis (Tatarková et al., 2013). The straw was carbonized in a sand bath at  
30  
31 633 200 °C for 24 h, and the residue was pyrolyzed at 300 °C. These papers studied a variety of 7  
32  
33  
34 634 pesticides, and the most studied were pyrazosulfuron-ethyl and HCB.

37  
38 635 As the previous feedstocks, various characteristics of wheat residue biochars changed in  
39  
40 636 different pyrolysis conditions. For instance, the pH varied between 5.4 and 10.6, CEC was in  
41  
42 637 the range of 22 and 68.7 cmol<sub>c</sub> kg<sup>-1</sup>, and SSA ranged from 2.7 to 62.6 m<sup>2</sup> g<sup>-1</sup>. In general, the  
43  
44 638 studies showed that increasing pyrolysis temperatures increases pH, SSA, and porosity (Cheng  
45  
46 639 et al., 2017; Manna and Singh, 2015). However, regarding the effect of pyrolysis temperature  
47  
48 640 on CEC, there were contrasting results. For example, Cheng et al. (2017) studied wheat straw  
49  
50 641 biochar produced at four peak temperatures (250, 350, 450, and 550 °C) and found that CEC  
51  
52 642 decreased (from 68.7 to 22.0 cmol<sub>c</sub> kg<sup>-1</sup>) with higher temperatures. Conversely, Manna and  
53  
54  
55 643 Singh (2015) showed that increasing pyrolysis temperature for wheat straw biochar did not  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 644 significantly affect CEC (62.0 and 62.6  $\text{cmol}_c \text{ kg}^{-1}$  for the 400 and 600 °C biochars,  
2 645 respectively).

3  
4  
5 646 Studies regarding wheat residue biochar provide evidence that the application of this material  
6  
7  
8 647 in soils can improve the sorption of pesticides (Cheng et al., 2017; Humera Aziz et al., 2018;  
9  
10 648 Manna and Singh, 2015; Tatarková et al., 2013). Tatarková et al. (2013) used a wheat straw  
11  
12 649 biochar produced at 300 °C (pH of 10.6 and SSA of was 4.59  $\text{m}^2 \text{ g}^{-1}$ ). They demonstrated that  
13  
14  
15 650 1 % biochar amended soil had 2.53 times higher MCPA sorption capacity than the non-amended  
16  
17  
18 651 soil. Furthermore, the quantity of MCPA leached in soil columns alone was 56 %, while 35 %  
19  
20 652 leached in columns with 1 % wheat straw biochar. Humera Aziz et al. (2018) demonstrated that  
21  
22 653 the application of wheat straw biochar prepared at 500°C (CEC of 85  $\text{cmol}_c \text{ kg}^{-1}$ , total organic  
23  
24 654 carbon of 43.80 % and SSA of 4.83  $\text{m}^2 \text{ g}^{-1}$ ) at 0.25 and 0.5 % in sandy clay loam soil (pH of  
25  
26 655 7.44, CEC of 5.2  $\text{cmol}_c \text{ kg}^{-1}$  and total organic carbon of 0.87 %) from Pakistan increased  
27  
28  
29 656 chlorpyrifos sorption. The data was adjusted by Freundlich isotherm, and  $K_f$  rose from 4.34 to  
30  
31  
32 657 218.83  $\text{L kg}^{-1}$  after the mixture with 0.5 % biochar. Manna and Singh (2015) reported a similar  
33  
34 658 effect for pyrazosulfuron-ethyl sorption by adding 0.1, 0.2, and 0.5 % wheat straw biochar in  
35  
36  
37 659 sandy loam soil from India. Increasing the pyrolysis temperature from 400 to 600 °C did not  
38  
39  
40 660 significantly increase the organic carbon and CEC, but it doubled the SSA (10.15 and 20.38  $\text{m}^2$   
41  
42 661  $\text{g}^{-1}$ , respectively) and improved the sorption capacity of the biochar. Manna and Singh (2019)  
43  
44 662 also demonstrated that amendment with these same biochars influences pesticide leaching  
45  
46  
47 663 because of its strong sorption. They reported that the application of wheat straw biochar made  
48  
49  
50 664 at 400 and 600 °C at 5 % in the soil led to a decrease in the pyrazosulfuron-ethyl loss by 9 e 39  
51  
52 665 %, respectively.

53  
54  
55 666 In general, the Freundlich isotherm adjusted pesticide sorption data (Humera Aziz et al., 2018;  
56  
57 667 Manna and Singh, 2015; Tatarková et al., 2013). Manna and Singh (2015) discussed the effect  
58  
59  
60 668 of biochar on the mean value (standard deviations were not provided) of the Freundlich  
61  
62  
63  
64  
65



669 exponent  $1/n$ , as nonlinearity in pyrazosulfuron-ethyl sorption increased with higher biochar  
670 doses. The  $1/n$  values were higher than 1 in the pure soil, suggesting nonlinear sorption  
671 isotherms (S-type). On the other hand,  $1/n$  becomes less than 1 in soils amended with 0.2 and  
672 0.5 % biochar, reflecting an L-type adsorption isotherm. It indicates that the adsorbate  
673 molecules did not suffer intense competition from the water molecule at low concentrations,  
674 but sorption sites become limited as the concentration increases. Unlike these results, Tatarková  
675 et al. (2013) showed that the Freundlich exponent values for MCPA sorption were close to 1  
676 for non-amended soil, biochar-amended soil, and biochar, indicating constant MCPA  
677 partitioning between sorbent and solution for all the materials within the concentration range  
678 (0.5 to 22 mg L<sup>-1</sup>). The use of classifications with numerical ranges should always be done with  
679 caution. Statistical analysis and presentation of standard deviations of results, especially close  
680 to class boundaries, may be of interest to reduce uncertainties and increase the reliability of data  
681 interpretations.

682 Although several studies involve biochar effects on pesticide sorption, questions have been  
683 raised about their impact on other processes, including degradation, volatilization, and plant  
684 uptake (Cheng et al., 2017; Manna and Singh, 2019; Wang et al., 2016). Recent evidence  
685 suggested that the adding wheat straw biochar improves the pesticide persistence in soil due to  
686 the strong affinity between pesticide and biochar particles. Some studies sustain this hypothesis,  
687 such as Cheng et al. (2017) for simazine and Manna and Singh (2019) for pyrazosulfuron-ethyl.  
688 Moreover, in the study of Tatarková et al. (2013), the half-life of MCPA increased from 5.2  
689 days for the non-amended soil to 21.5 days for the amended soil. On the other hand, the  
690 dissipation would increase in biochar amended soils because of the microbial activity  
691 stimulation by biochar nutrients. Song et al. (2012) carried out incubation tests with HCB and  
692 reported that in the first 2 weeks, faster HCB dissipation occurred in the 0.1 and 0.5 % biochar-

693 amended treatments compared to control. Nevertheless, a significant decrease in HCB  
694 dissipation was noted in the biochar-amended soils after 4 weeks.

695

## 696 **6. Key aspects, gaps, and uncertainties**

### 697 *6.1. Pesticide bioavailability and ecotoxicity*

698 Regarding the pesticide uptake by plants, Khorram et al. (2017) showed that rice biochar  
699 amendment could effectively suppress the undesirable inhibitory effect of fomesafen on corn  
700 growth. Song et al. (2012) tested the bioaccumulation of HCB in earthworms; their primary  
701 concern was that strong sorption of contaminants would result in accumulation by the  
702 mesofauna of ingestion of biochar particles. However, concentrations of HCB in earthworms  
703 decreased with increasing biochar application rate. After that, in another study, Song et al.  
704 (2016) investigated a biochar-plant tandem remediation scheme for HCB, adding 1 and 2 % of  
705 wheat biochar in soil and using ryegrass as the growing plant. Biochar amendment immobilized  
706 HCB in soils, and the uptake of HCB by ryegrass root was reduced by 93.05 % by 1 % biochar  
707 amendment compared to that of unamended soil. Unfortunately, no selected study presented  
708 ecotoxicological assessments of pesticide biochar sorption, despite two studies identified in the  
709 previous steps. For instance, Bielská et al. (2018) indicated reducing bioavailability and  
710 bioaccessibility of hydrophobic organic compounds by increasing 5 % the dose of biochar from  
711 mixed wood shavings and rice husks. However, this study identified biochar-induced toxicity  
712 by the highest amount (10 %) to the springtail *Folsomia candida*, which could be associated  
713 with pH-related effects or food sorption.

714 Although most studies focused on the remediation of isolated pesticides, these contaminants  
715 may have increased ecological risks in combination (Belden and Brain, 2018). Additionally,

1 716 some pesticide transformation products can be more toxic than the parental compound (Sinclair  
2 717 and Boxall, 2003), a challenge for pesticide management and environmental remediation  
3  
4 718 (Levine and Borgert, 2018). Nonetheless, in a co-contamination, hydrophobic organic  
5  
6  
7 719 compounds (including pyrene and DDE) in the mixture did not show competitive sorption to  
8  
9  
10 720 biochar (Bielská et al., 2018), which is an advantage considering multiple soil contaminants.

11  
12  
13 721 Biochar selection should allow pollution control while maintaining pesticide efficiency. For  
14  
15 722 instance, Lü et al. (2012) incorporated 2,4-D and acetochlor into the studied rice straw biochar  
16  
17  
18 723 (i.e., pyrolyzed at 350 °C, with 31 % ash, 20.6 m<sup>2</sup> g<sup>-1</sup> of SSA, and H/C ratio of 0.342) as a  
19  
20 724 sorptive carrier for retaining these herbicides. Doses of 0.5 and 1 % of biochar in soil decreased  
21  
22  
23 725 the acetochlor release rate, indicating the capacity to regulate herbicides' behavior. Besides,  
24  
25 726 Graber et al. (2011) suggested that higher SSA wood-derived biochar could negatively impact  
26  
27  
28 727 herbicide efficacy in terms of phytoavailability. Thus, depending on the biochar characteristics  
29  
30 728 and strength of sorption capacity, pesticide concentrations in the aqueous solution decrease and  
31  
32  
33 729 are not enough to promote pest control. This should be considered when planning the  
34  
35 730 application.

36  
37  
38 731

## 39 40 41 732 *6.2. Microbiological processes associated with biochar sorption*

42  
43  
44  
45 733 Most of the retrieved studies did not discuss microbial activity and its interactions with  
46  
47 734 pesticides and biochar. On the contrary, some studies sterilized soils before biochar application  
48  
49  
50 735 to avoid microbial interferences. Still, pesticides can affect the soil microbiota and inhibit  
51  
52 736 biodegradation. For example, Zhang et al. (2018) indicated that thiacloprid shifted the soil  
53  
54  
55 737 microbiota, and the biodegradation depended on soil pH, dissolved organic matter, and nutrients  
56  
57 738 (e.g., N and P). Nonetheless, biochar addition generally enhanced microbial activity and  
58  
59 739 multiplication, leading to more significant pesticide degradation, as shown by Das and  
60  
61  
62  
63  
64  
65

1 740 Mukherjee (2020). These authors considered that maize stalk biochar was a readily available  
2 741 source of energy for the microbiota.  
3

4  
5 742 With a combined application approach of biochar in soil together with a degrading bacteria  
6  
7 743 (i.e., *Acinetobacter lwoffii* DNS32), Tao et al. (2020) showed a 49 % greater capacity to degrade  
8  
9 744 100 mg L<sup>-1</sup> of atrazine in 1 day compared to experiments using only adding the bacteria to the  
10  
11 745 soil. In this scenario, microbial degradation could also increase pesticide dissipation, as  
12  
13 746 observed by Wu et al. (2019) for oxyfluorfen and maize straw and rice hull biochars.  
14  
15 747 Accordingly, Song et al. (2016) showed that wheat straw biochar stimulated microbial activity  
16  
17 748 for dissipating hexachlorobenzene, mainly when associated with plant roots. Wu et al. (2019)  
18  
19 749 also found that a rice hull biochar pyrolyzed at 500 °C (with 50.34 % ash, SSA of 95.67 m<sup>2</sup> g<sup>-1</sup>  
20  
21 750 <sup>1</sup>, and pH of 6.96) could improve oxyfluorfen degradation. The sorbent provided a suitable  
22  
23 751 environment for microorganisms and stimulated soil microbial activity. Wang et al. (2016)  
24  
25 752 showed that wheat midds biochar (i.e., pyrolyzed at 550 °C, with pH of 8.86, SSA of 24.73 m<sup>2</sup>  
26  
27 753 g<sup>-1</sup>, and pH of 8.82) alters 1,3-D degradation rates primarily through abiotic processes. They  
28  
29 754 indicated a 130 % increase in the half-life for pesticide dissipation in soil with 1 % biochar,  
30  
31 755 while 60 % higher in sterilized soils than control.  
32  
33  
34  
35  
36  
37  
38  
39

40 756 Furthermore, the biochar effect in the biodegradation of pesticides may vary with pyrolysis  
41  
42 757 temperatures. For example, Ren et al. (2016) studied carbaryl degradation in soils amended  
43  
44 758 with rice and maize biochars under sterile and unsterile conditions. There was an increase in  
45  
46 759 the biotic removal rate of carbaryl for soils amended with biochars produced at low  
47  
48 760 temperatures (350 °C). However, the addition of biochar from high temperatures (700 °C)  
49  
50 761 decreased biodegradation rates. The authors assumed that this difference is because of easily  
51  
52 762 degradable compounds in the low-temperatures biochars. Also, these biochars did not affect the  
53  
54 763 soil pH, which favors microbial communities. Zhang et al. (2020) observed similar behavior as  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

764 a function of pyrolysis temperature for biodegradation of imidacloprid and clothianidin in soil  
765 amended with maize straw biochar.

766

### 767 *6.3. The effects of the aging process on biochar sorption*

768 Ren et al. (2018) identified changes in wheat straw biochar properties (1 % dose) and its atrazine  
769 sorption capacity by aging in field soil (0.5, 1, 1.3, and 2 years). They reported that the SSA of  
770 the fresh biochar amended soil increased from 10.7 to 59.1 m<sup>2</sup> g<sup>-1</sup> after 0.5 years and then  
771 decreased with aging time, reaching 8.33 m<sup>2</sup> g<sup>-1</sup> after 2 years in the soil. According to the  
772 authors, this behavior is related to the initial exposure of micropores due to the dissipation of  
773 labile fraction from the biochar surface just after the application of biochar (increasing SSA),  
774 and then the adhesion of soil organic matter and minerals to the surface of the biochar  
775 (decreasing SSA). Another study (Khorram et al., 2017) investigated the aging process on  
776 fomesafen sorption-desorption and indicated that the rice husk biochar was still effective after  
777 6 months compared to the unamended soil. However, for 0.5 % of biochar, the total leaching  
778 rate of fomesafen increased from 67.5 % in freshly amended soil to 78.1 % in the soil aged 6  
779 months. So, the aging process generally reduces the sorption capacity of rice biochars,  
780 suggesting a need for reapplication.

781 Similarly, the decrease of sorption because of the aging effect was also reported for oxyfluorfen  
782 (Wu et al., 2019). In this case, to simulate the aging process, they used an artificial method  
783 (H<sub>2</sub>O<sub>2</sub> treatment). The soil's capacity to absorb oxyfluorfen increased linearly with added  
784 biochar, but the sorption capacity of biochar-amended soil decreased with aging. According to  
785 these authors, over time, changes might occur on physical or chemical properties in biochar  
786 surface or blocked sorption sites by acid, minerals, oxides, and native pollutants in soil.  
787 However, after 6 months of aging, the sorption capacity was still more significant than

1 788 measured for the unamended soils. Thus, desorption could result from pH changes,  
2 789 reintroducing the pesticide to the soil (Liu et al., 2018), and the long-term biochar application  
3  
4 790 and aging effects still need further research.  
5  
6  
7

8 791 As feedstock, different crop residues result in biochars with various properties and,  
9  
10 792 consequently, different pesticide sorption capacities. Some researchers in our final selection  
11  
12 793 compared biochars made of varying crop residues at the same conditions. For example, Ren et  
13  
14 794 al. (2016) studied rice straw and maize straw biochars pyrolyzed at 350 °C. The authors  
15  
16 795 reported that a 5 % rice straw biochar-amended soil showed higher sorption of carbaryl than a  
17  
18 796 5 % maize straw biochar-amended soil. Some of the factors that contributed to these results  
19  
20 797 were the higher SSA, ash content, and H/C ratio of the rice straw biochar (9.01 m<sup>2</sup> g<sup>-1</sup>, 29.1 %,  
21  
22 798 and 0.725, respectively) when compared with the properties of the maize straw biochar (6.71  
23  
24 799 m<sup>2</sup> g<sup>-1</sup>, 15.4 % and 0.685 respectively). Wu et al. (2019) also reported a higher sorption capacity  
25  
26 800 for oxyfluorfen by biochar derived from rice hull at 500 °C than maize straw biochar at 300  
27  
28 801 °C. Rice biochar also seems to be more effective than wheat biochar. Manna and Singh (2015)  
29  
30 802 compared the effect of these two materials on pyrazosulfuron-ethyl sorption in sandy loam soil.  
31  
32 803 Rice-derived biochar produced at 600 °C showed higher pH, O, C, and porosity than the wheat  
33  
34 804 biochar made at the same conditions. As a result, rice biochar sorbed 55.9 to 91.8 % of the  
35  
36 805 herbicide, while wheat biochar sorbed 6.9 to 86.0 %. Then, Manna and Singh (2019) reported  
37  
38 806 that rice biochar also had a more considerable effect on herbicide degradation. Under non-  
39  
40 807 flooded conditions, the application of 0.05 g kg<sup>-1</sup> of rice biochar produced at 600 °C resulted  
41  
42 808 in a 6-fold increase in the half-life of pyrazosulfuron-ethyl.  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

53 809

#### 56 810 *6.4. Potential pesticide desorption*

57  
58  
59  
60  
61  
62  
63  
64  
65

1 811 In addition to sorption, studying pesticide desorption is also essential to understand the strength  
2 812 of the bond between pesticide and biochar components, the reversibility of the reaction, and  
3  
4 813 pesticides' availability. However, only a few papers present discussed desorption results (10  
5  
6  
7 814 out of 32), including the pesticides fomesafen, 2,4-D, pyrazosulfuron-ethyl, acetochlor,  
8  
9 815 imazapic, imazapyr, flubendiamide, HCB, and MCPA. Only one study tested desorption with  
10  
11 816 maize residues biochar (total of 12). In addition, the retrieved studies from this review lack the  
12  
13 817 ecotoxicological analysis regarding the desorption process. In general, the Freundlich equation  
14  
15 818 satisfactorily fitted the desorption isotherms. These authors compared materials using the  
16  
17 819 percentage of desorbed pesticide, the desorption coefficient ( $K_f$ ), and the hysteresis coefficient  
18  
19 820 ( $H$ ), indicating the reaction potential reversibility.  
20  
21  
22  
23  
24

25 821 Most of the retrieved papers showed a decrease in desorption due to the addition of biochar in  
26  
27 822 soils. For instance, Khorram et al. (2017) studied the sorption of fomesafen in mixtures of soil  
28  
29 823 with rice straw biochar and showed that the desorption ( $K_f$ ) varied from  $0.42 \text{ mg}^{1-1/n} \text{ L}^{1/n} \text{ kg}^{-1}$   
30  
31 824 in the unamended soil to  $21.24 \text{ mg}^{1-1/n} \text{ L}^{1/n} \text{ kg}^{-1}$  for the mixture with 2 % biochar. In addition,  
32  
33 825 the value of  $H$  also decreased (from 0.76 in the unamended soil to 0.50 with 2 % biochar),  
34  
35 826 indicating less desorption. Similar results were found by Manna e Singh (2015) for  
36  
37 827 pyrazosulfuron-ethyl, Khorram et al. (2018) for fomesafen, Yavari et al. (2016; 2019) for  
38  
39 828 imazapic e imazapyr (all of them using rice straw biochar), Das and Mukherjee (2020) for  
40  
41 829 flubendiamide (with maize stalk biochar), and Song et al. (2016) for HCB (using wheat straw  
42  
43 830 biochar). Therefore, they suggest that the bond between pesticides and biochar particles is more  
44  
45 831 robust than between pesticides and unamended soil particles.  
46  
47  
48  
49  
50  
51  
52

53 832 On the contrary, some researchers pointed out that the increase of sorption capacity from the  
54  
55 833 same dose is not always proportional to the decrease in desorption. For instance, Tatarkova et  
56  
57 834 al. (2013) concluded that the bonds between wheat straw biochar and MCPA might be weak.  
58  
59  
60  
61  
62  
63  
64  
65

835 Despite decreased desorbed MCPA (from 64.2 % for the unamended soil to 55.1 % with 1 %  
836 biochar), these differences are less significant than the sorption results. Khorram et al. (2015)  
837 presented similar results for fomesafen in rice biochar as the increase of desorption  $K_f$  with  
838 increasing biochar dose; the value of H was between 0.41 to 0.71 in the unamended soils and  
839 between 0.95 to 0.99 in soils with 2 % biochar.

840 Furthermore, there is no consensus regarding the effect of different pyrolysis temperatures on  
841 desorption for various pesticides. For example, Yavari et al. (2016) studied soil with 1 % rice  
842 biochar and found more desorption of imazapic and imazapyr for mixtures using biochar  
843 produced at higher temperatures. On the contrary, Lu et al. (2012) studied the desorption of  
844 acetochlor and 2,4-D in pure rice biochar and pointed out that there was more reversibility for  
845 biochars produced at lower temperatures.

846

## 847 **7. Biochar transport, storage, and applicability**

848 Most studies on pesticide-contaminated soils consist of lab-scale batch experiments, and there  
849 is a gap on full-scale or *in situ* technologies (Morillo and Villaverde, 2017). For instance, only  
850 two papers reported results on pilot-scale experiments and biochar field application in the final  
851 paper selection. Among several challenges related to the utilization of biochar, easing handling  
852 and storage steps are critical for actual biochar implementation in the agriculture business.  
853 Therefore, this section describes some options for appropriate: i) transport of biochar produced  
854 in pyrolysis plants to the farm; ii) storage of biochar in the farm; and iii) dispersion of biochar  
855 in the soil.

856 Significant aspects impacting the transport of biochar from pyrolysis plants to farms are the  
857 distance between these two businesses, the size of the agricultural company, and its demand for  
858 biochar. In general, farms' demand for biochar could be fulfilled by packing powders in big



1 859 bags and distributing them to regional cooperative or logistic centers via road transport.

2 860 However, it might be feasible for farms with attached pyrolysis units to implement connected

3 861 transport and biochar storage systems. This initiative decreases human intervention in the

4 862 process, hence increasing overall farms' yield.

5 863 From the reactor unit, pneumatic conveying lines could transfer biochar to storage vessels or

6 864 silos, which means that the bulk material is transported through a pipeline with an airstream.

7 865 Biochar is fluidized with low gas flow rates due to its low particle density and particle size;

8 866 hence, pneumatic transport is an appealing economic way to handle such material. Some

9 867 authors measured the basic flow energy and permeability of biochar originated from lignin

10 868 residues (Toloue Farrokh et al., 2018) and birch tree chips (Toloue Farrokh et al., 2020), and

11 869 concluded that these powders (mean diameter from 52 to 73  $\mu\text{m}$ , and bulk density from 574 to

12 870 737  $\text{kg m}^{-3}$ ) might flow easily and with lower tendency to cause blockage of pipes during

13 871 pneumatic conveying operations.

14 872 Other relevant advantages of pneumatic conveying systems are that it requires very little

15 873 maintenance, and powders can be either conveyed in dilute or dense mode to prevent the

16 874 breakage of particles; the system's design is flexible as pipelines can be adjusted to the farm

17 875 layout; it is a dust-free system since the flow of biochar occurs inside of tubes; and the solids

18 876 flow rate can be monitored from simple pressure measurements as well as controlled with non-

19 877 mechanical valves, which are inexpensive and are resistant to wear and seizure (Geldart and

20 878 Jones, 1991; Massaro Sousa et al., 2020; Smolders and Baeyens, 1995).

21 879 Although there is limited information on the actual conveying of biochar, pneumatic transport

22 880 is a consolidated research field. For example, the efficient design of transport pipelines might

23 881 be performed using flow diagrams based on dimensionless numbers (e.g., Reynolds and

24 882 Archimedes) that account for different powder and air properties and pipeline dimensions

1 883 (Kalman, 2020; Kalman and Rawat, 2020). Moreover, practical experience and research  
2 884 literature on the conveying of pulverized coal are vast, which is helpful information to handle  
3  
4 885 biochar given the similarities between these two powders in terms of bulk density and particle-  
5  
6  
7 886 size distribution. Papers reporting experimental tests in various scale units and numerical  
8  
9  
10 887 simulations with different approaches are addressed in detail elsewhere (Chinnayya et al., 2009;  
11  
12 888 Cong et al., 2011; Jin et al., 2019; Lu et al., 2011; Zhou et al., 2020, 2018).

13  
14  
15 889 Silos are appealing equipment to meet process storage demand that also provides complete  
16  
17  
18 890 isolation from the external environment, minimizing biochar loss due to humidity degradation,  
19  
20 891 pests' contamination, or dust. Jenike's method is often used to design cylindrical or wedge-  
21  
22 892 shaped silo hoppers aiming at appropriate mass- or funnel-flow discharge of bulk solids with  
23  
24  
25 893 different properties (Enstad, 1975; Massaro Sousa et al., 2021; Oginni and Fasina, 2018).  
26  
27 894 However, to the best of our knowledge, the flow properties of biochar required for the design  
28  
29  
30 895 of silos have not been fully assessed in the literature, which is of great interest for future studies.

31  
32  
33 896 Despite this, successful design and operation of silo hoppers with pulverized coals are reported  
34  
35  
36 897 for both gravity-driven and air-assisted discharge of powders (Guo et al., 2015; Lu et al., 2015,  
37  
38 898 2012). Depending on samples' particle-size distribution and moisture content, minimum hopper  
39  
40  
41 899 inclination for mass-flow discharge ranges from 11 to 22°, whereas the outlet diameter must be  
42  
43 900 higher than 0.3 m (Chen et al., 2012; Y. Liu et al., 2015). Biochar generally presents better  
44  
45 901 flowability classification than pulverized coal under consolidated conditions (Toloue Farrokh  
46  
47  
48 902 et al., 2020, 2018), thus using steeper silos with cone angles of 10° and outlet diameter higher  
49  
50  
51 903 than 0.3 m might be sufficient to handle most biochar powders. However, measuring bulk and  
52  
53 904 flow properties is highly encouraged to optimize the silo design and prevent flow issues,  
54  
55 905 significantly whether the silo wall material or the biochar particle size and moisture content  
56  
57  
58 906 deviate from the ones shown in the previous studies.

1 907 In principle, biochar can be dispersed in soil with equivalent techniques for spreading fertilizers  
2  
3 908 and pH correctors. However, these methods should be optimized considering farm  
4  
5 909 particularities, business size, terrain conditions (e.g., slopes and soil wettability), local climate,  
6  
7 910 and duration of rainy seasons, among other aspects. Dust emissions can be reduced, for  
8  
9 911 example, with wet soil applications (Li et al., 2018). Moreover, using pelletized biochar might  
10  
11 912 be an alternative to prevent excessive biochar loss by wind action or water leaching. While  
12  
13 913 some papers reported the successful biochars densification for solid fuel applications (Bazargan  
14  
15 914 et al., 2014; Hu et al., 2016; Ranzi et al., 2018), future studies should consider pellet properties  
16  
17 915 for field applicability. Some relevant factors include pellet resistance when in contact with  
18  
19 916 different environmental conditions, total contact area for pesticide sorption, and optimal pellet  
20  
21 917 size to perform an economic and homogeneous distribution of biochar in soil.  
22  
23  
24  
25  
26

27 918 In general, adding biochar to the soil can result in several benefits, including improvements in  
28  
29 919 carbon sequestration, greenhouse gas mitigation, soil fertility, plant growth, and crop yields  
30  
31 920 (Ding et al., 2016; Hussain et al., 2017), solid advantages for agricultural areas. Besides, plant  
32  
33 921 growth stimulation could improve ecological restoration and ecosystem services (Rey Benayas  
34  
35 922 et al., 2009). The productivity increases due to more significant water holding capacity and  
36  
37 923 lower infiltration, which could also be an alternative for degraded or arid soils (Diatta et al.,  
38  
39 924 2020). The feedstock for biochar production is varied, including crop residues, wood biomass,  
40  
41 925 leaves, animal manure, solid waste, sewage, and industrial sludge (Ahmad et al., 2014;  
42  
43 926 Lehmann and Joseph, 2015). The alternative for applying crop residues to biochar production  
44  
45 927 could concatenate agricultural waste management and environmental remediation techniques.  
46  
47  
48  
49 928 Finally, note that the best selection of units and methods for handling biochars is intrinsically  
50  
51 929 related to the regional technology development, farm business size, and field particularities;  
52  
53 930 hence, performing techno-economic assessment under different farm contexts is encouraged to  
54  
55 931 clarify this point.  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 932 8. Final considerations

1  
2  
3 933 The increase in crop production had led to a demand for residue management, which could be  
4  
5 934 coupled with biochar production. Biochar has a potential for pesticide sorption in agricultural  
6  
7  
8 935 soils in this scenario, and studies have increased in the last few years. Particularly maize, rice,  
9  
10 936 and wheat residues are appropriate feedstocks for biochar production and contaminated soil  
11  
12  
13 937 amendment. In this review, we highlighted some advantages of these applications, such as:

- 14  
15  
16 938 ● Biochar amendments improved soil quality, restored soil properties, and contributed to  
17  
18 939 carbon sequestration.
- 20  
21 940 ● Biochar amendments have the potential sorption of pesticides, reducing their soil  
22  
23 941 mobility and bioavailability.
- 25  
26 942 ● Moreover, studies have shown several benefits from biochar application, including the  
27  
28 943 potential to ameliorate soil quality and properties (e.g., organic matter). Thus, biochar  
29  
30 944 could promote soil restoration and ecosystem services maintenance.
- 32  
33 945 ● Agricultural cooperatives can manage biochar production and logistics, and this could  
34  
35 946 benefit mini-farms as a low-cost amendment.
- 37  
38 947 ● Multiple feedstocks for biochar production could be used in rotation systems, as residue  
39  
40 948 management in the long-term could improve carbon sequestration in agriculture.

42  
43 949 On the other hand, these applications still need further research and development, mainly due  
44  
45 950 to some limitations and gaps, including:

- 47  
48 951 ● Most research focused on batch experiments in lab-scale, highlighting a need for field  
49  
50 952 application research and study of other contaminants. Also, there is a need to study the  
51  
52 953 long-term effects of biochar application and its potential desorption.
  - 54  
55 954 ● Retrieved studies were mainly for a few pesticides (only 21 different active substances).  
56  
57 955 Therefore, biochar sorption should be evaluated for different pesticides and co-
- 58  
59  
60  
61  
62  
63  
64  
65

956 contaminations since they behave differently depending on their chemical  
1  
2 957 characteristics.

3  
4  
5 958 ● Studies should consider the entire biochar life cycle. Monitoring all steps is essential to  
6  
7 959 ensure the biochar production meets the proposed goals for its application. Also, this  
8  
9 960 approach can avoid the risks of biochar containing contaminants (e.g., PAHs).

10  
11 961 ● Future studies should evaluate the influence of biochar particle size on remediation  
12  
13 962 dynamics, especially *in situ*. Micro- and nano-particles can show more significant  
14  
15 963 horizontal movement (runoff) and vertical movement. Thus, once they have absorbed  
16  
17 964 pesticides, they can be transported to greater distances.

18  
19 965 ● There is still limited knowledge about the aging process of biochar. A more systematic  
20  
21 966 understanding of the complexity of physical and chemical changes is needed, and also  
22  
23 967 the influence of different soil types and climate changes (e.g., wet/dry and  
24  
25 968 freezing/melting cycles).

26  
27 969 ● Biochar can reduce pesticide bioavailability and, consequently, its biodegradation.  
28  
29 970 Nonetheless, the microbiological processes still need investigation.

30  
31 971 ● Ecological risk assessment is still a gap regarding research on biochar applications.  
32  
33 972 Thus, bioassays are encouraged since several bioindicators are sensitive to pesticides,  
34  
35 973 besides the potential to assess desorption.

36  
37 974 ● Characterizing biochar bulk and flow properties is fundamental for designing solids  
38  
39 975 feeders, transport, and storage equipment. In addition, pelletized biochar's flow,  
40  
41 976 mechanical, and structural properties should be assessed to achieve cost-effective  
42  
43 977 dispersion of this material in farm soils.

44  
45 978 ● No studies showed an economic analysis of biochar applications. Therefore, performing  
46  
47 979 process techno-economic analysis under different farm contexts is encouraged.  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 980 ● Restoration methods must be feasible, technically achievable, and socially acceptable.

981 In addition, governments should provide financial incentives for environmental  
982 remediation in rural areas.

983

## 984 **Disclosure Statement**

985 No potential conflict of interest was reported by the authors.

986

## 987 **Author Contributions**

988 Conceptualization: A.P.O., J.Z.L., J.P.M., L.M.S.; Methodology: A.P.O., J.Z.L., J.P.M.;

989 Formal analysis: A.P.O., J.Z.L., J.P.M., L.M.S.; Investigation: A.P.O., J.Z.L., J.P.M., L.M.S.;

990 Data curation: A.P.O., J.Z.L., J.P.M.; Writing - Original Draft: A.P.O., J.Z.L., J.P.M., L.M.S.;

991 Writing – Review & Editing: V.G.S.R., E.L.G.E.; Visualization: A.P.O., J.Z.L., J.P.M.,

992 L.M.S.; Supervision: A.P.O.; Project administration: A.P.O.; Funding Acquisition: V.G.S.R.,

993 E.L.G.E.

994

## 995 **Funding**

996 This work was supported by the São Paulo Research Foundation (FAPESP) Process

997 2015/18790-3; A.P.O and J.P.M. have a Ph.D. scholarship from the National Council for

998 Scientific and Technological Development (CNPq) Processes 140411/2018-6 and

999 140024/2019-5; J.Z.L. has a Ph.D. scholarship from FAPESP Process 2017/16961-0.

## 1000 **References**

1001 Abbas, Z., Ali, S., Rizwan, M., Zaheer, I.E., Malik, A., Riaz, M.A., Shahid, M.R., Rehman,

- 1002 M.Z. ur, Al-Wabel, M.I., 2018. A critical review of mechanisms involved in the  
1  
2 1003 adsorption of organic and inorganic contaminants through biochar. Arab. J. Geosci. 11,  
3  
4  
5 1004 1–23. <https://doi.org/10.1007/s12517-018-3790-1>  
6  
7  
8 1005 Agbagla-Dohnani, A., Nozière, P., Gaillard-Martinie, B., Puard, M., Doreau, M., 2003. Effect  
9  
10 1006 of silica content on rice straw ruminal degradation. J. Agric. Sci. 140, 183–192.  
11  
12  
13 1007 <https://doi.org/10.1017/S0021859603003034>  
14  
15  
16 1008 Ahmad, M., Rajapaksha, A.U., Lim, J.E., Zhang, M., Bolan, N., Mohan, D., Vithanage, M.,  
17  
18 1009 Lee, S.S., Ok, Y.S., 2014. Biochar as a sorbent for contaminant management in soil and  
19  
20  
21 1010 water: A review. Chemosphere. <https://doi.org/10.1016/j.chemosphere.2013.10.071>  
22  
23  
24 1011 Ahmad, R., Kookana, R.S., Megharaj, M., Alston, A.M., 2004. Aging reduces the  
25  
26 1012 bioavailability of even a weakly sorbed pesticide (carbaryl) in soil. Environ. Toxicol.  
27  
28  
29 1013 Chem. 23, 2084. <https://doi.org/10.1897/03-569>  
30  
31  
32 1014 Aldana, G.O., Hazlerigg, C., Lopez-Capel, E., Werner, D., 2020. Agrochemical leaching  
33  
34 1015 reduction in biochar-amended tropical soils of Belize. Eur. J. Soil Sci.  
35  
36  
37 1016 <https://doi.org/10.1111/ejss.13021>  
38  
39  
40 1017 Ali, N., Khan, S., Yao, H., Wang, J., 2019. Biochars reduced the bioaccessibility and  
41  
42 1018 (bio)uptake of organochlorine pesticides and changed the microbial community  
43  
44  
45 1019 dynamics in agricultural soils. Chemosphere 224, 805–815.  
46  
47 1020 <https://doi.org/10.1016/j.chemosphere.2019.02.163>  
48  
49  
50 1021 Barber, J.L., Sweetman, A.J., Van Wijk, D., Jones, K.C., 2005. Hexachlorobenzene in the  
51  
52  
53 1022 global environment: Emissions, levels, distribution, trends and processes. Sci. Total  
54  
55 1023 Environ. <https://doi.org/10.1016/j.scitotenv.2005.03.014>  
56  
57  
58 1024 Bazargan, A., Rough, S.L., McKay, G., 2014. Compaction of palm kernel shell biochars for  
59  
60  
61  
62  
63  
64  
65

- 1025 application as solid fuel. *Biomass and Bioenergy* 70, 489–497.  
1  
2 1026 <https://doi.org/10.1016/j.biombioe.2014.08.015>  
3  
4  
5 1027 Beesley, L., Moreno-Jiménez, E., Gomez-Eyles, J.L., Harris, E., Robinson, B., Sizmur, T.,  
6  
7  
8 1028 2011. A review of biochars' potential role in the remediation, revegetation and  
9  
10 1029 restoration of contaminated soils. *Environ. Pollut.*  
11  
12 <https://doi.org/10.1016/j.envpol.2011.07.023>  
13 1030  
14  
15  
16 1031 Belden, J.B., Brain, R.A., 2018. Incorporating the joint toxicity of co-applied pesticides into  
17  
18 1032 the ecological risk assessment process. *Integr. Environ. Assess. Manag.* 14, 79–91.  
19  
20  
21 1033 <https://doi.org/10.1002/ieam.1957>  
22  
23  
24 1034 Bielská, L., Škulcová, L., Neuwirthová, N., Cornelissen, G., Hale, S.E., 2018. Sorption,  
25  
26 1035 bioavailability and ecotoxic effects of hydrophobic organic compounds in biochar  
27  
28  
29 1036 amended soils. *Sci. Total Environ.* 624, 78–86.  
30  
31 1037 <https://doi.org/10.1016/j.scitotenv.2017.12.098>  
32  
33  
34 1038 Biswas, B., Pandey, N., Bisht, Y., Singh, R., Kumar, J., Bhaskar, T., 2017. Pyrolysis of  
35  
36  
37 1039 agricultural biomass residues: Comparative study of corn cob, wheat straw, rice straw  
38  
39 1040 and rice husk. *Bioresour. Technol.* 237, 57–63.  
40  
41  
42 1041 <https://doi.org/10.1016/j.biortech.2017.02.046>  
43  
44  
45 1042 Brovini, E.M., de Deus, B.C.T., Vilas-Boas, J.A., Quadra, G.R., Carvalho, L., Mendonça,  
46  
47 1043 R.F., Pereira, R. de O., Cardoso, S.J., 2021. Three-bestseller pesticides in Brazil:  
48  
49  
50 1044 Freshwater concentrations and potential environmental risks. *Sci. Total Environ.* 771,  
51  
52 1045 144754. <https://doi.org/10.1016/j.scitotenv.2020.144754>  
53  
54  
55 1046 Chen, P., Yuan, Z., Shen, X., Zhang, Y., 2012. Flow properties of three fuel powders.  
56  
57  
58 1047 *Particuology* 10, 438–443. <https://doi.org/10.1016/j.partic.2011.11.013>  
59  
60  
61  
62  
63  
64  
65



- 1048 Cheng, H., Jones, D.L., Hill, P., Bastami, M.S., 2017. Biochar concomitantly increases  
1  
2 1049 simazine sorption in sandy loam soil and lowers its dissipation. *Arch. Agron. Soil Sci.*  
3  
4  
5 1050 63, 1082–1092. <https://doi.org/10.1080/03650340.2016.1261117>  
6  
7  
8 1051 Chinnayya, A., Chtab, A., Shao, J.Q., Carter, R.M., Yan, Y., Caillat, S., 2009.  
9  
10 1052 Characterization of pneumatic transportation of pulverised coal in a horizontal pipeline  
11  
12  
13 1053 through measurement and computational modelling. *Fuel* 88, 2348–2356.  
14  
15 1054 <https://doi.org/10.1016/j.fuel.2009.05.010>  
16  
17  
18 1055 Clay, S.A., Krack, K.K., Bruggeman, S.A., Papiernik, S., Schumacher, T.E., 2016. Maize,  
19  
20  
21 1056 switchgrass, and ponderosa pine biochar added to soil increased herbicide sorption and  
22  
23 1057 decreased herbicide efficacy. *J. Environ. Sci. Heal. Part B* 51, 497–507.  
24  
25  
26 1058 <https://doi.org/10.1080/03601234.2016.1170540>  
27  
28  
29 1059 Cong, X., Guo, X., Gong, X., Lu, H., Dong, W., 2011. Experimental research of flow patterns  
30  
31 1060 and pressure signals in horizontal dense phase pneumatic conveying of pulverized coal.  
32  
33  
34 1061 *Powder Technol.* 208, 600–609. <https://doi.org/10.1016/j.powtec.2010.12.027>  
35  
36  
37 1062 Dai, Y., Zhang, N., Xing, C., Cui, Q., Sun, Q., 2019. The adsorption, regeneration and  
38  
39 1063 engineering applications of biochar for removal organic pollutants: A review.  
40  
41  
42 1064 *Chemosphere.* <https://doi.org/10.1016/j.chemosphere.2019.01.161>  
43  
44  
45 1065 Damalas, C.A., Eleftherohorinos, I.G., 2011. Pesticide exposure, safety issues, and risk  
46  
47 1066 assessment indicators. *Int. J. Environ. Res. Public Health.*  
48  
49  
50 1067 <https://doi.org/10.3390/ijerph8051402>  
51  
52  
53 1068 Das, S.K., 2013. Mode of action of pesticides and the novel trends-A critical review. *Int. Res.*  
54  
55 1069 *J. Agric. Sci. Soil Sci.* 3, 393–401. <https://doi.org/10.14303/irjas.2013.118>  
56  
57  
58 1070 Das, S.K., Mukherjee, I., 2020. Low Cost Biomass Derived Biochar Amendment on  
59  
60  
61  
62  
63  
64  
65

- 1071 Persistence and Sorption Behaviour of Flubendiamide in Soil. *Bull. Environ. Contam.*  
1  
2 1072 *Toxicol.* 105, 261–269. <https://doi.org/10.1007/s00128-020-02936-4>  
3  
4  
5 1073 de Figueirêdo, L.P., Athayde, D.B., Daam, M.A., da Silva Guerra, G., Duarte-Neto, P.J.,  
6  
7  
8 1074 Sarmiento, H., Espíndola, E.L.G., 2020. Integrated ecosystem models (soil-water) to  
9  
10 1075 analyze pesticide toxicity to aquatic organisms at two different temperature conditions.  
11  
12  
13 1076 *Chemosphere* 270, 129422. <https://doi.org/10.1016/j.chemosphere.2020.129422>  
14  
15  
16 1077 Deng, H., Feng, D., He, J. xiong, Li, F. ze, Yu, H. mei, Ge, C. jun, 2017. Influence of biochar  
17  
18 1078 amendments to soil on the mobility of atrazine using sorption-desorption and soil thin-  
19  
20  
21 1079 layer chromatography. *Ecol. Eng.* 99, 381–390.  
22  
23 1080 <https://doi.org/10.1016/j.ecoleng.2016.11.021>  
24  
25  
26 1081 Diatta, A.A., Fike, J.H., Battaglia, M.L., Galbraith, J.M., Baig, M.B., 2020. Effects of biochar  
27  
28  
29 1082 on soil fertility and crop productivity in arid regions: a review. *Arab. J. Geosci.* 13, 1–  
30  
31 1083 17. <https://doi.org/10.1007/s12517-020-05586-2>  
32  
33  
34 1084 Ding, Y., Liu, Y., Liu, S., Huang, X., Li, Z., Tan, X., Zeng, G., Zhou, L., 2017. Potential  
35  
36  
37 1085 Benefits of Biochar in Agricultural Soils: A Review. *Pedosphere.*  
38  
39 1086 [https://doi.org/10.1016/S1002-0160\(17\)60375-8](https://doi.org/10.1016/S1002-0160(17)60375-8)  
40  
41  
42 1087 Ding, Y., Liu, Y., Liu, S., Li, Z., Tan, X., Huang, X., Zeng, G., Zhou, L., Zheng, B., 2016.  
43  
44  
45 1088 Biochar to improve soil fertility. A review. *Agron. Sustain. Dev.*  
46  
47 1089 <https://doi.org/10.1007/s13593-016-0372-z>  
48  
49  
50 1090 Duque-Acevedo, M., Belmonte-Ureña, L.J., Cortés-García, F.J., Camacho-Ferre, F., 2020.  
51  
52  
53 1091 Agricultural waste: Review of the evolution, approaches and perspectives on alternative  
54  
55 1092 uses. *Glob. Ecol. Conserv.* <https://doi.org/10.1016/j.gecco.2020.e00902>  
56  
57  
58 1093 Enstad, G., 1975. On the theory of arching in mass flow hoppers. *Chem. Eng. Sci.* 30, 1273–  
59  
60  
61  
62  
63  
64  
65

- 1094 1283. [https://doi.org/10.1016/0009-2509\(75\)85051-2](https://doi.org/10.1016/0009-2509(75)85051-2)  
1  
2
- 3 1095 Erickson, M.J., Dobbins, C., Tyner, W.E., 2011. The Economics of Harvesting Corn Cobs for  
4  
5 1096 Energy. *Crop Manag.* 10, 1–8. <https://doi.org/10.1094/CM-2011-0324-02-RS>  
6  
7
- 8 1097 FAO, 2020. Food and Agriculture Organization of the United Nations. FAOSTAT, Agri-  
9  
10 environmental Indicators / Pesticides.  
11 1098  
12  
13
- 14 1099 FAO, 2019. Food and Agriculture Organization of the United Nations. FAOSTAT, Crop  
15  
16 1100 statistics.  
17 1100  
18
- 19 1101 FAO, 2015. Food and Agriculture Organization of the United Nations. Prospects by Major  
20  
21 Sector Crop Production.  
22 1102  
23  
24
- 25 1103 Fernández Rodríguez, M.D., García Gómez, M.C., Alonso Blazquez, N., Tarazona, J.V.,  
26  
27 1104 2014. Soil Pollution Remediation. *Encycl. Toxicol.* 344–355.  
28  
29  
30 1105 <https://doi.org/10.1016/B978-0-12-386454-3.00579-0>  
31  
32
- 33 1106 Gan, J., Zhu, Y., Wilen, C., Pittenger, D., Crowley, D., 2003. Effect of Planting Covers on  
34  
35 1107 Herbicide Persistence in Landscape Soils. *Environ. Sci. Technol.* 37, 2775–2779.  
36  
37  
38 1108 <https://doi.org/10.1021/es026259u>  
39  
40
- 41 1109 Ganie, A.S., Bano, S., Khan, N., Sultana, S., Rehman, Z., Rahman, M.M., Sabir, S., Coulon,  
42  
43 F., Khan, M.Z., 2021. Nanoremediation technologies for sustainable remediation of  
44 1110  
45 contaminated environments: Recent advances and challenges. *Chemosphere.*  
46 1111  
47  
48 1112 <https://doi.org/10.1016/j.chemosphere.2021.130065>  
49  
50
- 51 1113 Gao, Y., Jiang, Z., Li, J., Xie, W., Jiang, Q., Bi, M., Zhang, Y., 2019. A comparison of the  
52  
53 characteristics and atrazine adsorption capacity of co-pyrolysed and mixed biochars  
54 1114  
55 generated from corn straw and sawdust. *Environ. Res.* 172, 561–568.  
56 1115  
57  
58 1116 <https://doi.org/10.1016/j.envres.2019.03.010>  
59  
60  
61  
62  
63  
64  
65

- 1117 Geldart, D., Jones, P., 1991. The behaviour of L-valves with granular powders. Powder  
1  
2 1118 Technol. 67, 163–174. [https://doi.org/10.1016/0032-5910\(91\)80153-A](https://doi.org/10.1016/0032-5910(91)80153-A)  
3  
4  
5 1119 Graber, E.R., Tsechansky, L., Khanukov, J., Oka, Y., 2011. Sorption, Volatilization, and  
6  
7  
8 1120 Efficacy of the Fumigant 1,3-Dichloropropene in a Biochar-Amended Soil. Soil Sci.  
9  
10 1121 Soc. Am. J. 75, 1365–1373. <https://doi.org/10.2136/sssaj2010.0435>  
11  
12  
13 1122 Guo, Z., Chen, X., Xu, Y., Liu, H., 2015. Study of flow characteristics of biomass and  
14  
15  
16 1123 biomass-coal blends. Fuel 141, 207–213. <https://doi.org/10.1016/j.fuel.2014.10.062>  
17  
18  
19 1124 Hall, K.E., Spokas, K.A., Gamiz, B., Cox, L., Papiernik, S.K., Koskinen, W.C., 2018.  
20  
21 1125 Glyphosate sorption/desorption on biochars - interactions of physical and chemical  
22  
23  
24 1126 processes. Pest Manag. Sci. 74, 1206–1212. <https://doi.org/10.1002/ps.4530>  
25  
26  
27 1127 Herath, I., Kumarathilaka, P., Al-Wabel, M.I., Abduljabbar, A., Ahmad, M., Usman, A.R.A.,  
28  
29 1128 Vithanage, M., 2016. Mechanistic modeling of glyphosate interaction with rice husk  
30  
31  
32 1129 derived engineered biochar. Microporous Mesoporous Mater. 225, 280–288.  
33  
34 1130 <https://doi.org/10.1016/j.micromeso.2016.01.017>  
35  
36  
37 1131 Hou, D., Al-Tabbaa, A., 2014. Sustainability: A new imperative in contaminated land  
38  
39  
40 1132 remediation. Environ. Sci. Policy 39, 25–34.  
41  
42 1133 <https://doi.org/10.1016/J.ENVSCI.2014.02.003>  
43  
44  
45 1134 Hu, Q., Yang, H., Yao, D., Zhu, D., Wang, X., Shao, J., Chen, H., 2016. The densification of  
46  
47  
48 1135 bio-char: Effect of pyrolysis temperature on the qualities of pellets. Bioresour. Technol.  
49  
50 1136 200, 521–527. <https://doi.org/10.1016/j.biortech.2015.10.077>  
51  
52  
53 1137 Huang, H., Zhang, C., Zhang, P., Cao, M., Xu, G., Wu, H., Zhang, J., Li, C., Rong, Q., 2018.  
54  
55  
56 1138 Effects of biochar amendment on the sorption and degradation of atrazine in different  
57  
58 1139 soils. Soil Sediment Contam. 27, 643–657.  
59  
60  
61  
62  
63  
64  
65

- 1140 <https://doi.org/10.1080/15320383.2018.1495691>  
1  
2
- 3 1141 Humera Aziz, Ghulam Murtaza, Muhammad Usman, Shahzad M. A. Basra, Abdullah Niaz,  
4  
5 1142 2018. Enhancing chlorpyrifos sorption potential of agricultural soil by biochar and  
6  
7  
8 1143 compost addition. *Pak. J. Agri. Sci.* 55, 833–841.  
9  
10 1144 <https://doi.org/10.21162/PAKJAS/18.7463>  
11  
12
- 13 1145 Hussain, M., Farooq, M., Nawaz, A., Al-Sadi, A.M., Solaiman, Z.M., Alghamdi, S.S.,  
14  
15  
16 1146 Ammara, U., Ok, Y.S., Siddique, K.H.M., 2017. Biochar for crop production: potential  
17  
18 1147 benefits and risks. *J. Soils Sediments* 17, 685–716. <https://doi.org/10.1007/s11368-016->  
19  
20  
21 1148 1360-2  
22  
23
- 24 1149 Jemec, A., Tišler, T., Drobne, D., Sepčić, K., Fournier, D., Trebše, P., 2007. Comparative  
25  
26 1150 toxicity of imidacloprid, of its commercial liquid formulation and of diazinon to a non-  
27  
28  
29 1151 target arthropod, the microcrustacean *Daphnia magna*. *Chemosphere* 68, 1408–1418.  
30  
31 1152 <https://doi.org/10.1016/j.chemosphere.2007.04.015>  
32  
33
- 34 1153 Jin, Y., Lu, H., Guo, X., Gong, X., 2019. Characteristics and formation mechanism of plug  
35  
36  
37 1154 flow in the industrial vertical pipeline of dense-phase pneumatic conveying of  
38  
39 1155 pulverized coal. *Chem. Eng. Sci.* 205, 319–331.  
40  
41 1156 <https://doi.org/10.1016/j.ces.2019.05.002>  
42  
43  
44
- 45 1157 Jones, A.D., Bruland, G.L., Agrawal, S.G., Vasudevan, D., 2005. Factors influencing the  
46  
47 1158 sorption of oxytetracycline to soils. *Environ. Toxicol. Chem.* 24, 761–770.  
48  
49 1159 <https://doi.org/10.1897/04-037R.1>  
50  
51  
52
- 53 1160 Junqueira, L.V., Mendes, K.F., Sousa, R.N. de, Almeida, C. de S., Alonso, F.G., Tornisielo,  
54  
55 1161 V.L., 2020. Sorption-desorption isotherms and biodegradation of glyphosate in two  
56  
57  
58 1162 tropical soils aged with eucalyptus biochar. *Arch. Agron. Soil Sci.* 66, 1651–1667.  
59  
60 1163 <https://doi.org/10.1080/03650340.2019.1686139>  
61  
62  
63  
64  
65

- 1164 Kadam, K.L., McMillan, J.D., 2003. Availability of corn stover as a sustainable feedstock for  
1  
2 1165 bioethanol production. *Bioresour. Technol.* 88, 17–25. <https://doi.org/10.1016/S0960->  
3  
4 1166 8524(02)00269-9  
5  
6  
7  
8 1167 Kalman, H., 2020. Role of Reynolds and Archimedes numbers in particle-fluid flows. *Rev.*  
9  
10 1168 *Chem. Eng.* <https://doi.org/10.1515/revce-2020-0005>  
11  
12  
13 1169 Kalman, H., Rawat, A., 2020. Flow regime chart for pneumatic conveying. *Chem. Eng. Sci.*  
14  
15 211, 115256. <https://doi.org/10.1016/j.ces.2019.115256>  
16 1170  
17  
18  
19 1171 Kazemi Shariat Panahi, H., Dehghani, M., Ok, Y.S., Nizami, A.S., Khoshnevisan, B.,  
20  
21 1172 Mussatto, S.I., Aghbashlo, M., Tabatabaei, M., Lam, S.S., 2020. A comprehensive  
22  
23 review of engineered biochar: Production, characteristics, and environmental  
24 1173 applications. *J. Clean. Prod.* <https://doi.org/10.1016/j.jclepro.2020.122462>  
25  
26 1174  
27  
28  
29 1175 Khalid, S., Shahid, M., Murtaza, B., Bibi, I., Natasha, Asif Naeem, M., Niazi, N.K., 2020. A  
30  
31 critical review of different factors governing the fate of pesticides in soil under biochar  
32 1176 application. *Sci. Total Environ.* 711, 134645.  
33  
34 1177 <https://doi.org/10.1016/j.scitotenv.2019.134645>  
35  
36 1178  
37  
38  
39  
40 1179 Khorram, M.S., Lin, D., Zhang, Q., Zheng, Y., Fang, H., Yu, Y., 2017. Effects of aging  
41  
42 1180 process on adsorption–desorption and bioavailability of fomesafen in an agricultural soil  
43  
44 amended with rice hull biochar. *J. Environ. Sci. (China)* 56, 180–191.  
45 1181  
46  
47 1182 <https://doi.org/10.1016/j.jes.2016.09.012>  
48  
49  
50 1183 Khorram, M.S., Sarmah, A.K., Yu, Y., 2018. The effects of biochar properties on fomesafen  
51  
52 adsorption-desorption capacity of biochar-amended soil. *Water. Air. Soil Pollut.* 229, 1–  
53 1184 13. <https://doi.org/10.1007/s11270-017-3603-2>  
54  
55 1185  
56  
57  
58 1186 Khorram, M.S., Wang, Y., Jin, X., Fang, H., Yu, Y., 2015. Reduced mobility of fomesafen  
59  
60  
61  
62  
63  
64  
65

- 1187 through enhanced adsorption in biochar-amended soil. *Environ. Toxicol. Chem.* 34,  
1  
2 1188 1258–1266. <https://doi.org/10.1002/etc.2946>  
3  
4  
5 1189 Kim, K.H., Kabir, E., Jahan, S.A., 2017. Exposure to pesticides and the associated human  
6  
7 health effects. *Sci. Total Environ.* <https://doi.org/10.1016/j.scitotenv.2016.09.009>  
8 1190  
9  
10  
11 1191 Kim, S., Chen, J., Cheng, T., Gindulyte, A., He, J., He, S., Li, Q., Shoemaker, B.A., Thiessen,  
12  
13 P.A., Yu, B., Zaslavsky, L., Zhang, J., Bolton, E.E., 2021. PubChem in 2021: New data  
14 1192 content and improved web interfaces. *Nucleic Acids Res.* 49, D1388–D1395.  
15  
16 1193 <https://doi.org/10.1093/nar/gkaa971>  
17  
18 1194  
19  
20  
21 1195 Kim, S., Dale, B.E., 2004. Global potential bioethanol production from wasted crops and  
22  
23 crop residues. *Biomass and Bioenergy* 26, 361–375.  
24 1196  
25  
26 1197 <https://doi.org/10.1016/j.biombioe.2003.08.002>  
27  
28  
29 1198 Kwon, G., Bhatnagar, A., Wang, H., Kwon, E.E., Song, H., 2020. A review of recent  
30  
31 advancements in utilization of biomass and industrial wastes into engineered biochar. *J.*  
32 1199  
33  
34 1200 *Hazard. Mater.* 400, 123242. <https://doi.org/10.1016/j.jhazmat.2020.123242>  
35  
36  
37 1201 Lal, R., 2008. Crop residues as soil amendments and feedstock for bioethanol production.  
38  
39  
40 1202 *Waste Manag.* 28, 747–758. <https://doi.org/10.1016/j.wasman.2007.09.023>  
41  
42  
43 1203 Lal, R., 2005. World crop residues production and implications of its use as a biofuel.  
44  
45 1204 *Environ. Int.* 31, 575–584. <https://doi.org/10.1016/j.envint.2004.09.005>  
46  
47  
48 1205 Lammoglia, S.K., Brun, F., Quemar, T., Moeys, J., Barriuso, E., Gabrielle, B., Mamy, L.,  
49  
50  
51 1206 2018. Modelling pesticides leaching in cropping systems: Effect of uncertainties in  
52  
53 1207 climate, agricultural practices, soil and pesticide properties. *Environ. Model. Softw.* 109,  
54  
55  
56 1208 342–352. <https://doi.org/10.1016/j.envsoft.2018.08.007>  
57  
58  
59 1209 Lehmann, J., Joseph, S., 2015. Biochar for environmental management: science, technology  
60  
61  
62  
63  
64  
65

- 1210 and implementation, 2nd ed. Routledge, London. <https://doi.org/0.4324/9780203762264>  
1  
2
- 3 1211 Levine, S.L., Borgert, C.J., 2018. Review and recommendations on criteria to evaluate the  
4  
5 1212 relevance of pesticide interaction data for ecological risk assessments. *Chemosphere*  
6  
7  
8 1213 209, 124–136. <https://doi.org/10.1016/J.CHEMOSPHERE.2018.06.081>  
9  
10
- 11 1214 Li, S., Harris, S., Anandhi, A., Chen, G., 2019. Predicting biochar properties and functions  
12  
13 1215 based on feedstock and pyrolysis temperature: A review and data syntheses. *J. Clean.*  
14  
15  
16 1216 *Prod.* 215, 890–902. <https://doi.org/10.1016/j.jclepro.2019.01.106>  
17  
18
- 19 1217 Li, X., Luo, J., Deng, H., Huang, P., Ge, C., Yu, H., Xu, W., 2018. Effect of cassava waste  
20  
21 1218 biochar on sorption and release behavior of atrazine in soil. *Sci. Total Environ.* 644,  
22  
23  
24 1219 1617–1624. <https://doi.org/10.1016/j.scitotenv.2018.07.239>  
25  
26
- 27 1220 Li, Z., A. Jennings, A., 2017. Implied Maximum Dose Analysis of Standard Values of 25  
28  
29 1221 Pesticides Based on Major Human Exposure Pathways. *AIMS Public Heal.* 4, 383–398.  
30  
31  
32 1222 <https://doi.org/10.3934/publichealth.2017.4.383>  
33  
34
- 35 1223 Liu, N., Charrua, A.B., Weng, C.H., Yuan, X., Ding, F., 2015. Characterization of biochars  
36  
37 1224 derived from agriculture wastes and their adsorptive removal of atrazine from aqueous  
38  
39  
40 1225 solution: A comparative study. *Bioresour. Technol.* 198, 55–62.  
41  
42 1226 <https://doi.org/10.1016/j.biortech.2015.08.129>  
43  
44
- 45 1227 Liu, W., Zheng, W., Ma, Y., Liu, K., 2006. Sorption and Degradation of Imidacloprid in Soil  
46  
47 1228 and Water. *J. Environ. Sci. Heal. Part B Pestic. Food Contam. Agric. Wastes* 41, 623–  
48  
49  
50 1229 634. <https://doi.org/10.1080/03601230600701775>  
51  
52
- 53 1230 Liu, Y., Guo, X., Lu, H., Gong, X., 2015. An investigation of the effect of particle size on the  
54  
55  
56 1231 flow behavior of pulverized coal, in: *Procedia Engineering*. Elsevier Ltd, pp. 698–713.  
57  
58 1232 <https://doi.org/10.1016/j.proeng.2015.01.170>  
59  
60  
61  
62  
63  
64  
65



- 1233 Liu, Y., Lonappan, L., Brar, S.K., Yang, S., 2018. Impact of biochar amendment in  
1  
2 1234 agricultural soils on the sorption, desorption, and degradation of pesticides: A review.  
3  
4  
5 1235 Sci. Total Environ. <https://doi.org/10.1016/j.scitotenv.2018.07.099>  
6  
7  
8 1236 Lu, H., Guo, X., Gong, X., Barletta, D., Poletto, M., 2015. Prediction of solid discharge rates  
9  
10 1237 of pulverized coal from an aerated hopper. Powder Technol. 286, 645–653.  
11  
12  
13 1238 <https://doi.org/10.1016/j.powtec.2015.09.017>  
14  
15  
16 1239 Lu, H., Guo, X., Gong, X., Cong, X., Liu, K., Qi, H., 2012. Experimental study on aerated  
17  
18 1240 discharge of pulverized coal. Chem. Eng. Sci. 71, 438–448.  
19  
20  
21 1241 <https://doi.org/10.1016/j.ces.2011.11.006>  
22  
23  
24 1242 Lu, H., Guo, X., Huang, W., Liu, K., Gong, X., 2011. Flow characteristics and pressure drop  
25  
26 1243 across the Laval nozzle in dense phase pneumatic conveying of the pulverized coal.  
27  
28  
29 1244 Chem. Eng. Process. Process Intensif. 50, 702–708.  
30  
31 1245 <https://doi.org/10.1016/j.cep.2011.03.009>  
32  
33  
34 1246 Lü, J., Li, J., Li, Y., Chen, B., Bao, Z., 2012. Use of rice straw biochar simultaneously as the  
35  
36 1247 sustained release carrier of herbicides and soil amendment for their reduced leaching. J.  
37  
38  
39 1248 Agric. Food Chem. 60, 6463–6470. <https://doi.org/10.1021/jf3009734>  
40  
41  
42 1249 Lunagariya, D.D., Patel, K.G., Singh, S., Parekh, V.B., Ahlawat, T.R., 2020. A Review on  
43  
44 1250 Adsorption and Desorption of Different Pesticides in Various Soil. Int. Res. J. Pure  
45  
46 1251 Appl. Chem. 35–41. <https://doi.org/10.9734/irjpac/2020/v21i2430332>  
47  
48  
49  
50 1252 Manna, S., Singh, N., 2019. Biochars mediated degradation, leaching and bioavailability of  
51  
52 1253 pyrazosulfuron-ethyl in a sandy loam soil. Geoderma 334, 63–71.  
53  
54  
55 1254 <https://doi.org/10.1016/j.geoderma.2018.07.032>  
56  
57  
58 1255 Manna, S., Singh, N., 2015. Effect of wheat and rice straw biochars on pyrazosulfuron-ethyl  
59  
60  
61  
62  
63  
64  
65

- 1256 sorption and persistence in a sandy loam soil. *J. Environ. Sci. Heal. Part B* 50, 463–472.  
1  
2 1257 <https://doi.org/10.1080/03601234.2015.1018757>  
3  
4  
5 1258 Massaro Sousa, L., Ferreira, M.C., Hou, Q.F., Yu, A.B., 2020. Feeding spent coffee ground  
6  
7  
8 1259 powders with a non-mechanical L-valve: Experimental analysis and TFM simulation.  
9  
10 1260 *Powder Technol.* 360, 1055–1066. <https://doi.org/10.1016/j.powtec.2019.11.005>  
11  
12  
13 1261 Massaro Sousa, L., Schulz, C.G., Condotta, R., Ferreira, M.C., 2021. On the design of conical  
14  
15  
16 1262 hoppers for spent coffee grounds: Moisture content and particle-size effects. *J. Food*  
17  
18 1263 *Eng.* 300, 110537. <https://doi.org/10.1016/j.jfoodeng.2021.110537>  
19  
20  
21 1264 Meena, R.S., Kumar, S., Datta, R., Lal, R., Vijayakumar, V., Brtnicky, M., Sharma, M.P.,  
22  
23  
24 1265 Yadav, G.S., Jhariya, M.K., Jangir, C.K., Pathan, S.I., Dokulilova, T., Pecina, V., Marfo,  
25  
26 1266 T.D., 2020. Impact of agrochemicals on soil microbiota and management: A review.  
27  
28  
29 1267 *Land.* <https://doi.org/10.3390/land9020034>  
30  
31  
32 1268 Moher, D., Liberati, A., Tetzlaff, J., Altman, D.G., 2009. Preferred Reporting Items for  
33  
34 1269 Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLoS Med.* 6,  
35  
36  
37 1270 e1000097. <https://doi.org/10.1371/journal.pmed.1000097>  
38  
39  
40 1271 Morillo, E., Villaverde, J., 2017. Advanced technologies for the remediation of pesticide-  
41  
42 1272 contaminated soils. *Sci. Total Environ.* <https://doi.org/10.1016/j.scitotenv.2017.02.020>  
43  
44  
45 1273 Nguyen, T.L.T., Hermansen, J.E., Mogensen, L., 2013. Environmental performance of crop  
46  
47  
48 1274 residues as an energy source for electricity production: The case of wheat straw in  
49  
50  
51 1275 Denmark. *Appl. Energy* 104, 633–641. <https://doi.org/10.1016/j.apenergy.2012.11.057>  
52  
53  
54 1276 Oginni, O., Fasina, O., 2018. Theoretical estimation of silo design parameters for fractionated  
55  
56 1277 loblolly pine grinds – Moisture content and particle size effects. *Ind. Crops Prod.* 123,  
57  
58 1278 379–385. <https://doi.org/10.1016/j.indcrop.2018.07.005>  
59  
60  
61  
62  
63  
64  
65

- 1279 Ouyang, W., Huang, W., Wei, P., Hao, F., Yu, Y., 2016a. Optimization of typical diffuse  
1  
2 1280 herbicide pollution control by soil amendment configurations under four levels of  
3  
4  
5 1281 rainfall intensities. *J. Environ. Manage.* 175, 1–8.  
6  
7 1282 <https://doi.org/10.1016/j.jenvman.2016.03.026>  
8  
9
- 10 1283 Ouyang, W., Zhao, X., Tysklind, M., Hao, F., 2016b. Typical agricultural diffuse herbicide  
11  
12  
13 1284 sorption with agricultural waste-derived biochars amended soil of high organic matter  
14  
15 1285 content. *Water Res.* 92, 156–163. <https://doi.org/10.1016/j.watres.2016.01.055>  
16  
17
- 18 1286 Palansooriya, K.N., Ok, Y.S., Awad, Y.M., Lee, S.S., Sung, J.K., Koutsospyros, A., Moon,  
19  
20  
21 1287 D.H., 2019. Impacts of biochar application on upland agriculture: A review. *J. Environ.*  
22  
23 1288 *Manage.* 234, 52–64. <https://doi.org/10.1016/j.jenvman.2018.12.085>  
24  
25
- 26 1289 Qin, X., Liu, Yetong, Huang, Q., Liu, Yiyun, Zhao, L., Xu, Y., 2019. In-Situ Remediation of  
27  
28  
29 1290 Cadmium and Atrazine Contaminated Acid Red Soil of South China Using Sepiolite and  
30  
31 1291 Biochar. *Bull. Environ. Contam. Toxicol.* 102, 128–133. <https://doi.org/10.1007/s00128->  
32  
33 1292 [018-2494-2](https://doi.org/10.1007/s00128-018-2494-2)  
34  
35  
36
- 37 1293 Racke, K.D., 1993. Environmental fate of chlorpyrifos. *Rev. Environ. Contam. Toxicol.*  
38  
39 1294 [https://doi.org/10.1007/978-1-4612-4362-5\\_1](https://doi.org/10.1007/978-1-4612-4362-5_1)  
40  
41
- 42 1295 Rani, L., Thapa, K., Kanojia, N., Sharma, N., Singh, S., Grewal, A.S., Srivastav, A.L.,  
43  
44  
45 1296 Kaushal, J., 2021. An extensive review on the consequences of chemical pesticides on  
46  
47 1297 human health and environment. *J. Clean. Prod.*  
48  
49 1298 <https://doi.org/10.1016/j.jclepro.2020.124657>  
50  
51  
52
- 53 1299 Ranzi, E., Costa, M., Bartocci, P., Barbanera, M., Skreiberg, Ø., Wang, L., Bidini, G.,  
54  
55 1300 Fantozzi, F., 2018. Biocarbon Pellet Production: Optimization of Pelletizing Process, in:  
56  
57 1301 *CHEMICAL ENGINEERING TRANSACTIONS.* <https://doi.org/10.3303/CET1865060>  
58  
59  
60  
61  
62  
63  
64  
65

- 1302 Regkouzas, P., Diamadopoulou, E., 2019. Adsorption of selected organic micro-pollutants on  
1  
2 1303 sewage sludge biochar. *Chemosphere* 224, 840–851.  
3  
4  
5 1304 <https://doi.org/10.1016/j.chemosphere.2019.02.165>  
6  
7  
8 1305 Ren, X., Sun, H., Wang, F., Zhang, P., Zhu, H., 2018. Effect of aging in field soil on  
9  
10 1306 biochar's properties and its sorption capacity. *Environ. Pollut.* 242, 1880–1886.  
11  
12  
13 1307 <https://doi.org/10.1016/j.envpol.2018.07.078>  
14  
15  
16 1308 Ren, X., Zhang, P., Zhao, L., Sun, H., 2016. Sorption and degradation of carbaryl in soils  
17  
18 1309 amended with biochars: influence of biochar type and content. *Environ. Sci. Pollut. Res.*  
19  
20  
21 1310 23, 2724–2734. <https://doi.org/10.1007/s11356-015-5518-z>  
22  
23  
24 1311 Rey Benayas, J.M., Newton, A.C., Diaz, A., Bullock, J.M., 2009. Enhancement of  
25  
26 1312 biodiversity and ecosystem services by ecological restoration: A meta-analysis. *Science*  
27  
28  
29 1313 (80-. ). 325, 1121–1124. <https://doi.org/10.1126/science.1172460>  
30  
31  
32 1314 Rouzies, E., Lauvernet, C., Barachet, C., Morel, T., Branger, F., Braud, I., Carlier, N., 2019.  
33  
34 1315 From agricultural catchment to management scenarios: A modular tool to assess effects  
35  
36  
37 1316 of landscape features on water and pesticide behavior. *Sci. Total Environ.* 671, 1144–  
38  
39 1317 1160. <https://doi.org/10.1016/j.scitotenv.2019.03.060>  
40  
41  
42 1318 Sadegh-Zadeh, F., Abd Wahid, S., Jalili, B., 2017. *Advances in Environmental Technology*  
43  
44  
45 1319 Sorption, degradation and leaching of pesticides in soils amended with organic matter: A  
46  
47 1320 review. *Adv. Environ. Technol.* 2, 119–132.  
48  
49  
50 1321 <https://doi.org/10.22104/AET.2017.1740.1100>  
51  
52  
53 1322 Saha, A., Bhaduri, D., Pipariya, A., Kumar Ghosh, R., 2017. Linear and nonlinear sorption  
54  
55 1323 modelling for adsorption of atrazine onto activated peanut husk. *Environ. Prog. Sustain.*  
56  
57  
58 1324 *Energy* 36, 348–358. <https://doi.org/10.1002/ep.12434>  
59  
60  
61  
62  
63  
64  
65

- 1325 Saleh, I.A., Zouari, N., Al-Ghouti, M.A., 2020. Removal of pesticides from water and  
1  
2 1326 wastewater: Chemical, physical and biological treatment approaches. *Environ. Technol.*  
3  
4  
5 1327 *Innov.* <https://doi.org/10.1016/j.eti.2020.101026>  
6  
7  
8 1328 Sanches, A.L.M., Vieira, B.H., Reghini, M.V., Moreira, R.A., Freitas, E.C., Espíndola,  
9  
10 1329 E.L.G., Daam, M.A., 2017. Single and mixture toxicity of abamectin and difenoconazole  
11  
12  
13 1330 to adult zebrafish (*Danio rerio*). *Chemosphere* 188, 582–587.  
14  
15 1331 <https://doi.org/10.1016/J.CHEMOSPHERE.2017.09.027>  
16  
17  
18 1332 Santos, B.M., Gilreath, J.P., Motis, T.N., Noling, J.W., Jones, J.P., Norton, J.A., 2006.  
19  
20  
21 1333 Comparing methyl bromide alternatives for soilborne disease, nematode and weed  
22  
23 1334 management in fresh market tomato. *Crop Prot.* 25, 690–695.  
24  
25 1335 <https://doi.org/10.1016/j.cropro.2005.09.015>  
26  
27  
28  
29 1336 Schäfer, R.B., Bundschuh, M., Rouch, D.A., Szöcs, E., von der Ohe, P.C., Pettigrove, V.,  
30  
31 1337 Schulz, R., Nugegoda, D., Kefford, B.J., 2012. Effects of pesticide toxicity, salinity and  
32  
33  
34 1338 other environmental variables on selected ecosystem functions in streams and the  
35  
36 1339 relevance for ecosystem services. *Sci. Total Environ.* 415, 69–78.  
37  
38 1340 <https://doi.org/10.1016/j.scitotenv.2011.05.063>  
39  
40  
41  
42 1341 Schiesari, L., Grillitsch, B., 2011. Pesticides meet megadiversity in the expansion of biofuel  
43  
44 1342 crops. *Front. Ecol. Environ.* 9, 215–221. <https://doi.org/10.1890/090139>  
45  
46  
47 1343 Scow, K.M., Johnson, C.R., 1996. Effect of Sorption on Biodegradation of Soil Pollutants.  
48  
49 1344 *Adv. Agron.* 58, 1–56. [https://doi.org/10.1016/S0065-2113\(08\)60252-7](https://doi.org/10.1016/S0065-2113(08)60252-7)  
50  
51  
52  
53 1345 Sinclair, C.J., Boxall, A.B.A., 2003. Assessing the ecotoxicity of pesticide transformation  
54  
55 1346 products. *Environ. Sci. Technol.* 37, 4617–4625. <https://doi.org/10.1021/es030038m>  
56  
57  
58 1347 Singh, S., Kumar, V., Chauhan, A., Datta, S., Wani, A.B., Singh, N., Singh, J., 2018.  
59  
60  
61  
62  
63  
64  
65

- 1348 Toxicity, degradation and analysis of the herbicide atrazine. *Environ. Chem. Lett.*  
1  
2 1349 <https://doi.org/10.1007/s10311-017-0665-8>  
3  
4  
5 1350 Sistani, K.R., Reddy, K.C., Kanyika, W., Savant, N.K., 1998. Integration of rice crop residue  
6  
7 into sustainable rice production system. *J. Plant Nutr.* 21, 1855–1866.  
8 1351  
9  
10 1352 <https://doi.org/10.1080/01904169809365528>  
11  
12  
13 1353 Smolders, K., Baeyens, J., 1995. The operation of L-valves to control standpipe flow. *Adv.*  
14  
15 *Powder Technol.* 6, 163–176. [https://doi.org/10.1016/S0921-8831\(08\)60525-7](https://doi.org/10.1016/S0921-8831(08)60525-7)  
16 1354  
17  
18  
19 1355 Song, Y., Li, Y., Zhang, W., Wang, F., Bian, Y., Boughner, L.A., Jiang, X., 2016. Novel  
20  
21 1356 Biochar-Plant Tandem Approach for Remediating Hexachlorobenzene Contaminated  
22  
23 Soils: Proof-of-Concept and New Insight into the Rhizosphere. *J. Agric. Food Chem.* 64,  
24 1357  
25 5464–5471. <https://doi.org/10.1021/acs.jafc.6b01035>  
26 1358  
27  
28  
29 1359 Song, Y., Wang, F., Bian, Y., Kengara, F.O., Jia, M., Xie, Z., Jiang, X., 2012. Bioavailability  
30  
31 assessment of hexachlorobenzene in soil as affected by wheat straw biochar. *J. Hazard.*  
32 1360  
33 *Mater.* 217–218, 391–397. <https://doi.org/10.1016/j.jhazmat.2012.03.055>  
34 1361  
35  
36  
37 1362 Song, Z., Feng, X., Lal, R., Fan, M., Ren, J., Qi, H., Qian, C., Guo, J., Cai, H., Cao, T., Yu,  
38  
39 Y., Hao, Y., Huang, X., Deng, A., Zheng, C., Zhang, J., Zhang, W., 2019. Optimized  
40 1363  
41 agronomic management as a double-win option for higher maize productivity and less  
42 1364  
43 global warming intensity: A case study of Northeastern China, in: *Advances in*  
44 1365  
45 *Agronomy*. Academic Press Inc., pp. 251–292.  
46  
47 1366  
48  
49 1367 <https://doi.org/10.1016/bs.agron.2019.04.002>  
50  
51  
52  
53 1368 Spark, K.M., Swift, R.S., 2002. Effect of soil composition and dissolved organic matter on  
54  
55 1369 pesticide sorption. *Sci. Total Environ.* 298, 147–161. <https://doi.org/10.1016/S0048->  
56  
57 9697(02)00213-9  
58 1370  
59  
60  
61  
62  
63  
64  
65

- 1371 Spurlock, F.C., Biggar, J.W., 1994. Thermodynamics of Organic Chemical Partition in Soils.  
1  
2 1372 2. Nonlinear Partition of Substituted Phenylureas from Aqueous Solution. Environ. Sci.  
3  
4  
5 1373 Technol. 28, 996–1002. <https://doi.org/10.1021/es00055a006>  
6  
7  
8 1374 Tang, F.H.M., Lenzen, M., McBratney, A., Maggi, F., 2021. Risk of pesticide pollution at the  
9  
10 1375 global scale. Nat. Geosci. 14, 206–210. <https://doi.org/10.1038/s41561-021-00712-5>  
11  
12  
13 1376 Tao, Y., Han, S., Zhang, Q., Yang, Y., Shi, H., Akindolie, M.S., Jiao, Y., Qu, J., Jiang, Z.,  
14  
15  
16 1377 Han, W., Zhang, Y., 2020. Application of biochar with functional microorganisms for  
17  
18 1378 enhanced atrazine removal and phosphorus utilization. J. Clean. Prod. 257, 120535.  
19  
20  
21 1379 <https://doi.org/10.1016/j.jclepro.2020.120535>  
22  
23  
24 1380 Tatarková, V., Hiller, E., Vaculík, M., 2013. Impact of wheat straw biochar addition to soil  
25  
26 1381 on the sorption, leaching, dissipation of the herbicide (4-chloro-2-methylphenoxy)acetic  
27  
28  
29 1382 acid and the growth of sunflower (*Helianthus annuus* L.). Ecotoxicol. Environ. Saf. 92,  
30  
31 1383 215–221. <https://doi.org/10.1016/j.ecoenv.2013.02.005>  
32  
33  
34 1384 Toloue Farrokh, N., Suopajarvi, H., Mattila, O., Sulasalmi, P., Fabritius, T., 2020.  
35  
36  
37 1385 Characteristics of wood-based biochars for pulverized coal injection. Fuel 265, 117017.  
38  
39 1386 <https://doi.org/10.1016/j.fuel.2020.117017>  
40  
41  
42 1387 Toloue Farrokh, N., Suopajarvi, H., Mattila, O., Umeki, K., Phounglamcheik, A., Romar, H.,  
43  
44  
45 1388 Sulasalmi, P., Fabritius, T., 2018. Slow pyrolysis of by-product lignin from wood-based  
46  
47 1389 ethanol production— A detailed analysis of the produced chars. Energy 164, 112–123.  
48  
49  
50 1390 <https://doi.org/10.1016/j.energy.2018.08.161>  
51  
52  
53 1391 Triques, M.C., Oliveira, D., Veloso Goulart, B., Carolina Montagner, C., Luiz Gaeta  
54  
55 1392 Espíndola, E., Bezerra de Menezes-Oliveira, V., 2021. Assessing single effects of  
56  
57  
58 1393 sugarcane pesticides fipronil and 2,4-D on plants and soil organisms. Ecotoxicol.  
59  
60 1394 Environ. Saf. 208, 111622. <https://doi.org/10.1016/j.ecoenv.2020.111622>  
61  
62  
63  
64  
65

- 1395 USEPA, 2021. United States Environmental Protection Agency. Chemical and Products  
1  
2 1396 Database (CPDat).  
3  
4
- 5 1397 Uwamungu, J.Y., Nartey, O.D., Uwimpaye, F., Dong, W., Hu, C., 2019. Evaluating Biochar  
6  
7  
8 1398 Impact on Topramezone Adsorption Behavior on Soil under No-Tillage and Rotary  
9  
10 1399 Tillage Treatments: Isotherms and Kinetics. *Int. J. Environ. Res. Public Health* 16, 5034.  
11  
12  
13 1400 <https://doi.org/10.3390/ijerph16245034>  
14  
15
- 16 1401 Varjani, S., Kumar, G., Rene, E.R., 2019. Developments in biochar application for pesticide  
17  
18 1402 remediation: Current knowledge and future research directions. *J. Environ. Manage.*  
19  
20  
21 1403 232, 505–513. <https://doi.org/10.1016/j.jenvman.2018.11.043>  
22  
23
- 24 1404 Villamil, M.B., Little, J., Nafziger, E.D., 2015. Corn residue, tillage, and nitrogen rate effects  
25  
26 1405 on soil properties. *Soil Tillage Res.* 151, 61–66.  
27  
28  
29 1406 <https://doi.org/10.1016/j.still.2015.03.005>  
30  
31
- 32 1407 Vryzas, Z., 2018. Pesticide fate in soil-sediment-water environment in relation to  
33  
34 1408 contamination preventing actions. *Curr. Opin. Environ. Sci. Heal.*  
35  
36  
37 1409 <https://doi.org/10.1016/j.coesh.2018.03.001>  
38  
39
- 40 1410 Wang, Q., Gao, S., Wang, D., Spokas, K., Cao, A., Yan, D., 2016. Mechanisms for 1,3-  
41  
42 1411 Dichloropropene Dissipation in Biochar-Amended Soils. *J. Agric. Food Chem.* 64,  
43  
44  
45 1412 2531–2540. <https://doi.org/10.1021/acs.jafc.5b04941>  
46  
47
- 48 1413 Wang, R., Yuan, Y., Yen, H., Grieneisen, M., Arnold, J., Wang, D., Wang, C., Zhang, M.,  
49  
50 1414 2019. A review of pesticide fate and transport simulation at watershed level using  
51  
52  
53 1415 SWAT: Current status and research concerns. *Sci. Total Environ.*  
54  
55 1416 <https://doi.org/10.1016/j.scitotenv.2019.03.141>  
56  
57
- 58 1417 Waqas, M., Asam, Z., Rehan, M., Anwar, M.N., Khattak, R.A., Ismail, I.M.I., Tabatabaei,  
59  
60  
61  
62  
63  
64  
65



- 1418 M., Nizami, A.S., 2020. Development of biomass-derived biochar for agronomic and  
1  
2 1419 environmental remediation applications. *Biomass Convers. Biorefinery*.  
3  
4  
5 1420 <https://doi.org/10.1007/s13399-020-00936-2>  
6  
7
- 8 1421 Wei, D., Li, B., Huang, H., Luo, L., Zhang, J., Yang, Y., Guo, J., Tang, L., Zeng, G., Zhou,  
9  
10 1422 Y., 2018. Biochar-based functional materials in the purification of agricultural  
11  
12 1423 wastewater: Fabrication, application and future research needs. *Chemosphere*.  
13  
14  
15 1424 <https://doi.org/10.1016/j.chemosphere.2017.12.193>  
16  
17
- 18 1425 Wu, C., Liu, X., Wu, X., Dong, F., Xu, J., Zheng, Y., 2019. Sorption, degradation and  
19  
20 1426 bioavailability of oxyfluorfen in biochar-amended soils. *Sci. Total Environ.* 658, 87–94.  
21  
22  
23 1427 <https://doi.org/10.1016/j.scitotenv.2018.12.059>  
24  
25
- 26 1428 Yaashikaa, P.R., Senthil Kumar, P., Varjani, S.J., Saravanan, A., 2019. Advances in  
27  
28 1429 production and application of biochar from lignocellulosic feedstocks for remediation of  
29  
30 1430 environmental pollutants. *Bioresour. Technol.*  
31  
32  
33 1431 <https://doi.org/10.1016/j.biortech.2019.122030>  
34  
35  
36
- 37 1432 Yavari, S., Abualqumboz, M., Sapari, N., Hata-Suhaimi, H.A., Nik-Fuaad, N.Z., Yavari, S.,  
38  
39 1433 2020. Sorption of imazapic and imazapyr herbicides on chitosan-modified biochars. *Int.*  
40  
41 1434 *J. Environ. Sci. Technol.* 17, 3341–3350. <https://doi.org/10.1007/s13762-020-02629-9>  
42  
43  
44
- 45 1435 Yavari, Saba, Malakahmad, A., Sapari, N.B., Yavari, Sara, 2017a. Sorption properties  
46  
47 1436 optimization of agricultural wastes-derived biochars using response surface  
48  
49 1437 methodology. *Process Saf. Environ. Prot.* 109, 509–519.  
50  
51  
52 1438 <https://doi.org/10.1016/j.psep.2017.05.002>  
53  
54
- 55 1439 Yavari, Saba, Malakahmad, A., Sapari, N.B., Yavari, Sara, 2017b. Synthesis optimization of  
56  
57 1440 oil palm empty fruit bunch and rice husk biochars for removal of imazapic and imazapyr  
58  
59 1441 herbicides. *J. Environ. Manage.* 193, 201–210.  
60  
61  
62  
63  
64  
65

- 1442 <https://doi.org/10.1016/j.jenvman.2017.02.035>  
1  
2
- 3 1443 Yavari, Saba, Malakahmad, A., Sapari, N.B., Yavari, Sara, 2016. Sorption-desorption  
4  
5 1444 mechanisms of imazapic and imazapyr herbicides on biochars produced from  
6  
7  
8 1445 agricultural wastes. *J. Environ. Chem. Eng.* 4, 3981–3989.  
9  
10 1446 <https://doi.org/10.1016/j.jece.2016.09.003>  
11  
12
- 13 1447 You, X., Jiang, H., Zhao, M., Suo, F., Zhang, C., Zheng, H., Sun, K., Zhang, G., Li, F., Li,  
14  
15  
16 1448 Y., 2020. Biochar reduced Chinese chive (*Allium tuberosum*) uptake and dissipation of  
17  
18 1449 thiamethoxam in an agricultural soil. *J. Hazard. Mater.* 390, 121749.  
19  
20  
21 1450 <https://doi.org/10.1016/j.jhazmat.2019.121749>  
22  
23
- 24 1451 Yu, H., Zou, W., Chen, J., Chen, H., Yu, Z., Huang, J., Tang, H., Wei, X., Gao, B., 2019.  
25  
26 1452 Biochar amendment improves crop production in problem soils: A review. *J. Environ.*  
27  
28  
29 1453 *Manage.* <https://doi.org/10.1016/j.jenvman.2018.10.117>  
30  
31
- 32 1454 Zhang, G., Zhang, Q., Sun, K., Liu, X., Zheng, W., Zhao, Y., 2011. Sorption of simazine to  
33  
34 1455 corn straw biochars prepared at different pyrolytic temperatures. *Environ. Pollut.* 159,  
35  
36  
37 1456 2594–2601. <https://doi.org/10.1016/j.envpol.2011.06.012>  
38  
39
- 40 1457 Zhang, L., Yan, C., Guo, Q., Zhang, J., Ruiz-Menjivar, J., 2018. The impact of agricultural  
41  
42 1458 chemical inputs on environment: Global evidence from informetrics analysis and  
43  
44  
45 1459 visualization. *Int. J. Low-Carbon Technol.* 13, 338–352.  
46  
47 1460 <https://doi.org/10.1093/ijlct/cty039>  
48  
49
- 50 1461 Zhang, P., Min, L., Tang, J., Rafiq, M.K., Sun, H., 2020. Sorption and degradation of  
51  
52  
53 1462 imidacloprid and clothianidin in Chinese paddy soil and red soil amended with biochars.  
54  
55 1463 *Biochar* 2, 329–341. <https://doi.org/10.1007/s42773-020-00060-4>  
56  
57
- 58 1464 Zhang, P., Sun, H., Min, L., Ren, C., 2018. Biochars change the sorption and degradation of  
59  
60  
61  
62  
63  
64  
65

- 1465 thiacloprid in soil: Insights into chemical and biological mechanisms. *Environ. Pollut.*  
1  
2 1466 236, 158–167. <https://doi.org/10.1016/j.envpol.2018.01.030>  
3  
4  
5 1467 Zhao, Z., Wu, Q., Nie, T., Zhou, W., 2019. Quantitative evaluation of relationships between  
6  
7  
8 1468 adsorption and partition of atrazine in biochar-amended soils with biochar  
9  
10 1469 characteristics. *RSC Adv.* 9, 4162–4171. <https://doi.org/10.1039/C8RA08544G>  
11  
12  
13 1470 Zhou, J., Han, X., Jing, S., Liu, Y., 2020. Efficiency and stability of lump coal particles  
14  
15  
16 1471 swirling flow pneumatic conveying system. *Chem. Eng. Res. Des.* 157, 92–103.  
17  
18 1472 <https://doi.org/10.1016/j.cherd.2020.03.006>  
19  
20  
21 1473 Zhou, J., Liu, Y., Du, C., Liu, S., Li, J., 2018. Numerical study of coarse coal particle  
22  
23  
24 1474 breakage in pneumatic conveying. *Particuology* 38, 204–214.  
25  
26 1475 <https://doi.org/10.1016/j.partic.2017.07.003>  
27  
28  
29 1476 Zhu, X., Li, J., Xie, B., Feng, D., Li, Y., 2020. Accelerating effects of biochar for pyrite-  
30  
31  
32 1477 catalyzed Fenton-like oxidation of herbicide 2,4-D. *Chem. Eng. J.* 391, 123605.  
33  
34 1478 <https://doi.org/10.1016/j.cej.2019.123605>  
35  
36  
37 1479 ZoBell, D.R., Okine, E.K., Olson, K.C., Wiedmeier, R.D., Goonewardene, L.A., Stonecipher,  
38  
39  
40 1480 C., 2005. Effects of feeding wheat straw and middlings ensiled with whey on  
41  
42 1481 digestibility and growth of cattle. *Can. J. Anim. Sci.* 85, 69–74.  
43  
44  
45 1482 <https://doi.org/10.4141/A04-038>  
46  
47  
48 1483  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65